



UNIVERSITY OF ILLINOIS  
LIBRARY

Class  
580.5

Book  
FB

Volume  
1  
cop 2

J3 09 20M

**CENTRAL CIRCULATION BOOKSTACKS**

The person charging this material is responsible for its return to the library from which it was borrowed on or before the **Latest Date** stamped below.

*Theft, mutilation, and underlining of books are reasons for disciplinary action and may result in dismissal from the University.*

**TO RENEW CALL TELEPHONE CENTER, 333-8400**

**UNIVERSITY OF ILLINOIS LIBRARY AT URBANA-CHAMPAIGN**

APR 22 1992  
APR 21 1992

When renewing by phone, write new due date below previous due date.

79521 L162







RIVER BIRCHES AT SUTTON, W. VIRGINIA.  
(*Betula Nigra L.*)

FIELD COLUMBIAN MUSEUM

PUBLICATION 9.

Botanical Series.

Vol. 1, No. 2.

---

FLORA  
OF  
WEST VIRGINIA

BY

CHARLES FREDERICK MILLSPAUGH,  
Curator, Department of Botany,

AND

LAWRENCE WILLIAM NUTTALL

---



CHICAGO, U. S. A.

January, 1896.

## PUBLICATIONS OF THE MUSEUM.

---

For the convenience of scientific workers it has been deemed expedient to issue the publications of the Museum in separate series for each of the sciences represented. The following series thus far have been established: Historical, Geological, Botanical, Zoölogical, Ornithological and Anthropological.

A consecutive number has been given the entire set of publications to indicate the order of issue. Each departmental series, however, has its own volume number and individual consecutive pagination, making the literature of the science, or general subject, independent and complete for separate binding; or they may be bound in order of publication without relation to subject. Proceedings, memoirs, monographs, bulletins, and hand-books and catalogues of collections, are included within the scope of the publications.

Full lists of the publications of the Museum may be found in the Annual Report of the Director.

Publications are sent to societies and institutions of a public character that reciprocate with their own literature and to a limited number of scientists who are able to exchange.

Two publications have been issued of the Botanical series: Contribution to the Flora of Yucatan (Pub. 4); and The Flora of West Virginia (Pub. 9). Contribution II to the Flora of Yucatan is in preparation.

FREDERICK J. V. SKIFF,  
*Director.*



## CONTENTS.

---

	PAGE
Introductory, - - - - -	69
Botanical History, - - - - -	70
Special Features of the Flora, - - - - -	71
The Sylva, - - - - -	76
Summary of the Flora, - - - - -	80
Catalogue of the Species, - - - - -	81
Host Index of the Fungi, - - - - -	234
Local Plant Names, - - - - -	261
Generic Index, - - - - -	264

---

## ILLUSTRATIONS.

---

Pl. V. River Birches at Sutton, West Virginia, - - -	FRONTISPIECE
Pl. VI. Map of West Virginia, - - - - -	169
Pl. VII. <i>Rhus radicans</i> , Linn., - - - - -	215



# The Flora of West Virginia.

BY

C. F. MILLSAUGH AND L. W. NUTTALL.

The State of West Virginia comprises about 24,780 square miles of territory lying between  $37^{\circ}30'$  and  $40^{\circ}30'$  north latitude, and  $0^{\circ}45'$  and  $5^{\circ}30'$  west longitude from Washington. Its outline of varied boundaries has become a synonym for irregularity, as a glance at the accompanying map will show; it might also comprise the topography, suggesting as it does an immense field over which a gigantic plow has left a confused maze of deep and irregular furrows. This topographical condition is mainly due to the great number of mountain ranges and a vast network of rapid streams, that, rising in the higher altitudes of the eastern and southern borders, pass in varied and tortuous courses through the State, to augment the Ohio on the west and northwest, and the Potomac on the northeast.

Along the low, as well as the lofty mountain ranges, there is comparatively little tableland, and in the wedge-like valleys there is a like absence of extensive bottoms, except along Tygart's Valley River in Randolph County, the Great Kanawha and the Ohio. Although there are many mountain glades, some nearly dry and others swampy, plainly indicating their late occupancy by small lakes, there is to-day neither pond nor lake within the limits of the State, and very little if any stagnant water.

As the major portion of the State lies west of the Alleghanies, the prevailing climate is much like that of western Pennsylvania, partaking little indeed of that southern atmosphere that we are wont to associate with the name Virginia.

The prevailing soil of the hills and valleys is stiff clay, and sandy and clayey alluvium, over which there is in general but little loam. The clay of the more open steep hillsides is so unctuous and unstable that frequent landslides occur during Spring, sometimes of great extent. This subsidence renders the valley streams muddy throughout the year. The rocks are principally sandstone and limestone, with some

outcroppings of shales on the northeastern heights. The special features of the now very fertile and then quite sterile soils, with the varied differences in altitude, as well as the vast areas of primitive forests, yield a flora of great variety, and often widely differing at points only a few miles apart.

The amount of exploration necessary under such conditions to gain a full knowledge of the flora, becomes an arduous undertaking, though the interest in searching an almost virgin field is so deep as to greatly lighten the labor.

#### BOTANICAL HISTORY.

The early botanical explorers, Pursh. and Nuttall, found many of their novelties among the eastern mountains of this State, and the old Dutch gardener, Kin, here sought oddities for horticulture, but either on account of their limited knowledge regarding the geography of this section, or from the undeveloped condition of the area they traversed, the localities of their collecting are in most cases but imperfectly detailed. Since their time, with the exception of a few transient botanists who have incidentally worked over the neighborhood of some vacation resort, the work done on the flora may be summarized as follows :

In 1867 and 1871, Dr. A. S. Todd, as chairman of a committee of the Medical Society of West Virginia, published a list of the "Medicinal Plants of West Virginia." This list contains an enumeration of nine trees, seven shrubs and sixty herbs.

In 1870, Mr. DissDebarr, State Commissioner of Immigration, in his "Handbook of West Virginia," compiled a list of the timber trees of the State, in which he enumerated fifty-two species and added twelve species of shrubs.

In 1876, Professor Fontaine in compiling his portion of the Centennial volume upon the "Resources of West Virginia," listed more carefully the forest trees, shrubs and medicinal plants of the State, drawing the last from the publication of Dr. Todd. This work contains an enumeration of sixty-nine trees and sixteen shrubs.

In 1878, Profs. H. N. Mertz and G. Guttenberg manifolded a check list of the "Flora of West Virginia," being an account of work done along the upper Ohio bottoms, and in the mountains of the northeastern portion of the State, the latter while located at Harper's Ferry. This list enumerates fifty-nine trees, thirty-seven shrubs and four hundred ninety-four herbs.

In 1888 and 1889, Miss Verona Mapel, Preceptress of the High School at Glenville, Gilmer County, quite thoroughly worked over her immediate vicinity in connection with her school duties. She reports

in manuscript forty-two trees, twenty-three shrubs and two hundred ninety herbs. Her list includes neither the commoner weeds and herbs, nor the grasses and sedges.

In 1890, '91 and '92, the present authors began a systematic survey of the State, publishing the results of their work in 1892 under the title of "A Preliminary Catalogue of the Flora of West Virginia." This catalogue included all known previous work and enumerated 1,645 species, giving localities, numerous critical notes, and descriptions and plates of several new forms.

In 1892, Mr. John K. Small and Miss Anna Murray Vail spent some time in Greenbrier County, exploring the vicinity of White Sulphur Springs. The results of their work are included in their "Report of the Botanical Exploration of Southwestern Virginia," published as one of the Memoirs of the Torrey Botanical Club.

From 1892 to the present, the authors have continued their investigation of the flora, adding the knowledge of over one thousand species to their preliminary list, and detecting many new localities for previously published species. This additional work forms the basis of the present publication. The small edition of the preliminary catalogue having been exhausted within a few months of its issue, and many institutions, libraries and personal workers being unable to secure copies of the work, it has been deemed expedient to include here all the species of that publication, without, however, repeating the body matter. These repeated species appear in small capitals, additional species new to the flora in black-faced type. As in the preliminary catalogue, all species that have been described from known West Virginia types are republished in full.

#### SPECIAL FEATURES OF THE FLORA.

The distribution of plants within the boundaries of the State is wonderfully comprehensive. Canada places representatives of her boreal flora upon its Alleghanian mountain tops, some that have even passed by the States of New York, New Jersey, Pennsylvania, Delaware and Maryland in the transit; the Southern States contribute to its flora through the influence of the mysterious New River on the southeast; the great trunk lines of railway, as well as the open condition of the western border line along the Ohio River, give entrance to individuals of a migratory character from the Western Plains; and from some not readily accountable reason, Eastern forms, and even coast line species stray within its limits.

Of the noteworthy species we connect in *Anemone trifolia*, L., Canby's and Curtiss' Virginian stations with Knipe's Pennsylvanian, and that in a direct and sequential manner through the State. Our

elevations for this species range from 850 to 2,300 feet. *Trautvetteria Carolinensis* (Walt.) Vail, is to be found along all such mountain rills as are deeply shaded, at altitudes above 1,000 feet. We extend the Manual distribution of *Caltha palustris*, L., southward nearly one degree, by finding it quite plentiful in the mountains of Randolph County, where *Isopyrum trifoliatum* (L.) Britt., keeps it company. *Helleborus viridis*, L., has its most western station on the north branch of the Potomac River, in Hardy County, whence it was first reported to Dr. Gray by Dr. Gamble, of Moorfield. *Aconitum uncinatum*, L., is at home all along the banks of Cheat River, at altitudes varying from 780 to 3,550 feet. Both species of *Actæa* are to be met with on the higher Alleghanies, where *Cimicifuga Americana*, Mx., is the principal representative of the tribe.

*Magnolia Fraseri*, Walt., is a striking vernal feature of the whole Alleghanian region, while *tripetala* and *acuminata* are common. *Berberis Canadensis*, Mill. becomes a veritable weed in many fields in the southern portion of the State, vying with *Papaver dubium*, L., and *Glaucium Glaucium* (L.), Karst, in the northeastern section. *Bicuculla eximiana* (L.), Millsp., we have found only upon the highest peaks of the Alleghanies.

Of the rambling crucifers, *Sisymbrium Thaliana* (L.) Celak, finds a home in Fayette County, and *Lepidium campestre* (L.), R. Br., is the worst and most prevalent weed in the cultivated portions of the northeastern section.

Nearly all the violets lend their beauty to the adornment of the woods and meadows; *Viola pedata*, var. *bicolor*, Pursh., with flowers as large as the cultivated pansy and fully as beautiful, is frequent among the Devonian shales of the northeast; *Viola primulæfolia*, L., comes in from the coast as far as four degrees; and *Viola hastata*, Mx., is quite plentiful in the southern section.

Although *Sida hermaphrodita* (L.), Rusby, has not been found in the Alleghanies as yet, it is not at all rare along the New and Great Kanawha Rivers, from Quinnimont to the Ohio, and thence down that river to the limits of the State. *Hibiscus Moscheutos*, L., is found not only along the bottoms of the Great Kanawha, but also in the northeastern part of the State, while *H. Trionum*, L., is a quite common weed in gardens.

All of the native species of *Aesculus*, together with the var. *hybrida* of *octandra*, are found in the State. *Ailanthus glandulosis* is becoming a most troublesome weed in many sections, especially in the northern counties.

Among the Leguminals we have re-discovered *Astragalus Carolinianus*, L., which from our specimens is considered by Professor Britton

to be indistinct from *A. Canadensis*, L., and as it has the priority of publication, the latter well-known name becomes a synonym. Another important discovery in this genus is that *A. distortus*, T. and G., habits the Devonian shales of Hardy County, the only station known for the species east of the Mississippi valley. This species is here associated with *Opuntia polyacantha*, Haw, in great quantity, giving this peculiar spot in the Alleghanies much the appearance of an arid waste in Arizona. *Stylosanthes hamata* (L.), Britt., here ventures farther east than has heretofore been supposed, and finds congenial soil along New River in Fayette County. *Lespedeza striata* (Thunb.), H. & A., spreads profusely throughout the southwestern portion of the State. A new clover, *Trifolium Virginicum*, Small, has been discovered in Greenbrier County, by Mr. J. K. Small.

The roses are striking in the many new forms they produce: *Rubus Millspaughii*, Britt., is so profuse in the mountains of Pocahontas and Pendleton Counties that, according to the mountaineers and hunters, it is upon it that the bears depend principally for fattening food prior to hibernation. *Rubus Canadensis roribaccus*, Baily, the Leucretia dewberry, came originally from Randolph County. Another peculiar *Rubus* (*R. Columbianus* Millsp.), has 5 to 7-incised leaves of striking character. A new *Spirea* (*S. Virginiana*, Britt.) grows plentifully within half a mile of the University at Morgantown. Mercer County in the southern section of the State, presents a wonderful array of *Crategi*: *C. spatulata*, *C. cordata*, *C. apiifolia*, *C. coccinea*, *C. tomentosa*, *C. punctata*, *C. Crus-galli*, *C. flava* and its variety *pubescens*, and *C. uniflora* were all found during one day's botanizing in this section.

Of the Calycanths we have all, even the two species recorded "Virginia doubtful" in the Manual.

Of the saxifrages we have notably, *Astilbe decandra*, Don., *Saxifraga crosa*, Pursh., *Boykinia aconitifolia*, Nutt., *Heuchera villosa*, Mx., and *H. Americana*, L., and even the Laboradorian *Ribes prostratum*, L' Her.

*Sedum Pulchellum*, *Nevii*, *ternatum*, *telephioides*, and *telephium*, are with us. The beautiful *Liquidambar styraciflua* extends limitedly down the Gauly and Great Kanawha: The Onagraceæ yield a new form in *Ludwegia alternifolia*, L., var. *linearifolia*, Britt. The purple and yellow passion-flowers (*P. lutea*, L., and, *incarnata*, L.) grace the thickets.

Passing many minor forms, the Compositæ gives us *Elephantopus Carolinianus*, Willd. and *tomentosus*, L. (called as a weed "The Devil's Grandmother"); *Eupatorium caelestinum*, L., profuse; *Solidago Curtisii*, *S. rupestris*, and *Riddellii*, *Silphium perfoliatum*, L.; *Rudbeckia speciosa*, Wend.; *Helianthus grosse-serratus*, Mart., *H. doronicoides*, Lam., and

*H. laevigatus*, T. and G., *Verbesina Virginica*, L.; *Cacalia suaveolens*, *reniformis*, and *atriplicifolia*; *Cnicus Virginianus*, and *pumilus*, Torr.; *Cichorium Intybus*, L.; *Tragopogon porrifolius*, L.; *Hieracium Canadense*, Mx., and *longipilum*, Torr. (extending both); and *Chondrilla juncea*, L., called as a weed "Naked-weed, Skeleton-weed."

The interesting bell-worts are: the white form of *Lobelia siphilitica*, L., and the rare *Campanula divaricata*, Mx.

Of the rarer heaths *Schollera erythrocarpa*, Mx., grows on the highest peak of the Alleghanies, alt. 4,800 ft.; *Chiogenes hispidula* at the Falls of Blackwater; *Menziesia globularis* Salisb., *Clethra acuminata*, Mx., *Moneses grandiflora*, Salisb. and all the Rhododendrons except *Rhodora* and *Lapponicum*; even the rare *R. canescens* (Mx.), Porter, being found along the Cacapon River.

*Naumbergia thyrsiflora* is found in Upshur County and *Mohrodendron Carolinum* (called Shittim-wood) is plentiful along the Gauly and New Rivers. *Polemonium Van-Bruntiae*, Britt., comes south to our flora, as well as all the *Hydrophyllums*, together with *Phacelia Purshii*, Buck, and *parviflora*, Pursh.

The beautiful morning-glories, *Impomoea coccinea*, *hederacea*, *purpurea*, and *pandurata*, are all too plentiful as weeds here; and *Cuscuta glomerata*, *Gronovii*, and *Epithymum*, have been found sparingly. *Physalis viscosa*, L., steals away from "near the coast" and is found along the Ohio River, keeping company with *Lycium vulgare*, Dun., and *Physalodes Physaloides*, Gaertn.

The notable *Scrophularias* are: *Collinsia verna*, Nutt., *Chelone obliqua*, L., and *Pentstemon Canescens*. Of the mints we have notably: *Koellia verticillata*, *clinopodioides*, *pycanthemoides*, and *Montana*. The other mints worthy of remark are: *Meehania cordata*, *Clinopodium vulgare*, *Scutellaria saxatilis*, *serrata*, *incana*, *parvula*, and *nervosa*; *Marrubium vulgare*, *Galeopsis tetrahit*, and *Stachys palustris* and *cordata*.

Of the ten Euphorbias the most notable are *E. Darlingtonii* and *E. Glyptosperma*, var., *pubescens*, Engl., the latter not having been previously found east of Iowa as far as we can learn.

The presence of *Quercus ilicifolia*, Wang, in Hardy Co., extends the Manual distribution southeastward; and the southing of *Q. macrocarpa*, Mx. is also extended by several stations in the State.

As to the conifers, we have about 470,000 acres of *Picea Mariana*, a few representatives of *Abies balsamea*, *Thuja occidentalis*, and several species of *Pinus*, as well as a few scant growths of *Taxus Minor*.

Among the sedges the principal item of interest is the re-discovery in Fayette County of what was doubtless the original type station of *Carex Fraseri*, And.



Of the *Equisetaceæ* the most notable form so far found is *E. lævigatum*, Braun., gathered in the southernmost part of the State thus extending its distribution southeastward.

Of the *Filices*, the rarer forms found with us are: *Polypodium polypodioides*; *Pellæa atropurpurea* in great quantities in the southern section; *Asplenium pinnatifidum*, *montanum* (plentiful) and *angustifolium*; *Dryopteris Goldieana*, and *marginalis*. *Cystopteris bulbifera*; *Dicksonia punctilobula*; and strange to say on the summit of Spruce Knob at an altitude of 4,800 ft. *Dryopteris fragrans*, in such great quantity that it is cut and stacked for fodder, this species being greatly relished by cattle.

*Lycopodium lucidulum*, L., *annotinum*, L., *obscurum* and its var. *dendroideum*, L. *clavatum*, and L. *complanatum* are all found in the forests of black spruce along the Alleghanies.

In the mosses, hepatics, and lichens, but little collecting has so far been done, no systematic searches having been made for specimens in these classes of plants. Among the lichens several new species have been discovered, the descriptions of which are still in manuscript with Professor Nylander. In the search for hepatics incidental to other exploration, in Mercer County, the dry bald face of a large limestone cave yielded a new species in *Plagiochila Virginica* Evans, as well as a rarity in the eastern flora of the United States, *Radula Xalapensis*, Mont. Among the mosses we have been rewarded in our itinerant work by finding two new forms *Dicranodontium Virginicus*, Britt. m. and *D. Millspaughi* Britt. m., as well as numerous noteworthy species.

The Fungi have been found to be of special interest, and it is to their collection and study that Mr. Nuttall has devoted most of his spare hours from business since 1893. His field of search for forms in this class of plants has been very limited, being almost wholly the immediate neighborhood of his home at Nuttallburg, in Fayette County, on New River. Even this small area has furnished the major part of the 980 fungi of this flora, and continues to present additional forms as well as unique hosts upon every search, no matter how casual the examination or short the time devoted to the trip. Fully two-thirds of the species collected have passed under the critical examination of Mr. J. B. Ellis, whose careful consideration of our numbers has been of incalculable assistance in this work.

The new species discovered in this area, and described in this Flora, are as follows:

Anthostoma micreecium.	<b>Cercospora Chionanthi.</b>	<b>Corticium leptaleum.</b>
Aposphæria pezizoides.	Cercospora (E)notherae.	Coryneum cupulatum.
<b>Aspergillus glaucus oblongisporus.</b>	<b>Cercospora septorioides.</b>	Cyathicula quisquillaris.
Botryodiplodia acerina.	<b>Clasterosporium Cornutum.</b>	Cylindrocolla Dendroctoni.
<b>Botrytis olivacea.</b>	Cladosporium nigrellum.	<b>Cylindrocolla flagellaris.</b>
<b>Camarsporium Linderæ.</b>	<b>Corticium albo-flavescens.</b>	Cylindrosporium Cratæg.

- Cytispora caryigena.*  
*Cytispora Celtidis.*  
*Cytispora exasperans.*  
*Cytispora Halesiæ.*  
*Cytispora Rhois-hirtæ.*  
*Cytispora Sassafra.*  
*Dermatella viticola.*  
*Diaporthe Araliæ.*  
*Diaporthe Halesiæ.*  
**Diaporthe Hydrangeæ.**  
*Didymella Physocarpi.*  
*Diaporthe Tetrapteræ.*  
*Diplodia caryigena.*  
*Diplodia Cercidis.*  
*Diplodia infuscans.*  
**Dothiorella Asiminæ.**  
**Dothiorella minor.**  
*Eutypella densissima.*  
*Fusicoccum ilicinum.*  
**Glœosporium Alni.**  
**Glœosporium rubicolum.**  
**Glœosporium Rumicis.**  
*Glœosporium Sanguinariæ.*  
*Haplosporella Araliæ.*  
*Haplosporella Celtidis.*  
**Helminthosporium brachypus.**  
*Helotiella Nuttallii.*  
*Hypocrea tenerrima.*  
*Hypocrea Virginiensis.*  
*Hypoxyton atroviride.*  
*Hymenula cerealis.*  
*Hypoxyton Nuttallii.*  
*Illosporium cæspitosum.*  
*Isaria Virginiensis.*  
*Lachnella Virginica.*  
*Lophidium nitidum.*  
*Macrosporium olivaceum.*  
**Massaria Magnoliæ.**  
*Microcera erumpens.*  
*Myxosporium luteum.*  
*Myxosporium platanicum.*  
*Myxosporium seriatum.*  
*Pestalozzia Toxicæ.*  
*Phlyctænâ Ipomœæ.*  
**Phoma Ascelepiadeæ.**  
*Phoma negundinicola ramicola.*  
*Phoma obscurans.*  
*Phoma pedunculi.*  
*Phoma Pennsylvanica.*  
*Phyllosticta Araliæ.*  
*Phyllosticta Castaneæ.*  
**Phyllosticta Celastri.**  
**Phyllosticta cercidicola.**  
**Phyllosticta globifera.**  
*Phyllosticta lindericola.*  
*Phyllosticta macrospora.*  
**Phyllosticta opaca.**  
**Phyllosticta Oxydendri.**  
*Phyllosticta Quercus-Prini.*  
*Phyllosticta rhoicola.*  
**Phyllosticta Ribis.**  
**Phyllosticta Xanthorrhizæ.**  
*Pilacre gracilipes.*  
*Pleosphæria corticola.*  
**Saccidium vitis.**  
**Septoria Brassicæ.**  
*Septoria hyalina.*  
*Septoria Polymniæ.*  
*Septoria Tecomæ.*  
*Septoria Trautvetteriæ.*  
*Septosporium Equiseti.*  
*Sirococcus Halesiæ.*  
**Sphæronæma infuscans.**  
*Sphæronæma Physocarpi.*  
*Sphæropsis Asiminæ.*  
**Sphæropsis Asiminæ fructigena.**  
*Sphæropsis Ipomœæ.*  
**Sphæropsis Cœnotheræ.**  
*Sphæropsis Physocarpi.*  
*Steganosporum pedunculi.*  
*Stagonospora petiolorum.*  
*Stagonospora Physocarpi.*  
*Stachylidium caricinum.*  
**Trematosphæria vitigena.**  
*Trichægum nodulosum.*  
*Tubercularia hamata.*  
*Tympanis Oxydendri.*  
**Valsa albo-puncta.**  
*Valsa Chionanthi.*  
*Valsa Diospyri.*  
*Valsa etherialis.*  
**Vermicularia Trautvetteriæ.**  
**Verticillium osteophilum.**

Of these those in black-faced type are here described for the first time.

Beside these unique forms, we report many species from our region that have not been before credited to the flora of North America, include many hitherto unpublished asci and spore measurements of species otherwise well described, and have transferred many not before well understood to their proper genera.

The host plants have proven also to be of special interest in that many of them yield certain species for the first time in the mycologic literature of this country, and many others pose as altogether new to Host Indices.

#### THE SYLVA.

The State is very happily located for the growth of forests, being in the favored belt of temperature between about 37° and 41° north latitude. Within its boundaries trending northeast and southwest, thus opening the country to the damp, warm winds from the Gulf, are numerous ranges of the great Appalachian, or Atlantic highlands, but by far the larger portion of the State lies on the westward slope of these mountains in the trans-Appalachian belt, the waters of which chiefly run northwestward and southwestward to the Ohio River. The altitude of the country descends from the Backbone or Alleghany range of the Appalachians, from an altitude of

from 2,500 and 4,800 feet to 500 at the southwestern corner of the State on the Ohio, at Kenova, and about 600 on the same river at Wheeling. The altitude of the eastern corner of the State at Harper's Ferry is 272 feet; thus the range of altitudes in the State is from 272 feet to about 4,800, giving a climatic range of 3,728 feet, or the equivalent of about 16° of latitude; consequently West Virginia has extensive areas of adaptability for every variety of forest growth that is found within the limits of the northern States east of the Rocky Mountains.

The most elevated portion of the State is the great eastern border of the ridgy plateau from which the trans-Appalachian country descends, a territory some 200 miles in length from the headwaters of the Big Sandy to those of the North Branch Potomac, this region is in the main from 2,500 to 4,800 feet in altitude, and furnishes a congenial home to the black spruce, the white pine, and other evergreen trees peculiar to northern latitudes.

West Virginia has a greater amount of hardwood timber in its forests than any other State in the Union. A thorough examination convinces us that nearly or quite two-thirds of the State remains uncleared, and by far the greater portion of the uncleared land is still in virgin forests where the ax of man has never found its way, and where magnificent specimens of forest growth stand thickly side by side and reach a towering height, no finer view of standing timber may be had within the confines of the Union. These splendid forests covering over sixteen thousand square miles yield nearly every species found in the north. Here trees grow to such size that ordinary methods will not suffice to handle them, and are frequently so densely compact that the light of day scarce penetrates their shade, and pathways must be cut before the ax men can find room to work.

The following list of the trees of the State, giving the diameter of trunks and width of board yield, is that of the State exhibit in the Forestry Building at the World's Columbian Exposition, 1893. The specimens were all gathered during the winter of 1892-3.

Ash (white), *Fraxinus Americana*. Logs 40 in., 62 in., 93 in., planks 27 and 33 in., finished boards, plain, 20 and 21 in., mottled, 9, 13 and 16 in., figured 9 and 13 in.

Ash (black), *F. nigra*.

Ash (mountain), *Sorbus Americana*. Trunk 8 in.

Ash (prickly), *Xanthoxylum Americanum*. Trunk 12 in.

Alder (mountain), *Alnus viridis*. Trunk 4 in.

Beech, *Fagus atropunicea*. Trunks 24 in., 27 in. and 38 in., plank 27 in., finished boards 27 and 31 in., quartered boards 7 and 9 in.

Beech (water), *Carpinus Caroliniana*. Trunk 10 in.

Birch (red), *Betula lenta*. Trunks 34 in., 93 in., 61 in., 48 in., planks 15 and 30 in., finished boards 12 and 16 in., figured boards 8, 13, and 14 in.

Birch (yellow), *Betula lutea*. Trunks 34 in., 36 in., 42 in., 49 in., plank 26 in.

Birch (hybrid), *Betula lenta* x *lutea*. Trunk 2 in.

Basswood (white), *Tilia Americana*. Trunks 32 in., 36 in., 43 in., 51 in., plank 28 in., finished board 16 in.

Basswood (yellow), *Tilia heterophylla*. Trunk 26 in.

Buckeye, *Aesculus glabra*. Trunk 29 in.

Buckeye (purple), *A. octandra hybrida*. Trunk 29 in.

Box Elder, *Acer Negundo*. Trunk 27 in.

Butternut, *Juglans cinerea*. Trunk 12 in.

Balsam Fir, *Abies balsamea*. Trunk 13 in.

Chestnut, *Castanea dentata*. Trunks 53 in., 40 in., 62 in., 78 in., plank 40 in., finished boards 26 in.

Cherry (wild), *Prunus serotina*. Trunks 36 in., 39 in., 48 in., 53 in., plank 27 in., finished boards 6 and 18 in., blistered 13, figured 6 to 19 in., curly 9 to 21 in.

Cherry (wild), *Prunus Pennsylvanica*. Trunk 14 in.

Cucumber, *Magnolia acuminata*. Trunk 35 in., plank 26 in.

Cottonwood, *Populus monilifera*. Trunk 26 in.

Crab (wild), *Pyrus coronaria*. Trunk 8 in.

Dogwood, *Cornus florida*. Trunks 7 in., 13 in., 28 in.

Elm (White), *Ulmus Americana*. Trunks 23 in., 42 in., 61 in.

Elm (red, slippery), *Ulmus pubescens*. Trunk 16 in.

Fringe tree, *Chionanthus Virginica*. Trunk 3 in.

Gum (black), *Nyssa sylvatica*. Trunk 36 in., plank 27 in.

Gum (sweet), *Liquidambar styraciflua*. Block 19 in.

Grape (Fox), *Vitis Labrusca*. Trunk 22 in., branches 18 in.

Hackberry, *Celtis occidentalis*. Trunk 18 in.

Haw (black), *Viburnum prunifolium*. Trunk 5 in.

Hemlock, *Tsuga Canadensis*. Trunk 42 in., plank 30 in., finished boards 12 and 19 in.

Hercules Club, *Aralia spinosa*. Trunk 5 in.

Hickory (bitter), *Hicoria minima*. Trunks 9 and 12 in.

Hickory (red), *Hicoria glabra*. Trunks 11 and 17 in.

Hickory (shagbark), *Hicoria ovata*. Trunk 27 in., planks 14 and 18 in.

Holly (white), *Ilex opaca*. Trunk 10 in.

Holly (mountain), *Ilex monticola*. Trunk 6 in.

Ironwood, *Ostrya Virginiana*. Trunks 12 in., 24 in., 38 in.

Juniper, *Juniperus Virginiana*. Trunk 7 in.

- Laurel, *Kalmia latifolia*. Trunk 8 in., root 24 in.  
 Locust (yellow), *Robinia Pseudacacia*. Trunks 18 in., 24 in., 38 in.  
 Locust (honey), *Gleditchia triacanthos*. Trunks 14 in., 28 in., 30 in., in thorn 10 in.  
 Magnolia, *Magnolia tripetala*. Trunk 15 in.  
 Magnolia (ear-leaf), *Magnolia Fraseri*. Trunk 12 in.  
 Mulberry, *Morus rubra*. Trunks 8 and 11 in.  
 Mulberry (paper), *Papyrius Papyrifera*. Trunk 9 in.  
 Maple (blistered), *Acer saccharum*, Marsh. Trunk 26 in., boards 9 and 11 in.  
 Maple (sugar), *Acer saccharum*, Marsh. Trunks 30 in., 40 in., 48 in., plank 28 in., finished boards 29 in., curly boards 13 to 16 in.  
 Maple (black), *Acer nigrum*. Trunk 16 in.  
 Maple (white), *Acer saccharinum* L. Trunks 35 in., 49 in., plank 26 in., birdseye boards 10 to 12 in.  
 Maple (red), *Acer rubrum*. Trunk 14 in.  
 Maple (striped), *Acer Pennsylvanicum*. Trunk 6 in.  
 Osage Orange, *Toxylon pomiferum*. Trunk 4 in.  
 Oak (black), *Quercus velutina*. Trunks 10 and 27 in., finished board 11 in.  
 Oak (Spanish), *Quercus digitata*. Trunks 13 and 18 in.  
 Oak (swamp), *Quercus palustris*. Trunk 15 in.  
 Oak (laurel), *Quercus imbricaria*. Trunk 10 in.  
 Oak (chestnut), *Quercus Prinus*. Trunks 14 in., 27 in., 36 in., finished board 21 in.  
 Oak (black jack), *Quercus nigra*. Trunks 18 in., 23 in.  
 Oak (white), *Quercus alba*. Trunks 50 in., 28 in., 32 in., 61 in., plank 37 in., boards (quartered) 5 to 13 in., quartered curly 10 in.  
 Oak (red), *Quercus rubra*. Trunks 42 in., 54 in., 84 in., plank 44 in.  
 Persimmon, *Diospyros Virginiana*. Trunk 12 in.  
 Poplar (yellow), *Liriodendron Tulipifera*. Trunk 48 in., trunk "wheels" 24, 36, 48, 60, 74, 90, and 138 in., the tree from which the last wheel came cut 25,000 feet of merchantable boards, plank 36 and 57 in., cube 36 in., finished boards 27 to 46 in.  
 Pine (white), *Pinus strobus*. Trunk 36 in.  
 Pine (yellow), *Pinus echinata*. Trunk 28 in.  
 Pine (pitch) *Pinus rigida*. Trunk 16 in.  
 Papaw, *Asimina triloba*. Trunk 8 in.  
 Rhododendron, *Rhododendron maximum*. Trunk 6 in.  
 Red Bud, *Cercis Canadensis*. Trunks 5½ and 9 in.  
 Scyamore, *Platanus occidentalis*. Trunks 52 in., 36 in., 40 in., finished board quartered 33 in.

- Sassafras, *Sassafras Sassafras*. Trunk 22 in., finished board "wavy" 18 in.
- Silver Bell, *Mohrodendron Carolinum*. Trunk 8 in.
- Sourwood, *Oxydendron arboreum*. Trunk 15 in.
- Spruce (black) *Picea Mariana*. Trunks 31, 34, and 49 in., planks 8 and 40 in., finished boards 18 and 20 in.
- Spicewood, *Benzoin Benzoin*. Trunk 2 in.
- Sumach, *Rhus typhina*. Trunk 8 in.
- Service, *Amalanchier Canadensis*. Trunks 8 and 9 in.
- Thorn, *Crategus coccinea*. Trunk 10 in.
- Tree of Heaven, *Ailanthus glandulosus*. Trunks 8 in., 24 in., 38 in.
- Witch Hazel, *Hamamelis Virginica*. Trunks 4 and 5 in.
- Walnut (black) *Juglans nigra*. Trunks 30 and 36 in., plank 29 in. finished boards "wavy" 20 in., "figured" 21 in.
- Willow (black), *Salix nigra*. Trunk 12 in.

## SUMMARY OF THE FLORA.

	Genera Species.			Genera Species.	
Fungi, etc.,	342	980	Equisitæ,	1	4
Lichens,	31	115	Filicinæ,	14	40
			Ophioglossæ,	2	7
Thallophyta, etc.,	373	1095	Lycopodæ,	1	5
			Selaginellæ,	1	1
Hepaticæ,	24	32			
Musci	42	90	Pteridophyta,	19	57
Sphagnæ,	1	1			
			Gymnospermæ,	7	13
Bryophyta,	67	123	Monocotyledons,	95	268
			Dicotyledons,	412	1028
			Anthophyta,	514	1309
Total number of species, varieties and forms detected in the State to the date of this Flora,					2584

# CATALOGUE.\*

## Fungi.

### PHALLACEÆ.

#### ITHYPHALLUS Fries.

##### **I. impudicus** (Linn.) Fr.

In the juvenile, egg stage, June 15, 1893, Oct. 28, 1893, we find the veil under pileus about one-third its length and part of it encircling the stipe about the middle (*Nuttall*, 983).

### NIDULARIACEÆ.

#### CRUCIBULUM Tul.

##### **C. vulgare** Tul.

On sticks and leaves, open woods, and on old cotton cloth, July 6, 1893 (*Nuttall*, 1079).

### LYCOPERDACEÆ.

#### TYLOSTOMA Pers.

##### **T. mammosum** (Mich.) Fr.

On light soil on rock, March 24, 1893 (*Nuttall*, 880).

### MITREMYCES Nees.

##### **M. lutescens** Schn.

On wet mossy banks, alt. 2,000 ft., March 25, 1893 (*Nuttall*, 881).

### GEASTER Mich.

##### **G. mammosus** Chev.

On earth, Short Creek, July 6, 1893 (*Nuttall*, 1080).

### BOVISTA Dill.

##### **B. lepidophora** (E. & E.) De Ton.

In grass on lawn, Sept. 5, 1893. Clavate 8 cm. high, 12 cm. broad, sterile base 3.5 cm. thick, white cuticle 1.5 mm. thick, gleba pale yellow (*Nuttall* 1193).

##### **B. PILA** Berk. & Curt.

Free on open ground. Monongalia Co., near Morgantown (*Millspaugh*).

\*Unless otherwise stated, all my numbers in this Flora are from Fayette Co., near Nuttallburg.—*L. W. Nuttall*.

## LYCOPERDON Tourn.

**L. gemmatum** Batsch.

On ground in woods, alt. 2,000 ft., Aug. 24, 1893 (*Nuttall*, 1180).

**L. furfuraceum** Schæff.

In grass on lawn, Sept. 3, 1893. Capillitium thicker than the spores, branches few, 2.5 to 4 cm. in diameter, outer coat pure white (*Nuttall*, 1188).

**L. pedicellatum** Peck.

On moss on ground in pine woods, alt. 2,000 ft., March 10, 1893. Pedicels 20  $\mu$  long (*Nuttall*, 864).

**L. PYRIFORME** Schæff.

Under bark of *Quercus palustris*, Monongalia Co., near Morgantown and near Little Falls (*Millspaugh*) on rotten wood, Feb. 2, 1893 (*Nuttall*, 832).

## SCLERODERMA Pers.

**S. VULGARE** Hornem.

On chips of *Picea Mariana*, Tucker Co. Falls of Blackwater (*Millspaugh*). Grant Co., on dead logs, Otter Fork of Cheat; and Monongalia Co., Tibbs Run, plentiful on clay of a path (*Millspaugh*). In laurel thickets, July 28, 1893. Spores 10  $\mu$ . (*Nuttall*, 1125).

**S. Bovista** Fr.

On ground, March 21, 1893 (*Nuttall*, 876).

## BOVISTELLA Morg.

**B. Ohioense** (Ell. & Morg.) Morg.

In grass on lawn, June 26, 1893 (*Nuttall*, 1003).

## MUCORACEÆ.

## MUCOR Mich.

**M. MUCEDO** Linn.

On open canned fruit left standing (*Millspaugh*). On dead dry fruit *Asimina triloba*, Oct. 1, 1895 (*Nuttall*, 1866, 754).

## SPORODINIA Link.

**S. Aspergillus** (Scop.) Schroet.

On dead *Agaricus* and *Boletus*, Sept. 10, 1893 (*Nuttall*, 1196).

## PERONOSPORACEÆ.

## CYSTOPUS Lev.

**C. CANDIDUS** (Pers.) Lev.

On living leaves of *Dentaria diphylla*, Monongalia Co., at Little Falls (*Millspaugh*). On *Brassica nigra*, July 8, 1893 (*Nuttall*, 1082).



**C. Portulacæ** (DC) Lev.

On *Portulacca oleracea*, July 8, 1893 (*Nuttall*, 1083).

**C. Bliti** (Biv.) DeB.

On *Amaranthus retroflexus*, July 20, 1893. Spores 18 to 22 x 15 to 18  $\mu$  (*Nuttall*, 1615).

PLASMOPARA Schroet.

**P. VITICOLA** (B. & C.) Berl. & De Ton.

On fruit of *Vitis Labrusca*, Monongalia Co., near Morgantown (*Millsbaugh*).

PHYTOPHTHORA DeBary.

**P. INFESTANS** (Mont.) DeB.

On living leaves and tubers *Solanum tuberosum*, Monongalia Co., near Morgantown (*Millsbaugh*).

BREMIA Regel.

**B. Lactucæ** Regel.

On *Lactucca Canadensis* and *L. hirsuta*, June 14, 1894 (*Nuttall*, 1549).

PERONOSPORA Corda.

**P. OBOVATA** Bon.

On living leaves *Spergula arvensis*, Preston Co., near Terra Alta (*Millsbaugh*).

ENTOMOPHTHORACEÆ Nowak.

EMPUSA Cohn.

**E. MUSCÆ** (Fr.) Cohn.

On *Musca domestica*, Monongalia Co., at Morgantown (*Millsbaugh*). Very prevalent on a species of *Tachina* found on maple leaves in great numbers in 1892 at Morgantown (*Millsbaugh*).

**E. GRYLLI** Fr.

On tufted caterpillars, on grasshoppers, and on the house fly, Monongalia Co., at Morgantown (*Millsbaugh*).

SCHIZOMYCETACEÆ.

BACCILLUS Cohn.

**B. TUBERCULOSIS** Koch.

In sputa of consumptive (*Millsbaugh*).

**B. ACIDI-LACTICI** (Zopf) Schroet.

In soured milk (*Millsbaugh*).

**B. SUBTILIS** (Ehrenb.) Cohn.

In infusion of hay and on exposed boiled potato (*Millsbaugh*).

**B. ULNA** Cohn.

On exposed coagulated egg albumen (*Millsbaugh*).

## SPIRILLUM Ehrenb.

- S. UNDULA (Muell) Ehrenb.  
In infusion of hay (*Millspaugh*).

## BACTERIUM Ehrenb.

- B. LINEOLA (O. F. Muell) Cohn.  
In infusion of radish (*Millspaugh*).
- B. TERMO (O. F. Muell.) Ehrenb.  
In various decomposing organic substances (*Millspaugh*).

## MICROCOCCUS (Hall.) Cohn.

- M. CREPUSCULUM (Ehrenb.) Cohn.  
Found associated with *Bacterium Termo* in decomposing vegetable infusions (*Millspaugh*).
- M. AMYLOVORUS Burrill.  
On fruit of *Pyrus communis*, Monongalia Co., near Morgantown (*Millspaugh*).
- M. AURANTIACUS, Cohn.  
Caught on sterilized potato in laboratory (*Millspaugh*).
- M. LUTEUS Cohn.  
Caught on sterilized potato in laboratory (*Millspaugh*).

## STREPTOCOCCUS Billr.

- S. PYOGENES (Rosenb.) Zopf.  
*Micrococcus septicus*, Cohn. In blood of dead calf (*Millspaugh*).
- S. UREÆ (Cohn.) Trev.  
*Micrococcus ureæ*, Cohn. In decomposing urine (*Millspaugh*).

## SACCHAROMYCETACEÆ.

## SACCHAROMYCES Meyen.

- S. CEREVISIÆ Meyen.  
In Pasteur's liquid left uncorked in laboratory (*Millspaugh*).
- S. MYCODERMA Reess.  
On same liquid as above at same date (*Millspaugh*).

## DIATOMACEÆ.

- CYMBELLA GASTROIDES, Kuetz.  
CYMBELLA TURGIDA (Grun.) Greg.  
STAURONEIS PHENICENTRON Ehrb.  
NAVICULA VIRIDIS, Kuetz.  
NAVICULA MAJOR, Kuetz.  
NAVICULA NOBILIS (Ehrb.) Kuetz.

NAVICULA RHOMBOIDES, Ehrb.  
 NAVICULA BOREALIS (Ehrb.) Kuetz.  
 NAVICULA TRINODIS, Lewis.  
 ACHNANTHES LANCEOLATA, Breb.  
 SYNEDRA ULNA (Nitzsch.) Ehrb.  
 NITZSCHIA AMPHIOXYS INTERMEDIA, Grun.

MYXOMYCETEÆ.

PHYSARUM Pers.

- P. citrinum** Schum.  
 On moss, alt. 2,000 feet, Aug. 2, 1893, (*Nuttall*, 1160).  
**P. pulcherrimum** B. & R.  
 On dead wood, July 25, 1893 (*Nuttall*, 1133).  
**P. psittacinum** Ditm.  
 On fruit cones of *Magnolia Fraseri*, alt. 2,000 ft., Aug. 12, 1893  
 (*Nuttall*, 1165).  
**P. sinuosum** (Bull.) Rost.  
 On bark of *Aralia spinosa*, Aug. 16, 1893 (*Nuttall*, 1154).

LEOCARPUS Link.

- L. fragilis** (Dicks.)  
 On *Aspidium spinulosum* and twigs of *Tsuga Canadensis*, Nov.  
 25, 1895 (*Nuttall*, 1888).

TILMADOCHÉ Fr.

- T. nutans** (Pers.) Rost.  
 On dead *Asparagus* leaves, and dead limbs *Magnolia Fraseri*,  
 Nov. 4, 1893 (*Nuttall*, 1264).  
**T. gyrocephala** (Mont.) Rost.  
 On dry bark of *Hicoria ovata*, living leaves of *Hydrangea*,  
 etc., Short Creek, alt. 1,300 ft., Aug. 21, 1893 (*Nuttall*, 1168).  
**T. viridis** (Gmel.) ?  
 On dead limbs *Magnolia Fraseri*, Sept., 1895 (*Nuttall*, 1856,  
 744?)

DIDYMIUM Schrad.

- D. squamulosum** (Alb. & Schw.) Fr.  
 On new timber in mine near the entrance, Sept., 1893 (*Nuttall*,  
 1342).

DIACHÆA Fries.

- D. leucopoda** (Bull.) Rost.  
 On dead wood and leaves in damp, shady place, June 23, 1893  
 (*Nuttall*, 998).

LAMPRODERMA Rost.

- L. violaceum** (Fr.) Rost.  
 On very rotten wet log, Short Creek, alt. 1,300 ft., Oct. 25,  
 1893 (*Nuttall*, 1255).

## COMATRICHA P. Hoyer.

**C. Friesiana** (DeB.) Rost.

On rotten log, Nov. 6, 1893 (*Nuttall*, 1237).

## STREMONITIS Gled.

**S. fusca** Roth.

On rotten log, and on *Hydnum* sp., April 7, 1893 (*Nuttall*, 889).

**S. ferruginea** Ehrh.

On log, Keeney's Creek, June 9, 1893 (*Nuttall*, 973).

## TUBULINA Pers.

**T. cylindrica** (Bull.) DeC.

On dead log. Spores brown, rough (reticulate?) approx. 6  $\mu$  diameter, Oct. 20, 1893 (*Nuttall*, 1230).

## LINDBLADIA Fries.

**L. effusa** (Ehr.) Rost.

Forms large patches on dead *Tsuga Canadensis* at Short Creek Cliff and Masterson's Glade, July 15, 1893 (*Nuttall*, 1130).

## CRIBRARIA Pers.

**C. aurantiaca** Schrad.

On dead wet log, July 25, 1895 (*Nuttall*, 1826, 712).

## RETICULARIA Bull.

**R. Lycoperdon** Bull.

On dead log, Short Creek, April 6, 1893 (*Nuttall*, 888).

## ARCYRIA Hall.

**A. punicea** Pers.

On rotten stump, June 15, 1893 (*Nuttall*, 979).

**A. cinera** (Bull.) Schum.

On damp dead wood and weeds in shade, July 24, 1893 (*Nuttall*, 1118).

**A. nutans** (Bull.) Grev.

On dead log, Short Creek. Spores 7.5  $\mu$ , July 24, 1893 (*Nuttall*, 1119).

## LYCOGALA Mich.

**L. epidendron** Buxb.

On wet dead logs, June 15, 1893 (*Nuttall*, 995). On *Magnolia Fraseri*, Nov. 1895.

## TRICHIA Hall.

**T. fallax** Pers.

On side of dead log, Oct. 19, 1893. Rich reddish-brown color when fresh, shining pale-brown when dry; elaters long and slender pointed, spores 10 to 12.5  $\mu$  (*Nuttall*, 1209).

**T. CHRYSOSPERMA** (Bull.) DeC.

On decayed wood, Monongalia Co., near Morgantown (*Mills-  
paugh*). On dead logs, Feb. 8, 1893 (*Nuttall*, 842).

**T. proximella** Karst.

On *Liquidambar styraciflua*, Feb. 6, 1893 (*Nuttall*, 836).

HEMIARCYRIA Rost.

**H. RUBIFORMIS** (Pers.) Rost.

Under bark of *Fraxinus Americana* and *Quercus alba*, Monongalia Co., near Morgantown (*Mills-  
paugh*). On bark of dead *Quercus* sp. Feb. 13, 1893 (*Nuttall*, 843).

**H. CLAVATA** (Pers.) Rost.

Under bark of wet decaying log *Quercus alba*. Monongalia Co., near Morgantown (*Mills-  
paugh*). On dead log, June 15, 1893 (*Nuttall*, 831).

**H. stipata** Schw.

On dead wet limbs of *Magnolia Fraseri*, Sept. 16, 1895 (*Nutt-  
all*, 1847).

HYPHOMYCETACEÆ.

OOSPORA Wallr.

**O. fasciculata** (Berk.) Sacc.

On decaying orange, Jan. 15, 1893 (*Nuttall*, 1134).

MONILIA Pers.

**M. aureo-fulva** C. & E.

On under side dead log *Liquidambar styraciflua*, Aug. 29, 1893 (*Nuttall*, 1186).

**M. FRUCTIGENA** Pers.

On ripe fruit *Prunus cerasus* cult. Monongalia Co., near Mor-  
gantown (*Mills-  
paugh*). On fruit of *Prunus domestica* cult. and *Amygdalus Persica* cult., June 28, 1893 (*Nuttall*, 1006).

POLYSCYTALUM Riess.

**P. sericeum** Sacc.

On *Quercus Prinus*, Nov., 1893 (*Nuttall*, 1335, 293).

OIDIUM Link.

**O. erysiphoides** Fr.

On *Zizia cordata*, *Eupatorium purpureum* and *Rubus odoratus*,  
Nov. 18, 1893 (*Nuttall*, 1266, 210).

**O. LEUCOCONIUM** Desm.

On leaves of *Rosa* cult., Cabell Co., near Huntington (*Mills-  
paugh*).

**O. MONILIOIDES** Link.

On living leaves *Poa pratensis*, Preston Co., near Terra Alta  
(*Mills-  
paugh*). On same host, May 30, 1894 (*Nuttall*, 1512).

## BOTRYOSPORIUM Corda.

**B. pulchrum** Corda.

On *Lactuca Canadensis*, Aug. 23, 1895. Tips of branches inflated. Spores  $7.5 \times 2.5 \mu$  (Nuttall, 1844, 735).

## TRICHODERMA Pers.

**T. lignorum** (Tode) Harz.

Under bark rotten *Hicoria ovata* Short Creek, alt. 1,250 ft., Dec. 8, 1893 (Nuttall, 1299, 248).

## ASPERGILLUS Mich.

**A. glaucus** (Linn.) Link.

On *Crategus* sp., April 30, 1894 (Nuttall, 1474).

**A. glaucus oblongisporus** E. & E. n. var?

Found on *Lachnocladium semivestitum* after neglect in plant press (Nuttall, 1871, 760).

This is either a new species or a distinct variety of *A. glaucus*. The smooth oblong-elliptical conidia are quite different from the usual form. Spores  $5$  to  $7.5 \times 2.5$  to  $3 \mu$ .

**A. flavus** Lk.

Also found on *Lachnocladium semivestitum* neglected in plant press (Nuttall, 1877, 760).

Conidia  $4$  to  $6 \mu$  diameter.

## STERIGMATOCYSTIS Cram.

**S. nigra** v. Tiegh.

On dried peach in garden, Aug. 15, 1893 (Nuttall, 1170).

## PENCILLIUM Link.

**P. glaucum** Link.

On *Hydnum* sp. and *Polyporus varians*, Aug. 16, 1893 (Nuttall, 1164, 136).

## BOTRYTIS Mich.

**B. olivacea** E. & E. sp. nov.

TYPE HABITAT: On dead log. Short Creek, alt. 1,800 ft., Feb. 26, 1894 (Nuttall, discov. 1411, 381). *Fungi Columbiana*, 593. N. A. F., 3187.

Forming thin olivaceous patches 2 to 4 cm. in extent, composed of olive-brown, septate threads, about  $4 \mu$  thick, and dichotomously or oppositely branched above, the short ( $15$  to  $25 \mu$ ) branches sub-attenuated above, and bearing the obovate or elliptical, olive-brown,  $6$  to  $8 \times 4$  to  $4.5 \mu$  conidia at their tips.

**B. torta** E. & E., Proc. Phila. Acad., 1894, 375.

TYPE HABITAT: On dead leaves *Carex Fraseri*, Dec. 10, 1893 (Nuttall, discov. 1248, 257).

Hyphæ simple, sparingly branched, twisted above as in *B. streptothrix* or in *Streptothrix atra* B. & C., brown, 80 to 100 x 3 to 4  $\mu$ , forming numerous small brownish-black tufts, effused or gregarious, on both sides of the leaf. Conidia elliptical brown, 5 to 6.5 x 3 to 3.5  $\mu$ .

Differs from *B. streptothrix* (C. & E.) in its much smaller conidia and more dwarfish growth.

**B. vulgaris** Fr.

On leaves *Magnolia Fraseri*, alt. 2,000 ft., July 12, 1893; on burrs of *Xanthium Canadense*, Nov. 1, 1893; on pedicels of *Ipomœa pandurata*, Feb. 17, 1894; on decaying *Brassica oleracea*, Jan. 18, 1894; on *Cicuta maculata*, March 18, 1894 (*Nuttall*, 1116).

HYPHODERMA Fr.

H. DESMAZIERI Duby.

On living leaves *Pinus echinata*. Wood Co., near Lockhart's Run (*Millsbaugh*).

VERTICILLIUM Nees.

**V. osteophilum** E. & E., sp. nov.

TYPE HABITAT: On jaw bone of calf, in woods, Nov. 27, 1894 (*Nuttall*, discov. 1761, 647.)

Prostrate sterile hyphæ yellowish-brown, distantly septate, rough, 6 to 7  $\mu$  diameter; fertile hyphæ, loosely cespitose, erect septate, smooth, yellowish-hyaline, 7 to 15  $\mu$  diameter, the primary branches alternate, and issuing at right angles, bearing opposite or ternately-verticillate branches, the ultimate division 12 to 15 x 4  $\mu$  3 to 4 in a terminal verticil, with tips often curved and bearing 1 to 4 terminal, globose, hyaline, 3.5 to 4  $\mu$ ; conidia.

**V. puniceum** (Cke. & E.) Grev.

On dead sticks, June, 1893 (*Nuttall*, 1115).

TRICHOTHECIUM Link.

**T. roseum** (Pers.) Link.

On dead dried peach on ground; on decaying leaves *Ilex verticillata*, Sept. 14, 1893 (*Nuttall*, 1199).

CEPHALOTHECIUM Corda.

**C. roseum** Corda.

On old *Polyporus pergamenus*, Short Creek, alt. 1,300 ft., Nov. 10, 1893 (*Nuttall*, 1260, 205).

DACTYLIUM Nees.

**D. dendroides** (Bull.) Fr.

On *Polystictus versicolor*, and apparently checking its development. Oct. 12, 1893 (*Nuttall*, 1206). Spores 28 x 10  $\mu$ .

## RAMULARIA Ung.

**R. Celastri** Ell. & Mart.

On leaves of *Celastrus scandens* near the ground. \* Oct. 11, 1894. Conidia 15 to 28  $\mu$  (*Nuttall*, 1700).

**R. Tulasnei** Sacc.

On *Fragaria* cult., Oct. 27, 1894 (*Nuttall*, 1724).

**R. Taraxaci** Karst.

On *Taraxacum Taraxacum*, Oct. 20, 1894 (*Nuttall*, 1722).

## DEMATIACEÆ.

## CONIGSPORIUM Link.

**C. harknessioides** (Ell. & Holl.) Sacc.

On *Rumex acetosella*, Oct. 20, 1894 (*Nuttall*, 1727, 613).

## TORULA Pers.

**T. dimidiata** Penz.

On *Rhus hirta*, alt. 1,300 ft.; Nov. 21, 1893 (*Nuttall*, 1273, 220).

**T. herbarum** Link.

On *Asparagus officinalis*, Nov. 1, 1893 (*Nuttall*, 1240, 186).

## PERICONIA Bon.

**P. pycnospora** Fres.

On *Parthenocissus quinquefolia*, Feb. 20, 1894. Conidia 12.5 to 15  $\mu$  diameter. On *Rubus odoratus*. On *Polymnia Uvedalia*. On *Phytolacca decandra*, hyphæ 600 x 15 to 16  $\mu$  (*Nuttall*, 1413). On *Polymnia Uvedalia*, spores 12.5 to 15  $\mu$  Nov. 16, 1894 (*Nuttall*, 1743).

## GONATOBOTRYUM Sacc.

**G. maculicolum** (Wint.) Sacc.

On young sprouts *Hamamelis Virginica* 1 to 2 feet high; Short Creek, alt. 1,850 ft., Aug. 15, 1894 (*Nuttall*, 1656).

## STREPTOTHRIX Corda.

**S. atra** B. & C.

On dead limbs on ground. Conidia 6 to 8 x 4.5 to 5  $\mu$ , Dec. 16, 1893 (*Nuttall*, 1329).

## ZYGODESMUS Corda.

**Z. graminicola** E. & E.

On *Carex Fraseri*, spores 7.5  $\mu$  Feb. 11, 1894 (*Nuttall*, 1374, 334).

**Z. pannosus** B. & C.

On charred bark, May 28, 1894 (*Nuttall*, 1531, 502).



STACHYLIDIUM Link.

**S. caricinum** E. & E., Proc. Phila. Acad., 1894, 377.

TYPE HABITAT: On dead-leaves *Carex Fraseri* Feb. 11, 1894 (*Nuttall*, discov. 1375, 335).

Hyphæ fasciculate brown, septate 600 to 700 x 3  $\mu$  simple or occasionally forked above towards the tip, with short cylindrical hyaline branches opposite or in whorls of three, bearing at their tips the elliptical, hyaline 4 to 5 x 1.5 to 2  $\mu$  conidia, collected into a globose head 10 to 12  $\mu$  diameter.

FUSICLADIUM Bonord.

**F. dendriticum** (Waller.) Fckl.

On living *Pyrus Malus* leaves and fruit. Monongalia Co., near Morgantown (*Millsbaugh*).

**F. Virginiense** E. & E. sp. nov?

TYPE HABITAT: On living leaves *Aster infirmus*, June 29, 1894 (*Nuttall*, discov. 1593, 541).

POLYTHRINCIUM Kunze & Schm.

**P. trifolii** Kunze.

On *Trifolium repens*, July 4, 1894 (*Nuttall*, 1588).

CLADOSPORIUM Link.

**C. herbarum** (Pérs.) Link.

On *Polygonatum biflorum*, *Menispermum Canadense*, and leaves of *Rhus hirta*, June 28, 1894 (*Nuttall*, 1564).

**C. herbarum fasciculare** Corda.

On *Reseda* cult., Jan. 28, 1894. Conidia 15 to 18 x 6 to 7.5  $\mu$  1 to 2 septate (*Nuttall*, 1354).

**C. nigrellum** E. & E., Proc. Phila. Acad., 1893, 463.

TYPE HABITAT: On inner bark of *Robinia Pseudacacia* railroad ties, Keeney's Creek, Oct. 1893 (*Nuttall*, discov. 1227, 172).

Hyphæ densely tufted, septate, subequal, 150 to 200 x 5 to 6  $\mu$ , tufts effused, subconfluent, forming a black, velvety coat extending over the surface of the bark indefinitely, with the same habit as *Macrosporium nigrellum* C. & E. Conidia smoky-hyaline, becoming pale brown, variable in size, the smaller ones ovate, continuous or uniseptate, 6 to 8 x 5  $\mu$ , the larger ones oblong-elliptical or sub-cylindrical, 2 to 3-septate, 12 to 15 x 5 to 6  $\mu$ .

**C. Triostei** Peck.

On living leaves *Triosteum perfoliatum*, alt. 1,200 ft., July 26, 1894 (*Nuttall*, 1824).

**C. epiphyllum** (Pers.) Mart.

On dying leaves *Robinia Pseudacacia*, July 31, 1893 (*Nuttall*, 1139).

**C.** ————— sp.

On dead decorticated limb *Magnolia Fraseri*, Sept., 1895 (*Nuttall*, 1853, 739).

A most beautiful velvety black form.

**C. epimyces** Cooke.

On pileus *Polyporus varians*, Jan. 25, 1894 (*Nuttall*, 1341).

## CLASTEROSPORIUM Sz.

**C. Cornutum** E. & E., sp. nov.

TYPE HABITAT: On decaying wood, Oct. 21, 1895 (*Nuttall*, discov. 1883).

Hyphæ prostrate, septate, branched, brown, about  $6\mu$  diameter. Effused in black velvety patches 1 to 2 cm. across. Conidia in pairs or threes, horn-shaped, 10 to 14 septate, 100 to 225 x  $14\mu$ , broadest below, gradually tapering above to an obtuse point, curving outward at the base, rising and spreading out above like the horns of an ox. The conidia are but slightly constricted at the septa and are sessile on the hyphæ, appearing at first as a simple nodule or tubercle on the side of the thread.

Allied to *C. Hirudo* Sacc., but that has solitary multiseptate (55 to 65) conidia and evanescent hyphæ.

## HELMINTHOSPORIUM Link.

**H. persistens** Cooke.

On branches of *Acer* in pine woods, alt. 2,000 ft., March 9, 1894. Conidia 75 to 150 x 10 to 15  $\mu$  (*Nuttall*, 1421, 392).

**H. macrocarpon** Grev.

On decorticated limbs *Platanus occidentalis*. Oct. 6, 1895 (*Nuttall*, 1887). On *Magnolia Fraseri*, Nov. 1895.

**H. folliculatum** Corda.

On *Zea Mays*, Aug. 2, 1895 (*Nuttall*, 1835).

**H. attenuatum** Peck & Cooke.

On dead log, Short Creek, alt. 1,800 ft., June 10, 1894 (*Nuttall*, 1547).

**H. septemseptatum** Peck.

On *Magnolia Fraseri*, June 20, 1894 (*Nuttall*, 1574, 534).

**H. brachypus** E. & E. sp. nov.

TYPE HABITAT: On dry wood of old log. Oct. 8, 1895 (*Nuttall*, discov. 1873, 757).

Effused in brownish-black patches of several centimeters in extent, velutinous, thin. Prostrate hyphæ only sparingly branched, obscurely septate, crooked; fertile hyphæ cespitose, erect, 40 to 50 x 4 to 6  $\mu$ , 2 to 3 septate, abruptly constricted at the tips and subtruncate. Conidia terminal, oblong-fusoid, 5-septate

brown, not constricted at the septa, 30 to 40 x 12 to 14  $\mu$ , with a short (8 to 10  $\mu$ ) obconical, hyaline or pale-brown, persistent pedicel, which is sometimes prolonged into a hyaline thread as long as the spore. This prolongation may be something of the nature of a pith or medulla drawn out of the supporting hypha and remaining attached to the pedicel.

CERCOSPORA Fres.

**C. Armoraciæ** Sacc.

On *Roripa Armoracia*. Conidia 175 x 5  $\mu$ , Aug. 26, 1894 (*Nuttall*, 1687, 593).

**C. CAULOPHYLLI** Peck.

On living leaves *Caulophyllum thalictroides*, Grant Co., near Bayard (*Millsbaugh*).

**C. Violæ** Sacc.

On *Viola obliqua*, Aug. 10, 1895 (*Nuttall*, 1841, 731).

**C. granuliformis** Ell. & Hol.

On living leaves *Viola obliqua*, June 28, 1894 (*Nuttall*, 1565).

**C. oculata** E. & K.

On leaves *Vernonia noveboracensis*, Aug. 6, 1894 (*Nuttall*, 1682).

**C. Vernoniæ** E. & K.

On leaves *Vernonia noveboracensis*, Oct. 10, 1894 (*Nuttall*, 1698).

**C. omphacodes** Ell. & Hol.

On *Phlox amœna*, July 14, 1894 (*Nuttall*, 1605).

**C. Dirantheræ** E. & K.

On leaves of *Dianthera Americana*. Oct. 23, 1895 (*Nuttall*, 1879).

**C. avicularis** Wint.

On living leaves *Polygonum aviculare*, July 5, 1895 (*Nuttall*, 1815).

**C. dubia** (Riess) Wint.

On *Chenopodium album viride*, July 4, 1894 (*Nuttall*, 1587).

**C. beticola** Sacc.

On leaves *Beta vulgaris*, Oct. 4, 1894 (*Nuttall*, 1692).

**C. Bœhmeriæ** Peck.

On leaves *Bœhmeria cylindrica*, Oct. 5, 1894 (*Nuttall*, 1694).

**C. Acalyphæ** Peck.

On leaves *Acalypha Virginica*, Oct. 3, 1894 (*Nuttall*, 1691).

**C. Ampelopsidis** Peck.

On *Parthenocissus quinquefolia*. Conidia + 40 x 3  $\mu$ , June 30, 1894 (*Nuttall*, 1578).

**C. Menispermī** E. & H.

On leaves *Menispermum Canadense*. Conidia + 77 x 5  $\mu$  June 26, 1894 (*Nuttall*, 1562).

**C. ageratoides** E & E.

On *Eupatorium ageratoides*, Oct. 14, 1894 (*Nuttall*, 1706).

**C. anthelmintica** Atk.

On living leaves *Chenopodium anthelminticum*, Oct. 15, 1894 (*Nuttall*, 1713, 601).

**C. Kalmiæ** E. & E.

On leaves *Kalmia latifolia*, Aug. 15, 1893 (*Nuttall*, 1167).

**C. Oenotheræ** E. & E., Proc. Phila. Acad., 1894, 380.

TYPE HABITAT: On leaves *Onagra biennis*, Oct. 1894 (*Nuttall*, discov. 1704, 599).

Spots irregular, mostly elongated, grayish-brown, sub-angular, 3 to 5 x 2 to 3 mm. subconfluent. Hyphæ amphigenous, subhyaline, continuous or faintly 1 to 2 septate, 15 to 20 x 3  $\mu$  in minute scattered tufts, few in a tuft, spreading subundulate. Conidia linear or only slightly attenuated above, smoky-hyaline, nucleate and faintly 3 to 5 or more septate, 25 to 80 x 2 to 2.5  $\mu$  straight or only slightly curved.

**C. Chionanthi** E. & E., sp. nov.

TYPE HABITAT: On living leaves *Chionanthus Virginica*, Sept. 1895 (*Nuttall*, discov. 1852, 738).

Spots variable in shape, subindefinite, dark brown, grayish above, 2 to 4 mm. diameter. Hyphæ epiphyllous, cespitose, 75 to 150 x 3.5 to 5  $\mu$ , brown, septate, and subgeniculate or subundulate above. Conidia fusoid or clavate, becoming brown, 3 to 5-septate, 30 to 60 x 4 to 4.5  $\mu$ .

**C. septorioides** E. & E. sp. nov.

TYPE HABITAT: On leaves *Rubus Canadensis*, Oct. 16, 1894 (*Nuttall*, discov. 1725, 610).

Spots dirty-brown, irregular, subangular, more or less limited by the veinlets, 2 to 3 mm. diameter, definite, but without any differently colored border. Hyphæ epiphyllous, cespitose on a small tubercular base, simple, sub-entire, continuous, brownish, 20 to 25 x 3  $\mu$ . Conidia cylindrical, slightly curved, hyaline, nucleate, 35 to 63 x 2 to 2.5  $\mu$ .

Differs from *C. Rubi* Sacc. in its narrower conidia without septa, and its shorter hyphæ.

**C. Rubi** Sacc.

On leaves *Rubus Canadensis*, Nov. 1894 (*Nuttall*, 1785, 676).

**C. depazeoides** (Desm.) Sacc.

On *Sambucus Canadensis*, July 26, 1894 (*Nuttall*, 1626).

**C. SMILACIS** Thum.

On living leaves *Smilax rotundifolia*, Monongalia Co., near Camp Eden (*Millsbaugh*). On *Smilax rotundifolia* (*Nuttall*, 1573, 533).

**C. smilacina** Sacc.

On leaves *Smilax glauca*, Aug. 26, 1894 (*Nuttall*, 1653, 573).

**C. columnaris** E. & E.

On *Phaseolus vulgaris* cult. Oct. 13, 1895 (*Nuttall*, 1870).

SPORODESMIUM Link.

**S. toruloides** E. & E.

On dead twigs *Cornus florida*, June 10, 1894 (*Nuttall*, 1548, 514).

**S.** ————— sp.

TYPE HABITAT: On *Tsuga Canadensis*, Feb. 3, 1894 (*Nuttall*, discov. 331).

**S. moriforme** Peck.

On *Opulaster opulifolius*, May 12, 1894. Spores 25 to 40 x 25  $\mu$  (*Nuttall*. 1503, 483).

**S. aurantiacum** B. & C.

On a dead stick, Short Creek, July 25, 1893 (*Nuttall*, 1132, 115).

**S. concinnum** Berk.

On wet dead sticks in association with *Clavaria mucida*, Oct. and Nov., 1893. Conidia 112 to 115 x 30 to 32  $\mu$  65 to 100 x 30  $\mu$  (*Nuttall*, 1233).

SPEIRA Corda.

**S. minor** Sacc.

On *Pinus Virginiana*, Dec. 15, 1894. Conidia 30 x 10  $\mu$  (*Nuttall*, 1775, 664).

MACROSPORIUM Fr.

**M. commune** Rabh.

On dead stems of garden *Asparagus officinalis*, conidia + 40 x 15  $\mu$  Nov. 1, 1893 (*Nuttall*, 1222). On petioles of *Aralia spinosa*, conidia 75 x 20  $\mu$  (*Nuttall*, 358). On *Rumex obtusifolius*, *Rhus radicans* and *Cicuta maculata*? (*Nuttall*).

**M.** ————— sp.

TYPE HABITAT: On dead stems *Polymnia Uvedalia*, Nov. 3, 1893 (*Nuttall*, discov. 1736, 623).

**M. Abutilonis** Speg.

On *Abutilon Abutilon*, July 21, 1895 (*Nuttall*, 1836).

**M.** ————— sp.

TYPE HABITAT: On decayed fruit *Diospyros Virginiana*, April 25, 1895 (*Nuttall*, discov. 1825, 711).

**M. asclepiadeum** Cooke?

On *Asclepias Syriaca*, Nov. 3, 1894 (*Nuttall*, 1730).

**M. caudatum** C. & E.

On dead flowers of *Yucca* cult., Sept. 5, 1894 (*Nuttall*, 1680, 587). On stems of *Phytolacca decandra*, conidia 88 x 20  $\mu$ .

**M. Saponariæ** Peck.

On *Saponaria officinalis*, Oct. 6, 1894 (*Nuttall*, 1695).

**M. olivaceum** E. & E., Proc. Phila. Acad., 1894, 383.

TYPE HABITAT: Parasitic on *Sphaeropsis Asiminae* on dead limbs of *Asimina triloba*, March, 1894 (*Nuttall*, discov. 1418, 388).

Forms a light olive velutinous coat over the pustules of the *Sphaeropsis*. Hyphæ tufted, yellowish brown (under the microscope), septate, erect, nearly straight or subundulate, 80 to 100 x 4 to 5  $\mu$ . Conidia obovate or obpiriform, 3 to 5-septate and muriform, brown, 24 to 38 x 15 to 20  $\mu$  terminal, sessile. Conidia also occur subcubical or subglobose 15 to 20  $\mu$  diameter with 2 septæ crossing each other at right angles.

**M. antennæforme** B. & C.

On *Celtis occidentalis*, Aug. 11, 1894 (*Nuttall*, 1652). Conidia + 95 x 15  $\mu$ .

**M.** \_\_\_\_\_ sp.

TYPE HABITAT: On pods of *Datura Stramonium*, Feb. 10, 1894 (*Nuttall*, discov. 1369, 326).

**M. TOMATO** Cooke?

On ripe fruit *Lycopersicum esculentum* cult. Spores shorter than described. Sept. 9, 1893 (*Nuttall*, 1195). Same host Monongalia Co., at Morgantown, prevalent 1891 (*Millsbaugh*).

**M. Maydis** C. & E.

On leaves *Zea Mays* cult., with conidia smaller than described, Nov. 1893 (*Nuttall*, 1334, 292).

## TRICHÆGUM Corda.

**T. nodulosum** E. & E., Proc. Phila. Acad., 1894, 385.

TYPE HABITAT: On dead leaves *Carex Fraseri*, Feb. 11, 1894 (*Nuttall*, discov. 1373, 333).

Erumpent, tufted, becoming subeffused, black, tufts gregarious, forming subvelutinous patches 2 to 4 mm. across, or when standing singly the hyphæ and conidia forming a compact mass .5 to 1 mm. diameter, and resembling somewhat the sorus of a *Puccinia*. Hyphæ simple sparingly fasciculate, brown, septate, often swollen at the septa, about 4  $\mu$  thick and 200 to 300  $\mu$  long. Conidia near the base of the hyphæ, at first elliptical, yellowish-hyaline, uniseptate, 8 to 10 x 6 to 7  $\mu$ , soon becoming 4 to 6-septate, muriform and opaque, 10 to 25  $\mu$  diameter, subglobose, obovate, or elliptical.

## SEPTOSPORIUM Corda.

**S. EUISETI** PECK, Rep. State Bot. N. Y., 1892, 25.

TYPE HABITAT: Tips of living leaves *Equisetum arvense*, Doddridge Co., near Center Point, and Monongalia Co., on College campus, Morgantown (*Millsbaugh*, discov. 1891).

Hypæ forming minute tufts, the fertile very short, bearing

acrogenous spores, the sterile longer, septate, colored; spores elliptical, usually with three transverse septæ and one or two longitudinal ones, colored, .001 in. long, .005 in. broad.

SARCINELLA Sacc.

**S. heterospora** Sacc.

On *Cercis Canadensis*, Oct. 16, 1894 (*Nuttall*, 1726, 610).

STILBACEÆ.

STILBUM Tode.

**S. magnum** Peck.

In cracks of bark, Nov. 6, 1893. Spores  $2.5 \times 1.25 \mu$  (*Nuttall*, 1278, 225).

**S. flavipes** Peck.

On dead driftwood *Platanus occidentalis*, Dec. 12, 1894 (*Nuttall*, 1767).

**S. erythrocephalum** Ditm.

On dung of Rabbit, Jan. 18, 1894 (*Nuttall*, 1345, 304).

**S. vulgare** Tode?

On fallen dead leaves, Short Creek, July 25, 1893 (*Nuttall*, 1137, 120).

PILACRE Fries.

**P. gracilipes** E. & E., Proc. Phila. Acad., 1894, 386.

TYPE HABITAT: On rotten wood, Short Creek, alt. 1,750 ft., Dec. 16, 1893 (*Nuttall*, discov. 1219, 274).

Scattered, stem slender, white-pruinose, 3 mm. long, .25 to .33 mm. thick. Head hemispherical, olivaceous, about 1 mm. diameter. Fertile hyphæ hyaline, dichotomously branched, 2 to  $2.5 \mu$  thick, bearing the conidia laterally. Conidia globose or subglobose, yellow-brown under the microscope, 4 to  $5.5 \mu$  diameter.

Smaller and of a more slender growth than *P. Petersii* B. & C.

**P. Petersii** B. & C.

On *Acer saccharum* and *Ilex opaca*, Feb. 10, 1894 (*Nuttall*, 1371).  
On *Magnolia Fraseri*, Nov. 1895.

ISARIA Pers.

**I. Virginiensis** E. & E., Proc. Phila. Acad., 1893, 465.

TYPE HABITAT: On the young stroma of *Hypoxylon rubiginosum*?  
June 30, 1893 (*Nuttall*, discov. 1109, 95, 102).

Stromata gregarious, simple, slender-clavate, 1.5 to 2 mm. high. Yellowish-white, obtuse and subcapitate at the apex, curved, often decumbent, clothed nearly to the base with spreading, hyaline, dendroid, 1.5 to  $2 \mu$  branching hyphæ (sporophores)  $40$  to  $45 \times 2.5 \mu$  their tips often toothed and bearing 1 to 4 elliptical or ovate, hyaline  $3.5 \times 2.5 \mu$  conidia.

Analogous to *I. umbrina* Pers (*Institale acariforme* Fr.), but differs in several respects.

**I. clavata** Ditm.

On dead sticks on ground, June, 1893 (*Nuttall*, 1110).

## CERATIUM A. &amp; S.

**C. hydroides** (Jacq.) Alb. & Schw.

On rotten log, June 10, 1893 (*Nuttall*, 976).

## SPOROCYBE Fr.

**S. Rhois** (B. & C.) Sacc.

On *Rhus copallina*, March 24, 1894, spores  $7.5 \times 2$  to  $2.5 \mu$ .

On *Rhus hirta*, March 28, 1894, spores  $10 \times 2.5 \mu$  (*Nuttall*, 1430).

**S. Azaleæ** (Peck) Sacc.

On capsules of *Rhododendron maximum*, Dec. 12, 1893. Conidia varying from globular through elliptical to cylindrical, dark brown, outer coat easily ruptured,  $10 \times 10$ ,  $8 \times 5$ ,  $12 \times 5$ ,  $15 \times 7.5 \mu$  (*Nuttall*, 1307).

## TUBERCULARIACEÆ.

## TUBERCULARIA Tode.

**T. VULGARIS** Tode.

On twigs, Feb. 23, 1893. On *Robinia Pseudacacia*, Dec. 1, 1893. On *Aralia spinosa*, Dec. 13, 1893. On *Sambucus pubens*. On *Acer Negundo*, April 20, 1894 (*Nuttall*, 942). On limbs of dead *Rhus hirta*, Monongalia Co., near Morgantown, 1891 (*Millsbaugh*).

**T. hamata** E. & E., Proc. Phila. Acad., 1894, 386.

TYPE HABITAT: On dead limbs of *Celtis occidentalis*. Feb. 2, 1894 (*Nuttall*, discov. 1357, 313).

Depressed hemispherical, umbonate, soon becoming black, .5 to 1.5 mm. diameter, Conidia oblong, slightly curved, hyaline, 5 to  $8 \times 1.5$  to  $2 \mu$  on slender simple sporophores 30 to  $40 \mu$  long, incurved or involute at the tips.

**T.** ——— sp.

On *Asimina triloba*, Feb. 22, 1894. Flesh colored spores 18 to  $22 \times 3 \mu$  (*Nuttall*, 1391, 361).

**T.** ——— sp.

Conidia of *Nectria verrucosa*. On dead twigs *Morus rubra* July 25, 1895 (*Nuttall*, 1827, 715).

**T. Sambuci** Corda.

On *Sambucus pubens*, Feb. 26, 1894 (*Nuttall*, 1409).

**T. Celastris** Schw.

On *Celastrus scandens*, April 25, 1895. Spores 5 to  $6 \times 2 \mu$  (*Nuttall*, 1811).

## DENDRODOCHIUM Bon.

**D. affine** Sacc.

On bark wet dead stumps, Oct. 24, 1893 (*Nuttall*, 1229, 174).



**D. rubellum microsporium** Sacc.

On dead limb *Magnolia acuminata*, Aug. 16, 1894, Short Creek, alt. 1,000 ft. On *Liriodendron Tulipifera* (Nuttall, 1659).

TUBERCULINA Sacc.

**T. persicina** (Ditm.) Sacc.

Parasitic on *Uredo* (Cæoma) *nitens*, that on *Rubus Baileyanus* June 10, 1894 (Nuttall, 1557, 527).

ILLOSPORIUM Mart

**I. cæspitosum** E. & E., Proc. Phila. Acad., 1894, 385.

TYPE HABITAT: On rotten log, Nov. 15, 1893 (Nuttall, discov. 1286, 231).

Sporodochia globose, 100 to 110  $\mu$  diameter, cespitose, forming tufts about 1 mm. diameter. Hyphæ 6 to 8  $\mu$  thick, branched, the branches curved or tortuous. Conidia globose or ovate 6 to 12  $\mu$  diameter.

Differs from *I coccinellum* Cke. in its color, and larger cespitose sporodochia.

HYMENULA Fr.

**H. cerealis** E. & E., Proc. Phila. Acad., 1894, 386.

TYPE HABITAT: On wheat straw *Triticum* sp. brought in from Painesville, Ohio, May 24, 1894 (Nuttall, discov. 1520, 495).

Sporodochia gelatinous, orbicular, yellowish-amber color becoming darker, at first sub-pulvinate, becoming depressed or flattened, .5 to .75 mm. diameter. Basidia slender, 25 to 30 x 1.25  $\mu$  simple or oftener branched. The branches erect. Conidia hyaline, oblong, minute, 3 to 4 x 1 to 1.25  $\mu$ .

CYLINDROCOLLA, Bon.

**C. DENDROCTONI** Peck, Flora, W. Va., 1892, 516.

TYPE HABITAT: On dead insects, *Dendroctonus frontalis*, beneath the bark of pine. Hampshire Co., near Romney, (Millsbaugh).

The insects are probably killed by this fungus, as they lie dead in their burrows in the inner bark of the tree (*Pinus Virginiana*).

Sporodochia minute, forming irregular masses, soft, somewhat waxy, white or whitish; sporophores slender, abundantly branched above, often compacted below into a short stem-like base, spores catenulate, short cylindrical, subtruncate, colorless, .00016 to .0002 in. long, .00008 to .0001 in. broad.

On some of the insects there is a cottony or flocculent mass of white mycelium interwoven in a somewhat reticulate manner, and collected in strings or bundles. It bears no fruit but is probably a luxuriant growth of the mycelium of this fungus.

Occasionally the fungus seems to spread from the insect to bark immediately adjacent to it.

**C. flagellaris** E. & E. sp. nov.

TYPE HABITAT: On dead stem of *Helianthus decapetalus* Dec. 3, 1894 (*Nuttall*, discov. 1762, 650).

(*C. lactea* S. & E. partly, Sacc. No. 3550.)

Sporodochia gregarious, subglobose, subhyaline and gelatinous when fresh, lens-shaped, with a thin spreading margin, orbicular, .33 to .50 mm. diameter yellowish-amber color (whitish in the center) when dry. Sporophores densely fasciculate, flagelliform 2 to 3 times dichotomously branched, 80 to 100  $\mu$  long, separating into the short cylindrical, hyaline, 4 to 5 x 1  $\mu$  conidia.

This was included by Saccardo in *Michelia* II, p. 581, in *Cylindrocolla lactea*, S. & E. (on rotten wood of *Kalmia*), but differs in the shape and color of the sporodochia and the sporophores only 2 to 3 times dichotomously branch. *C. lactea* retains its white color when dry, and resembles young *Lastosphaeria ovina*.

## SCORIOMYCES Ell. &amp; Sacc.

**S. Cragini** E. & S.

Under loose bark of dead *Hicoria* sp., Short Creek, alt. 1,250 ft., Nov. 21, 1893 (*Nuttall*, 1272, 219):

## VOLUTELLA Tode.

**V. ciliata** (A. & S.) Fr.

On decaying leaves of *Prunus* cult. in grass, June 20, 1893 (*Nuttall*, 1010).

## BACTRIDIIUM Kunze.

**B. FLAVUM** K. & S.

Under decaying bark of *Quercus alba*, Monongalia Co., near Morgantown (*Millspaugh*).

## HELISCUS Sacc.

**H. Lugdunensis** Sacc.

On dead limbs of *Ilex opaca*, thrown in wet places one year previously, March 16, 1894; sporodochia + 2 mm. diameter, conidia 30 to 33 x 5  $\mu$  (*Nuttall*, 1427, 400).

## FUSARIUM Link.

**F. CULMORUM** Smith.

On heads of ripe living wheat. Monongalia Co., Laurel Point (*Millspaugh*).

"This specimen combines the characters of a number of so-called species, making it difficult to say which it really is. Probably they are all forms of one species." Professor Peck (in letter).

**F. sarcochroum** (Desm.) Sacc.

On *Acer Negundo*, April 24, 1894 (*Nuttall*, 1482, 462).

**F. roseum** Link.

On follicles of *Asclepias Syriaca*, May 31, 1894. Conidia  $55 \times 4 \mu$  + 6-septate (*Nuttall*, 1530, 503).

**F. SOLANI** Mart.

Found associated with "black rot" on Tomato fruits that have fallen to the ground. Monongalia Co., at Morgantown, 1891 (*Millsbaugh*).

**F. roseum** — Var. nov?

On dead fruit of *Diospyros Virginiana*, Aug. 1895 (*Nuttall*, 1831, 721).

**F.** ————— sp.

On stems of *Asparagus officinalis*, Oct. 28, 1893. Color, light pink. Conidia oblong to obovate  $7.5$  to  $10 \times 2.5 \mu$  (*Nuttall*, 1239, 1248, 185).

MICROCERA Desm.

**M. erumpens** E. & E., Proc. Phila. Acad., 1894, 386.

TYPE HABITAT: On dead limbs of *Tsuga Canadensis*, March, 1895 (*Nuttall*, discov. 1398, 371).

Sporodochia scattered, depressed-globose, .5 mm. diameter, at first covered by the epidermis, raising it into little whitish pustules, then erumpent and closely embraced by the ruptured epidermis, at first orange-red, then becoming nearly black, and finally leaving subcupuliform cavities in the bark, when dry. Conidia falcate to fusiform, multinucleate, and finally three or more septate,  $60$  to  $83 \times 3$  to  $4.5 \mu$  hyaline, narrowing to a slender point at each end, borne on short sporophores ( $20$  to  $35 \mu$ ), which are more or less branched above.

Differs from *M. coccophila* Desm. in the shape of the sporodochia and their subcuticular origin.

(The additional description, incorporated in this relation of the specific characters, is by E. & E.)

EPICOCCUM Link.

**E. purpurascens** Ehrenb.

On cardboard box in grass, April 13, 1894 (*Nuttall*, 1453).

**E. NEGLECTUM** Desm.

On living leaves of *Avena sativa* and *Catalpa Catalpa*, Monongalia Co., at Morgantown, 1891 (*Millsbaugh*).

**E. Durianum** Mont.

Underside of outer bark of *Robinia Pseudacacia*, Nov. 10, 1893 (*Nuttall*, 1285, 229).

EPIDOCHIUM Fries.

**E. melanochlorum** Desm. ?

On *Carex Fraseri*, Feb. 11, 1894 (*Nuttall*, 1376, 337).

## TRIMMATOSTROMA Corda.

**T. Americana** Thum.

On *Salix nigra*, March 21, 1894 (*Nuttall*, 1437, 410).

## HYPHELIA Fries.

**H. terrestris** Fr.

On damp ground. June 18, 1893 (*Nuttall*, 1016).

## SPHÆROPSIDÆ.

## SPHÆRIOIDAECÆ.

## PHYLLOSTICTA Pers.

**P. Paviæ** E. & E.

On *Aesculus octandra hybrida* and *Hamamelis Virginica*, June 10, 1894 (*Nuttall*, 1539). (*P. sphærospidea* E. & E.)

**P. caryigena** Sacc.

*P. Caryæ* E. & E. On living leaves of *Hicoria avata* and *H. microcarpa*, June 30, 1894 (*Nuttall*, 1580).

**P. Celtidis** E. & K.

On living leaves of *Celtis occidentalis*, Oct. 10, 1894 (*Nuttall*, 1715, 603).

**P. Hamamelidis** Pk.

On living leaves of *Hamamelis Virginica*, June 10, 1894 (*Nuttall*, 1556).

**P. Catalpæ** E. & M.

On living leaves of *Catalpa Catalpa*, July 25, 1895 (*Nuttall*, 1623).

**P. Sanguinariæ** Wint.

On living leaves of *Sanguinaria Canadensis*, June 29, 1894 (*Nuttall*, 1567).

**P. Ipomœæ** E. & K.

On *Ipomœa pandurata*, Sept. 19, 1895. Spores 5 to 10 x 2 to 2.5  $\mu$  (*Nuttall*, 1861, 749).

**P. Haynaldi** Roum.

On *Ilex verticillata*. Oct. 5, 1895. Spores 5 to 8 x 2.5 to 3  $\mu$  (*Nuttall*, 1872, 756).

**P. Ribis** E. & E. sp. nov.

TYPE HABITAT: On leaves of *Ribes* cult. Oct. 21, 1895 (*Nuttall* discov., 1882, 768).

Spots large, irregular in shape, .5 to 1 cm., or often confluent along the margin of the leaf for 2 cm., rusty brown, becoming whitish or grayish, and mostly zonate, definite but without any differently colored border. Perithecia epiphyllous, scattered, 150 to 200  $\mu$  diameter, the apex prominent and black. Sporules oblong-elliptical, hyaline, granular and nucleate, 15 to 22 x 6 to 9  $\mu$ .

Distinguished from *P. ribicola* Fr., and *P. Grossularie* Sacc. by its much larger sporules.

**P. Galacis** (Cke).\*

On leaves of *Galax aphylla*, May 4, 1894 (*Nuttall*, 1486, 474).

**P. Ipomœæ** E. & K.

On *Ipomœa pandurata*, Sept. 19, 1895. Spores 5 to 10 x 2 to 2.5  $\mu$  (*Nuttall*, 1861, 749).

**P. Haynaldi** Roum.

On *Ilex verticillata*, Oct. 5, 1895. Spores 5 to 8 x 2.5 to 3  $\mu$  (*Nuttall*, 1872, 756).

**P. Ampelopsidis** E. & M.

On *Parthenocissus quinquefolia*, June 18, 1893 (*Nuttall*, 1014).

**P. ASIMINÆ** E. & K.

On living leaves *Asimina triloba*, Monongalia Co. near Camp Eden, 1891 (*Millsbaugh*). On same host June 1, 1894, spores 8 to 10 x 6  $\mu$  (*Nuttall*, 1517).

**P. Quercus-prini** E. & E., Proc. Phila. Acad., 1894, 356.

TYPE HABITAT: On leaves *Quercus Prinus*, June 29, 1894 (*Nuttall*, discov. 1594, 542).

Spots orbicular, rusty brown, definite, with a very narrow border, 2 to 3 mm. diameter. Perithecia epiphyllous, scattered, erumpent, 90 to 110  $\mu$  diameter. Sporules ovate or oblong, hyaline, mostly a little curved, 5 to 6.5 x 2 to 2.5  $\mu$ .

This comes very near *Ph. Ludoviciana* E. & E., but in that species the spots are larger, perithecia more prominent below, and sporules rather larger. *Ph. marginalis* E. & E. also has similar sporules, but the perithecia are hypophyllous.

**P. macrospora** E. & E., Proc. Phila. Acad., 1894, 355.

TYPE HABITAT: On leaves of *Liriodendron Tulipifera*, Aug. 7, 1894 (*Nuttall*, discov. 1654, 574).

Spots few, suborbicular, ferruginous-gray, 3 to 5 mm. diameter, with a narrow, raised, darker border. Sporules oblong elliptical hyaline, nucleate, 15 to 30 (mostly 20 to 25) x 6 to 7  $\mu$ .

**P. pirina** Sacc.

On leaves *Pyrus Malus*, at Blue Sulphur Springs, July 25, 1894 (*Nuttall*, 1624).

**P. Rosæ** Desm.

On *Rosa humilis*. Spores 6 to 7.5 x 2.5  $\mu$  Sept. 18, 1895 (*Nuttall*, 1863).

**P. gentianicola** (DC).

On leaves of *Gentiana Andrewsii*, Sept. 19, 1895. Spores oval, 7.5 x 5  $\mu$  (*Nuttall*, 1857).

**P. rhoicola** E. & E., Proc. Phila. Acad., 1894, 356.

TYPE HABITAT: Rare, on leaves *Rhus radicans*, Aug. 18, 1894 (*Nuttall*, discov. 1668, 567 in part).

\* *Phoma Galacis* Cke., Messrs. Ellis & Everhart decide from our specimens that this species belongs in *Phyllosticta*.

Spots 4 to 6 mm. diameter, deciduous, grayish-white, with a narrow black border. Perithecia epiphyllous, scattered, 100 to 110  $\mu$  diameter, convex-prominent above, visible also below. Sporules narrow-elliptical, hyaline, 2-nucleate, 5 to 6 x 2 to 3  $\mu$ .

Differs from *P. toxica*, E. & M., in its larger deciduous spots with a narrow border, and its narrow-elliptical sporules.

**P. lindericola** E. & E., Proc. Phila. Acad., 1894, 354.

TYPE HABITAT: On leaves *Benzoin Benzoin*, Aug. 26, 1894 (*Nuttall*, discov. 1688, 593).

Spots various; punctiform and minute without any definite border, often irregular in shape 1 mm. to 1 or 1.5 cm. diameter, or marginal, forming a narrow strip along the edge of the leaf for half its length, pale brown, nearly the color of weather-beaten wood, with a dark (almost black) border. Perithecia epiphyllous, hemispherical, 100 to 120  $\mu$  diameter, covered by the blackened epidermis which is raised into pustules barely pierced at the apex. Sporules oblong-elliptical, hyaline, 4 to 7 x 2 to 3  $\mu$ .

This differs in almost every respect from *Ph. Linderæ* E. & E.

**P. Smilacis subeffusa** E. & E.

On *Smilax rotundifolia*, Sept. 21, 1894 (*Nuttall*, 1701, 594). Perithecia scattered over surface of dead dry leaves of the host.

**P. Oxydendri** E. & E., sp. nov.

TYPE HABITAT: On leaves of *Oxydendron arboreum*, Oct. 10, 1894 (*Nuttall*, discov. 1717, 605).

Spots suborbicular or elliptical, reddish-gray, 3 to 8 mm. diameter, with a narrow, slightly raised dark red border, beyond which the leaf is generally shaded reddish-purple. Perithecia epiphyllous, subapplanate, 100 to 120  $\mu$  diameter. Sporules oblong-elliptical, 2-nucleate, hyaline, 7 to 8 x 2.5 to 3  $\mu$ . The spots finally become of a lighter, dirty-white color.

**P. Celastris** E. & E., sp. nov.

TYPE HABITAT: On leaves *Celastrus scandens*, Oct., 1894 (*Nuttall*, discov. 1718, 606).

Spots mostly marginal, 8.5 to 1 cm. diameter, ferruginous, definite, with a narrow, darker border. Perithecia epiphyllous, subapplanate, 150  $\mu$  diameter. Sporules oblong, 7 to 10 x 2 to 2.5  $\mu$  hyaline, 2-nucleate.

**P. globifera** E. & E., sp. nov.

TYPE HABITAT: On leaves *Cornus florida*, Oct. 24, 1894 (*Nuttall*, discov. 1728, 615).

Spots marginal, light-brown, continuous, extending across the tip of the leaf and partly down each side, 5 to 1 cm. wide, separated from the green part of the leaf by a narrow, purple margin. Perithecia amphigenous, discoid, black, 100 to 112  $\mu$  diameter. Sporules globose or short-elliptical, hyaline, granular, 6 to 8 x 5 to 6  $\mu$ .

**P. Negundinis** Sacc.

On *Acer Negundo*, July 5, 1894. Sporules 5 to 10 x 3 to 7.5  $\mu$  (*Nuttall*, 1631, 560, 561).

**P.** ————— sp.

On *Geum Canadense*, Oct. 24, 1894 (*Nuttall*, 1733). Spores irregular 4 to 7.5 x 1.5 to 2.5  $\mu$  2-nucleate.

**P. ACERICOLA** E. & E.

On living leaves *Acer saccharinum*, Putnam Co., near Buffalo, 1891 (*Millsbaugh*). On leaves *A. rubrum*, June 24, 1894 (*Nuttall*, 1561).

**P. Xanthorrhizæ** E. & Nuttall, sp. nov.

TYPE HABITAT: On leaves *Xanthorrhiza apiifolia*, Nov. 3, 1893 (*Nuttall*, discov. 1738, 625).

Spots subelliptical, dirty-white in the center with a dark shaded margin, 3 mm. to 1 cm. long x 2 to 8 mm. wide, finally more or less deciduous. Perithecia seated on the white part of the spots, epiphyllous, subdiscoid, perforated above, 60 to 75  $\mu$  diameter. Sporules oblong-elliptical, hyaline, 2-nucleate, 5 to 6 x 2.5 to 3  $\mu$ .

Accompanied by a *Macrosporium* on the same spots.

**P.** ————— sp.

On leaves *Rubus Canadensis*, Nov. 3, 1894. Sporules variable in form, 5 x 1.25 to 2  $\mu$  (*Nuttall*, 1747, 627).

**P. cercidicola** E. & E., sp. nov.

TYPE HABITAT: On leaves *Cercis Canadensis*, Nov. 21, 1894 (*Nuttall*, discov. 1751).

Spots suborbicular, .5 to 1 cm. diameter, rusty-brown, becoming lighter, paler below, margin narrow reddish-purple. Perithecia epiphyllous, convex, suberumpent, but covered (except the apex) by the epidermis, 110 to 150  $\mu$  diameter. Sporules oblong, slightly narrowed and rounded or obtusely pointed at the ends, hyaline, 2 to 3-nucleate, 15 to 20 x 6 to 7  $\mu$ .

Differs in the character of the spots, and in the presence of a perithecium, from *Glomosporium Cercidis*, E. & E.

**P. Araliæ** E. & E., Proc. Phila. Acad., 1894, 355.

TYPE HABITAT: On leaves *Aralia spinosa*, Sept. 20, 1895 (*Nuttall*, discov. 1703, 598, 588?).

Spots suborbicular, light-brown with the margin a little darker, .5 to 1 cm. diameter. Perithecia epiphyllous, somewhat flattened, 100 to 150  $\mu$  diameter. Sporules oblong-elliptical, hyaline, 2-nucleate, 10 to 15 x 5 to 6  $\mu$ .

**P. Sassafras** Cooke.

On leaves *Sassafras Sassafras*, June 28, 1894. Sporules 5 to 7 x 2 to 3  $\mu$  (*Nuttall*, 1566).

**P. Sambuci** Desm.

On leaves *Sambucus Canadensis*, July 20, 1894 (*Nuttall*, 1614).

**P. LABRUSCÆ** Thum.

On living leaves *Vitis* cult. (Concord), Wood Co., near Lockhart's Run, 1891 (*Millsbaugh*). On leaves of *V. æstivalis* and *rupestris*, June 29, 1894 (*Nuttall*, 1569).

**P. cornicola** (D. C.) Rabh.

On *Cornus florida*, Nov. 5, 1894 (*Nuttall*, 1748, 638).

**P. Cookei** Sacc.

On leaves *Magnolia acuminata*, Aug. 9, 1894. Sporules 15 to 20 x 6 to 7.5  $\mu$  (*Nuttall*, 1655, 577).

**P. Tiliæ** Sacc.

On leaves *Tilia Americana*, Aug. 10, 1894. Sporules 7 to 8 x 2.5 to 3  $\mu$  (*Nuttall*, 1672, 579).

**P. Chionanthi** Thum.

On *Chionanthus Virginica*, Aug. 10, 1895. Spots darker than in type (from Portugal). Spores round or oval, 5  $\mu$  or 7.5 x 5  $\mu$  (*Nuttall*, 1833, 724).

**P. lirioidendrica** Sacc.

On leaves *Liriiodendron Tulipifera*, June 16 and Oct. 29, 1894. Sporules 7.5 to 10 x 2 to 3  $\mu$  (*Nuttall*, 1571). *P. Liriiodendri*, Cke.

**P. Ulmi** West?

On leaves *Ulmus pubescens*, July 9, 1894. Sporules + 16 x 7.5  $\mu$  (*Nuttall*, 1631, 563).

**P. phomiformis** Sacc.

On leaves *Quercus Prinus* and *Q. alba*, July 4, 1894. Sporules 15 to 30 x 6 to 8  $\mu$  (*Nuttall*, 1585).

**P. Castaneæ** E. & E., Proc. Phila. Acad., 1894, 357.

TYPE HABITAT: On leaves *Castanea pumila* (not on *Castanea vesca* as originally published), July 26, 1894 (*Nuttall*, discov. 1639, 570).

Spots orbicular, rust-color, with narrow dark-shaded margin. Perithecia epiphyllous, scattered on the spots, dark, semierumpent, 100  $\mu$  diameter, often collapsing. Sporules oblong, hyaline, 5 to 7 x 2 to 2.5  $\mu$ .

**P. Lappæ** Sacc.

On *Arctium Lappa*, July 20, 1894 (*Nuttall*, 1613).

**P. Dioscoreæ** Cooke.

On leaves *Dioscorea villosa*, June 29, 1894 (*Nuttall*, 1570).

**P. opaca** E. & E. sp. nov.

TYPE HABITAT: On leaves of *Ilex opaca*. Oct. 5, 1895 (*Nuttall*, discov., 1881).

Spots mostly marginal or terminal, of irregular shape, 1 to 2 cm. across, dirty-white, with a raised subferruginous narrow border. Perithecia evenly scattered, subepidermal, sublenticular, black, raising the epidermis into pustules, 200  $\mu$  diameter. Sporules elliptical, hyaline, 3 x 1.5  $\mu$ .

Differs from *P. Ilicis* E. & E. only in its minute sporules.

## PHOMA Fries.

**P.** ——— sp.

On *Opulaster opulifolius*, May 7, 1894 (*Nuttall*, 1510, 494). Spores 7 to 10 x 2.5 to 3  $\mu$ .



**P. negundinicola ramicola** E. & E., Proc. Phila. Acad., 1894, 357-

TYPE HABITAT: On dead twigs *Acer Negundo*, April, 1894 (*Nuttall*, discov. 1483; 465, 466?).

Perithecia evenly but not thickly scattered, globose, small, .25 to .33 mm. diameter, covered by the slightly ruptured epidermis, but distinctly prominent. Sporules oblong, elliptical, or ovate, hyaline, with a single large nucleus, 10 to 13 x 4 to 5.5  $\mu$ .

Differs from the typical form, on the dead peduncles, in its broader sporules, the former being but 2.5 to 3  $\mu$ .

**P. Pennsylvanica** E. & E., Proc. Phila. Acad., 1894, 357.

TYPE HABITAT: On dead limbs *Acer Pennsylvanicum*, Feb. 2, 1894 (*Nuttall*, discov. 1361, 318).

Perithecia numerous, evenly scattered, subepidermal, .33 to .5 mm. diameter, whitish inside, raising the epidermis into minute pustules, but scarcely rupturing it. Sporules subglobose, 6 to 7.5  $\mu$  diameter, nearly hyaline.

**P. Asclepiadea** E. & E., sp. nov.

TYPE HABITAT: On dead stems of *Asclepias Syriaca*, Oct. 10, 1894 (*Nuttall*, discov. 1717, 604).

Perithecia thickly but evenly scattered, subelliptical, 110 to 120  $\mu$  in the longer diameter, subcuticular, covered by the blackened epidermis, which is raised into minute pustules but scarcely ruptured. Sporules oblong-elliptical, 2-nucleate, hyaline, 5 to 6 x 2 to 2.5  $\mu$ . On curved (almost hooked) basidia 12 to 15  $\mu$  long.

Differs from the next form which occurs with it on the same stems, in its larger perithecia and sporules and its curved basidia.

**P.** ————— sp.

On dead stems *Asclepias Syriaca*, Oct. 10, 1894 (*Nuttall*).

**P. melaleuca** B. & C.

On petioles *Aralia spinosa*, April 25, 1895 (*Nuttall*, 1810).

**P. leucostoma** Lév.

On *Opulaster opulifolius*, May 13, 1894. Sporules 2.5 to 4 x 1  $\mu$  (*Nuttall*, 1506, 487).

**P. mixta** B. & C.

On fire-killed *Liriodendron Tulipifera*, Oct. 16, 1894. Sporules appendiculate at each end, 2-nucleate, 7 to 10 x 2 to 3  $\mu$ , oblong fusoid; the accompanying filiform process hooked at the apex, 20 x .75 to 1.5  $\mu$  (*Nuttall*, 1709).

**P. scabra** Sacc.

On *Platanus occidentalis*, March 30, 1894 (*Nuttall*, 1459, 434).

**P. obscurans** E. & E., Proc. Phila. Acad., 1894, 357.

TYPE HABITAT: On leaves of *Fragaria* cult., July 8, 1894 (*Nuttall*, discov. 1600, 554).

Spots 5 to 8 mm. diameter, with a ferruginous center and broad purple border, paler below. Perithecia few, scattered, convex-prominent. Sporules oblong-elliptical, hyaline 2-nucleate, 4.5 to 5.5 x 1.5 to 2  $\mu$ . Basidia simple, lanceolate-fusoid, 8 to 12 x 1.5  $\mu$ .

*Phyllosticta fragaricola* Desm. has similar sporules, but the spots are much smaller, with a white center. The basidia also indicate *Phoma* and not *Phyllosticta*.

**P. illicicola** (C. & E.) Sacc.

On leaves of *Ilex opaca*, June 18, 1893 (*Nuttall*, 1017).

**P. Spirææ** Desm?

On *Opulaster opulifolius*, May 13, 1894. Sporules 10 to 13 x 2  $\mu$  (*Nuttall*, 1508, 489).

**P. herbarum** West.

On *Onagra biennis*, March 18, 1894 (*Nuttall*, 1435, 406).

**P. Phytolaccæ** B. & C.

On stems *Phytolacca decandra*, July 31, 1893 (*Nuttall*, 1140).

**P. pedunculi** E. & E., Proc. Phila. Acad., 1894, 357.

TYPE HABITAT: On old peduncles *Magnolia Fraseri*, Jan., 1894 (*Nuttall*, discov. 1339, 297).

Perithecia scattered, flattish pustuliform, .5 to .75 mm. diameter, with a prominent papilliform ostiolum. Sporules fusoid-oblong, hyaline, 2 to 3 nucleate, 7 to 11 x 2 to 2.5  $\mu$  ends acute.

Differs from *Stagonospora pedunculi* in its larger, flatter perithecia and fusoid-oblong (not cylindrical), rather shorter sporules.

**P. uvicola** B. & C.

On *Vitis* cult. (Concord), July 4, 1893 (*Nuttall*, 1108, 94).

**P. nyssæcarpa** Cooke.

On dead limbs *Nyssa aquatica*, April 15, 1895 (*Nuttall*, 1809, 696). Sporules 10 x 3  $\mu$ .

**P. samararum** Desm.

On samaras *Acer Pennsylvanicum*, Feb. 2, 1894 (*Nuttall*, 1363, 321).

**P. smilacina** (Peck) Sacc.

On *Smilax rotundifolia*, Aug. 17, 1894. Sporules about 18 x 6 to 7.5  $\mu$  (*Nuttall*, 1660).

### AOSPHÆRIA Berk.

**A. pezizoides** E. & E., Proc. Phila. Acad., 1894, 358.

TYPE HABITAT: On decorticated trunks of *Platanus occidentalis*, *Salix nigra falcata*, *Betula nigra*, *Fraxinus* sp., and *Liquidambar styraciflua*, on abrasions made by driftwood and ice, where it seems to grow exclusively and always below high-water mark, April, 1894 (*Nuttall*, discov. 1458, 432).

Perithecia erumpent-superficial, seriate-aggregate, hemispherical, about .75 mm. diameter, with a distinct papilliform ostiolum, at length collapsing and pezizoid or discoid. Sporules elliptical or ovoid, 4 to 6 x 2.5  $\mu$  ends obtuse.

DENDROPHOMA Sacc.

**D. Therryana** Sacc. & Roum.

On *Platanus occidentalis*, April 19, 1894 (*Nuttall*, 1463).

SPHÆRONÆMA Fries.

**S. Physocarpi** E. & E., Proc. Phila. Acad., 1894, 358.

TYPE HABITAT: On dead stems *Opulaster opulifolius*, May, 1894 (*Nuttall*, discov. 1504, 484).

Perithecia gregarious, erumpent-superficial, hemispherical, 150  $\mu$  diameter, with a straight, stout beak 90 to 115  $\mu$  long. Sporules narrow-elliptical, hyaline, 2 to 3-nucleate, 7 to 9 x 2.5  $\mu$ .

**S. infuscans** E. & E. sp. nov.

TYPE HABITAT: On dead dry wood *Juglans cinerea.*, Dec. 5, 1894 (*Nuttall*, discov. 1766, 661).

Perithecia gregarious, erumpent-superficial, hemispherical, brownish-black, about .25 mm. diameter, with a short cylindrical (.33 mm.), shining-black, obtuse, perforated ostiolum. Sporules abundant, elliptical, continuous, hyaline 3 to 4 x 1.5  $\mu$  on basidia simple or branched below, about 12  $\mu$  long and .5  $\mu$  thick at the base, narrowed above.

The surface of the wood is blackened.

**S. acerinum** Peck.

On *Acer rubrum*, June 12, 1894 (*Nuttall*, 1543).

**S. corneum** C. & E.

On *Onagra biennis*, Jan. 21, 1895 (*Nuttall*, 1791, 682 in part).

**S. Magnoliæ** Peck.

On *Magnolia acuminata*, Short Creek, alt. 1,000 ft., Aug. 16, 1894 (*Nuttall*, 1658). Sporules 10 x 6  $\mu$  nucleus large.

SIROCOCCUS Preuss.

**S. Halesiæ** E. & E., Proc. Phila. Acad., 1894, 358.

On dead limbs *Mohrodendron Carolinum* (*Halesia tetraptera*), Feb. 26, 1894 (*Nuttall*, discov. 1407, 377, 376?).

Perithecia scattered, or 2 to 3-cornate, erumpent, subglobose, about .33 mm. diameter, black, with a minute papilliform ostiolum. Sporules subglobose, about 3  $\mu$  diameter, greenish-hyaline, 3 to 6-concatenate, the terminal sporule oblong and paler. Chains simple or sparingly branched, arising directly from the proliferous layer without any distinct basidia, collected in stellate clusters.

VERMICULARIA Fries.

**V. phlogina** Fairm.

On dead leaves *Phlox amœna*, Sept. 15, 1894 (*Nuttall* 1606).

**V. compacta** C. & E.

On *Rubus odoratus*, May 30, 1894; sporules 20 to 25 x 30  $\mu$  (*Nuttall*, 1513). On *Caulophyllum thalictroides* (552).

**V. petiolorum** Schw.

On *Aralia spinosa*, and on *Robinia Pseudacacia*? Feb. 17, 1894  
(Nuttall, 1388, 356).

**V. Toxica** E. & E., sp. nov?

On *Rhus radicans*, Aug. 18, 1894 (Nuttall, 1669, 569).

**V. Dematium** (Pers.) Fr.

On *Cassia Marilandica*, Oct. 28, 1893 (Nuttall, 1251, 193). On  
*Heuchera Americana*.

**V. Dematium** ————— var.

On *Onagra biennis*, March 9, 1894 (Nuttall, 1434, 405).

**V. Arctii** Schw.

On dead leaves and petioles *Arctium Lappa*, on ground, July  
21, 1894. Sporules fusiform slightly curved + or — 20 x 3  $\mu$ .  
Setæ 35 to + 125 x 3 to 5  $\mu$  (Nuttall, 1617).

**V. subeffigurata** Schw.

On fruit peduncles *Magnolia Fraseri* in association with STAG-  
NOSPORA PEDUNCULI E. & E., July 25, 1893 (Nuttall, 1136).

**V. subeffigurata scapincola** Schw.

Bases of dead leaves *Yucca filamentosa* cult., Feb. 19, 1894.  
Sporules 18 x 2.5  $\mu$ , largest setæ 200  $\mu$  (Nuttall, 1367).

**V. Trautvetteriæ** Nuttall, sp. nov.

TYPE HABITAT: On dead leaves *Trautvetteria Carolinensis*, July 15,  
1894 (Nuttall, discov. 1633).

Perithecia very small, scattered. Sporules curved, sharp  
pointed, 18 to 25 x 2.5 to 3  $\mu$ . Setæ + 100 x 10  $\mu$ .

**V. liliacearum** West.

On *Asparagus officinalis*, *Vagnera racemosa*, *Clintonia umbellulata*  
and *Iris cristata*? Nov. 1, 1893 (Nuttall, 1241, 188).

## DOTHIORELLA Sacc.

**D. Asiminæ** E. & E., sp. nov.

On *Asimina triloba*, Feb. 2, 1894 (Nuttall, 1392, 362).

Sporules 5 to 7 x 2 to 3  $\mu$  oval or oblong, some of them  
slightly curved. Perithecia suberumpent, seriate, .25 mm. diam-  
eter, white inside and out, ovoid, in series of 3 to 5, raising the  
epidermis into short ridges split along the top so as to expose the  
pallid white apices of the perithecia. Sporules oval or ellip-  
tical, hyaline, 5 to 7 x 2 to 3  $\mu$ .

**D. minor** E. & E., sp. nov.

TYPE HABITAT: On dead limbs *Liriodendron Tulipifera* April 11,  
1895 (Nuttall, discov. 1804, 691).

Perithecia seriate-subconfluent, .5 to .75 mm. diameter, irregularly  
ruptured above, white inside, erumpent, splitting the epidermis  
into short (2 to 4 mm.) longitudinal cracks. Sporules elliptical, hy-

aline continuous, 5 to 7.5 x 3 to 3.5  $\mu$ . Differs from *D. Liriiodendri*, Cke. in its much smaller sporules.

**D. glandulosa** (Cooke) Sacc?

On *Robinia Pseudacacia*, March 14, 1894 (*Nuttall*, 1423, 395).

FUSICOCCUM Corda.

**F. ilicinum** E. & E., Proc. Phila. Acad., 1894, 359.

TYPE HABITAT: On dying, transplanted, *Ilex opaca* (transplanted Dec. 28, 1893), May 24, 1894 (*Nuttall*, discov. 1521, 496).

Stromata cortical, convex, about 1 or 1.5 mm. diameter, multilocular, whitish inside, the tuberculiform apex pierced with a single pore, rupturing the epidermis and slightly raising it. Sporules fusoid, hyaline, nucleate 15 to 22 x 2.5 to 3  $\mu$ .

CYTISPORA Ehrenb.

**C. Celtidis** E. & E., Proc. Phila. Acad., 1894, 360.

TYPE HABITAT: On dead limbs *Celtis occidentalis*, Feb. 2, 1894 (*Nuttall*, discov. 1358, 314).

Stroma valsoid, flat, thin, 1.5 to 2 mm. diameter, only penetrating the surface of the bark, multilocular, gray inside, raising the bark into small pustules, and finally rupturing it, cells resembling perithecia. Sporules allantoid, 6 to 7 x 1 to 1.5  $\mu$ .

**C. Halesiæ** E. & E., Proc. Phila. Acad., 1894, 361.

TYPE HABITAT: On dead limbs of *Mohrodendron Carolinum* (*Halesia tetraptera*), June 7, 1894 (*Nuttall*, discov. 1540).

Stromata convex-conical, sunk in the bark, orbicular, about 1 mm. diameter, white inside, unilocular, the inner surface of the cavity lined with simple, straight basidia about 15  $\mu$  long, bearing the oblong-fusoid, hyaline, 2-nucleate, straight, 5 to 7 x 1 to 1.5  $\mu$  sporules, which are expelled through a single orifice perforating the raised epidermis.

This probably is the spermogonial stage of *Diaporthe Halesiæ* or *D. tetraptera*, both of which are found in company with it.

**C. leucostoma** (Pers.) Sacc.

On cultivated *Prunus domestica* and *Amygdalus Persica*, Dec. 12, 1894 (*Nuttall*, 1769).

**C. exasperans** E. & E., Proc. Phila. Acad., 1894, 360.

TYPE HABITAT: On dead limbs *Acer Pennsylvanicum*, Feb. 2, 1894, Short Creek, alt. 1,300 ft. (*Nuttall*, discov. 1366).

Stroma buried in the bark, orbicular, about 1 mm. diameter, 4 to 6-celled (at length one-celled), prolonged above into a stout, conical beak opening by a single round pore at the apex. Sporules oblong or allantoid, hyaline, 5 to 7 x 1.25  $\mu$ .

**C.** ———— sp.

On *Chionanthus Virginica*, March 28, 1894 (*Nuttall*, 1455, 427).

**C. Persicæ** Schw.

On *Amygdalus Persica* cult. "Early Rivers," June 30, 1893 (*Nuttall*, 1106).

**C. —————** sp.

On *Cercis Canadensis*, April 10, 1894 (*Nuttall*, 1471, 447). Spores 5 to 6 x .75  $\mu$ .

**C. rhoïna** Fr.

On *Rhus hirta*, June 20, 1894. Sporules 5 to 8  $\mu$  (*Nuttall*, 1551, 636).

**C. caryigena** E. & E., Proc. Phila. Acad., 1894, 359.

TYPE HABITAT: On dead limbs *Hicoria ovata*, May 31, 1894 (*Nuttall*, discov. 1524, 513).

Stroma cortical, convex, 1.5 to 2 mm. diameter, slaty-black, 4 to 6-celled, cells yellowish-white, subglobose, small. Sporules allantoid, hyaline 3.5 to 4.5 x .75 to 1  $\mu$ .

(Spermatogonia of *Valsa caryigena*, B. & C.)?

**C. Rhois-hirtæ** Nuttall, sp. nov.

On *Rhus hirta*, June 10, 1894 (*Nuttall*, 1533, 635).

This form is clearly distinct from *C. Rhoïna* Fr. Cirrus reddish yellow. Perithecia very prominent. Spores 8 to 10 x 2 to 2.5  $\mu$ .

**C. Sassafras** E. & E. sp. nov.

TYPE HABITAT: On fire-killed twigs of *Sassafras Sassafras*, Dec. 20, 1894 (*Nuttall*, discov. 1766, 666). Jamaica, Long Island (F. C. Stewart).

Stromata thickly scattered, conical, rising from an orbicular base about 1 mm. diameter, the black, shining apices erumpent and roughening the bark, at first multilocular, becoming unilocular. Sporules allantoid, hyaline, only slightly curved, 4 to 5 x 1  $\mu$  borne on simple, straight basidia about 12  $\mu$  long.

**C. betulina** Ehr?

On *Betula nigra*, Feb. 20, 1894. Cirrus yellowish, sporules, 4 to 6 x 1  $\mu$  (*Nuttall*, 1396, 367).

**C. Salicis** (Corda) Rabenh?

On *Salix nigra*, March 22, 1894 (*Nuttall*, 1429).

**C. Platani** Fckl.

On *Platanus occidentalis*, March 14, 1894 (*Nuttall*, 1424, 396)

**C. leucosperma** (Pers) Fr.

On *Carpinus Caroliniana*, Nov. 17, 1893. Sporules 5 x 1  $\mu$  (*Nuttall*, 1257).

**C. ceratophora** Sacc.

On *Fraxinus Pennsylvanica*, Oct. 17, 1893 (*Nuttall*, 1234, 163).

**C. Curreyi** Sacc?

On *Pinus Virginiana*, Dec. 13, 1844; on *Tsuga Canadensis*, April 15, 1895 (*Nuttall*, 1771, 697).

**C. coccinea** (Reb.) Fr.

On *Robinia Pseudacacia*, Feb. 2, 1894. Sporules 3 x .75  $\mu$  (Nuttall, 1355).

**C. ———** sp.

Spermogonia of *Valsa goniosoma* Sz.? On fire-killed *Liquidambar styraciflua*, Oct. 21, 1895 (Nuttall, 1884).

**C. orthospora** B. & C?

On *Robinia Pseudacacia*, April 20, 1894. Sporules irregular in shape 12 to 18 x 2.5 to 3  $\mu$  (Nuttall, 1478, 458).

**C. carposperma** Fr.

On *Asimina triloba*, Nov. 5, 1893 (Nuttall, 1292, 237).

SPHÆROPSIS Lév.

**S. Linderæ** Peck.

On dead twigs *Benzoin Benzoin*, April 8, 1895 (Nuttall, 1795).

**S. Asiminæ** E. & E., Proc. Phila. Acad., 1894, 361.

TYPE HABITAT: On dead limbs *Asimina triloba*, Feb. 24, 1894 (Nuttall, discov. 1397, 370).

Perithecia scattered or subseriate, buried in the bark which is raised into little pustules over them and soon ruptured. Perithecia small (.25 mm.) thick-walled, with an obscurely papilliform ostiolum. Sporules oblong-elliptical, brown, obtuse, 18 to 22 x 8 to 10  $\mu$ .

**S. Asiminæ fructigena** E. & E. var. nov.

TYPE HABITAT: On old dried-up fruits of *Asimina triloba*, Oct. 3, 1895 (Nuttall, discov. 1871, 755).

A fructigenous form of *S. Asiminæ* E. & E. Perithecia smaller than in the species (110 to 140  $\mu$ ) and more abundant. Sporules a little larger (20 to 24 x 9 to 12  $\mu$ ).

**S. Ampelopsidis** C. & E.

On *Parthenocissus quinquefolia*, March 30, 1894. Sporules 18 to 20 x 10 to 12  $\mu$  (Nuttall, 1460, 437).

**S. cerasina** Peck.

On *Prunus serotina*, alt. 1,200 ft., Nov. 22, 1894 (Nuttall, 1746).

**S. Physocarp**i E. & E., Proc. Phila. Acad., 1894, 361.

TYPE HABITAT: On dead *Opulaster opulifolius*, May 2, 1894 (Nuttall, discov. 1502, 482, 486).

Perithecia scattered, .33 mm. diameter, buried in the inner bark, with their apices and papilliform ostiola erumpent, white inside (at first), sometimes, as in the preceding species 2 to 3 confluent. Sporules varying from short-elliptical 12 to 15 x 10  $\mu$  to oblong or obovate-elliptical 15 to 20 x 10  $\mu$ .

Differs from *S. Nielliæ* E. & E. in its buried perithecia and much larger sporules.

**S. celastrina** Peck.

On branches *Celastrus scandens*, April 27, 1895 (Nuttall, 1812).

**S. Œnotheræ** E. & E., sp. nov.

TYPE HABITAT: On dead stems *Onagra biennis*, Jan. 21, 1895 (*Nuttall*, discov. 1792, 683).

Perithecia scattered, erumpent, superficial, globose, 200 to 225  $\mu$  diameter, with a papilliform ostiolum, occasionally subcollapsing above. Sporules oblong-elliptical, pale brown, 20 to 25 x 10 to 12  $\mu$  on stout basidia shorter than the sporules. There are indications that the sporules become finally uniseptate.

**S. Sumachi** (Schw.) C. & E.

On branches *Rhus hirta*, March 29, 1894. Sporules 20 to 26 x 8 to 12.5  $\mu$  (*Nuttall*, 1439).

**S. Sassafras** C. & E.

On *Sassafras Sassafras*, April 10, 1895. Sporules 23 x 10  $\mu$  (*Nuttall*, 1801, 688).

**S. phomatella** Peck.

On *Fraxinus Americana*, March 31, 1894 (*Nuttall*, 1441).

**S. Caryæ** C. & E.

On hickory barrel-hoop, April 9, 1894. Sporules 18 to 25 x 8 to 11  $\mu$  (*Nuttall*, 1448).

**S. Ipomœæ** E. & E., Proc. Phila. Acad., 1894, 362.

TYPE HABITAT: On dead peduncles of *Ipomœa pandurata*, Dec. 9, 1893 (*Nuttall*, discov. 1305, 249).

Perithecia scattered, ovate-globose, 250 to 300  $\mu$  diameter, the upper part prominent and closely covered by the shining black epidermis. Sporules elliptical, brown, 18 to 23 x 10 to 13  $\mu$  on pedicels of about the same length as the sporules.

**S. Menispermii** Peck.

On *Menispermum Canadensis*, Jan. 17, 1894 (*Nuttall*, 1343, 302).

## CONIOTHYRIUM Corda.

**C. Fuckelii** Sacc.

On *Parthenocissus quinquefolia*, sporules globular, 2 to 2.5  $\mu$  March 31, 1894 (*Nuttall*, 1442).

**C. concentricum** (Desm.) Sacc ?

On leaves *Yucca filamentosa*, Sept. 1, 1894 (*Nuttall*, 1187).

## HAPLOSPORELLA Speng.

**H. Celtidis** E. & E., Proc. Phila. Acad., 1894, 362.

TYPE HABITAT: On dead limbs *Celtis occidentalis*, Feb. 2, 1894 (*Nuttall*, discov. 1359, 315).

Perithecia mostly in valsoid clusters of 3 to 10, small, 200  $\mu$  white inside, slightly sunk in the inner bark, their papilliform ostiola rupturing the epidermis. Sporules elliptical, brown, continuous, 18 to 22 x 10 to 12  $\mu$  on basidia of about the same length as the sporules.



**H. Araliæ** E. & E., Proc. Phila. Acad., 1894, 362.

TYPE HABITAT: On dead limbs *Aralia spinosa*, Feb. 26, 1894 (Nuttall, discov. 1406, 375).

Stromata seriate-connate, erumpent through longitudinal cracks in the bark, and extending from 4 to 5 mm. to 2 or more centimetres. Perithecia ovate-globose, buried in the black, sub-carbonaceous stroma, 3 to 6 in each single stroma, about .33 mm. diameter. Sporules elliptical, brown, 20 to 25 x 10 to 12  $\mu$ .

This may be the pycnidia of *Botryosphaeria fuliginosa* (M. & N.).

DIPLODIA Fries.

**D. Liriodendri** Peck.

On fire killed *Liriodendron Tulipifera*, Oct. 16, 1894 (Nuttall, 1710).

**D. Salicina** Lév.

On dead twigs *Salix nigra*, March 25, 1894 (Nuttall, 1457).

**D. Cercidis** E. & E., Proc. Phila. Acad., 1894, 363.

TYPE HABITAT: On dead limbs *Cercis Canadensis*, April 9, 1894. (Nuttall, discov. 1475, 449).

Perithecia subseriate, globose, .33 to .5 mm. diameter, slightly sunk in the inner bark and splitting the epidermis with short longitudinal clefts. Sporules elliptical, 20 to 23 x 10 to 15  $\mu$  on stout basidia, uniseptate.

**D. infuscans** E. & E., Proc. Phila. Acad., 1894, 363.

TYPE HABITAT: On bark of dead limbs *Fraxinus Americana*, April 27, 1894 (Nuttall, discov. 1492, 459).

Perithecia ovate-globose, small, 110 to 150  $\mu$  diameter, thickly scattered, blackening both the outer and inner surface of the bark. Ostiolum not conspicuous, obscurely papilliform. Sporules oblong-elliptical, 12 to 15 x 8 to 10  $\mu$  scarcely constricted.

In *D. inquinans* West, the sporules and perithecia are larger, and the bark is not blackened within.

**D. atrata** (Desm.) Sacc.

On dead limbs *Acer Negundo*, April 20, 1894 (Nuttall, 1465).

**D. caryigena** E. & E., Proc. Phila. Acad., 1894, 363.

TYPE HABITAT: On dead limbs *Hicoria ovata*, May 31, 1894 (Nuttall, discov. 1525). Also Canada (Dearness).

Perithecia subseriate, sunk in the inner bark, covered by the epidermis, which is raised into pustules and ruptured, about .5 mm. diameter, black. Sporules elliptical, brown, uniseptate, scarcely constricted, 15 to 20 x 8 to 10  $\mu$  (Pycnidia of *Valsa caryigena* B. & C.)

**D. viticola** Desm.

On *Vitis*, March 26, 1894. No septum visible in sporules (Nuttall, 1443, 412).

**D. illicicola** Desm.?

On *Ilex opaca*, Dec. 20, 1894. Sporules 22 to 25 x 11 to 13  $\mu$  (Nuttall, 1777, 667).

**D. Rubi** Fr.

On *Rubus* cult., Feb. 19, 1895. Sporules 20 x 8 to 12  $\mu$  (Nuttall, 1789).

**D. inquinans** West.

On *Fraxinus Americana*, April 27, 1894 (Nuttall, 1491, 459).

**D. Juglandis** Fr.

On *Juglans cinerea*, April 9, 1894 (Nuttall, 1449).

**D. Maydis** (Berk.) Sacc.

On *Zea Mays*, Aug. 10, 1895. Spores 30 x 4  $\mu$  (Nuttall, 1840, 729).

## BOTRYODIPLODIA Sacc.

**B. acerina** E. & E., Proc. Phila. Acad., 1894, 363.

TYPE HABITAT: On dead limbs *Acer Pennsylvanicum*, Feb. 2, 1894 (Nuttall, discov. 1362, 319).

Perithecia erumpent in botryoidal clusters, often seriatly confluent for 1 or more cm., about .5 mm. diameter, white inside, flattish above, with a broad papilliform ostiolum. Sporules elliptical, brown, uniseptate, 20 to 25 x 12 to 15  $\mu$  on basidia of about the same length.

**B.** ——— sp.

Stylosporustage? of *Pseudovalsa sigmoidea* (C. & E.). On dead limbs *Quercus Prinus*, with that species. Spores brown, 1-septate 25 x 12.5  $\mu$  (Nuttall, 1829).

## ASCOCHYTA Lib.

**A. clematidina** Thum.

On *Clematis Virginiana*, Aug. 10, 1895 (Nuttall, 1842, 733).

## ACTINONEMA Fries.

**A. Rosæ** (Lib.) Fr.

On *Rosa* cult., June 27, 1894 (Nuttall, 1563).

## HENDERSONIA Berk.

**H. Lirella** Cooke.

On *Salix nigra falcata*, April 23, 1894 (Nuttall, 1480, 460). Sporules 15 x 7.5  $\mu$ .

**H. Desmazieri** Mont.

On bark of limbs *Platanus occidentalis*, April 24, 1894. Sporules + 40 x 22  $\mu$  (Nuttall, 1466).

## STAGONOSPORA Sacc.

**S. petiolorum** E. & E., Proc. Phila. Acad., 1894, 365.

TYPE HABITAT: On dead petioles *Aralia spinosa*, Feb. 17, 1894 (Nuttall, discov. 1389, 357).

Perithecia scattered, innate, small, slightly prominent and covered by the shining, blackened epidermis, 150 to 250  $\mu$  diameter,

mostly sub-elliptical. Sporules oblong, hyaline, nucleate, becoming one or more septate, 12 to 20 x 2 to 5  $\mu$ .

**S. pedunculi** E. & E., Proc. Phila. Acad., 1893, 457.

TYPE HABITAT: On old fruit peduncles *Magnolia Fraseri* (not *LIRIODENDRON TULIPIFERA*, loc. cit.). Short Creek, July 25, 1893 (*Nuttall*, discov. 1135, 119).

Perithecia gregarious, pustuliform, about .33 mm. diameter, covered by the blackened epidermis. Sporules cylindrical, 18 to 22 x 2 to 2.5  $\mu$ , multinucleate, hyaline, straight.

**S. Physocarpi** E. & E., Proc. Phila. Acad., 1894, 365.

TYPE HABITAT: On dead stems and limbs *Opulaster opulifolius*, May 12, 1894 (*Nuttall* discov. 1505, 485).

Perithecia scattered, depressed-hemispherical; 200 to 250  $\mu$  diameter, sunk in the bark, with the upper part prominent but covered by the epidermis, which is pierced by the papilliform ostiolum, white inside. Sporules linear, multiseptate, hyaline 25 to 35 x 3 to 4  $\mu$ .

**S. collapsa** (C. & E.) Sacc.

On *Acer saccharinum*, Jan. 26, 1894. Sporules 15 to 23 x 3  $\mu$  (*Nuttall*, 1353, 309).

**S.** ——— sp.

On *Chionanthus Virginica* (*Nuttall*, 1462, 440). Spores + 25 x 7  $\mu$  4-septate. This may prove to be only a stylosporous stage of some Dothidaceous species—Ellis.

CAMAROSPORIUM Schulz.

**C. Linderæ** E. & E., sp. nov.

TYPE HABITAT: On dead limbs *Benzoin Benzoin*, April 15, 1895 (*Nuttall* discov. 1808, 694).

Perithecia scattered, semierumpent .75 mm. diameter, the upper part, except the papilliform, erumpent ostiolum, covered by the closely adherent epidermis, which is raised into distinct, hemispherical pustules. Sporules oblong, 3-septate with 1 to 2 cells divided by a longitudinal septum, brown, obtuse, not constricted 12 to 18 (mostly 12 to 15) x 5 to 7  $\mu$ . The perithecia are of a firm consistence and slaty-black inside.

SEPTORIA Fries.

**S. Tecomæ** E. & E., Proc. Phila. Acad., 1894, 367.

TYPE HABITAT: On leaves *Tecoma radicans*, Aug. 11, 1894 (*Nuttall*, discov. 1671, 580).

Spots light brown (wood color), irregular in shape, small, 1.5 to 2 mm., inconspicuous and indistinctly margined. Perithecia immersed, small (65 to 70  $\mu$ ) barely visible with a lens. Sporules 40 to 50 x 2 to 2.5  $\mu$ , not strongly curved, nucleate, hyaline.

**S. Brassicæ** E. & E., sp. nov.

TYPE HABITAT: On leaves *Brassica nigra*, Nov. 26, 1894 (*Nuttall*, discov. 1759, 646).

Spots suborbicular or subangular, 3 to 4 mm. diameter, dull white with a narrow, darker and sometimes slightly raised border. Perithecia epiphyllous, numerous, sublenticular, pale brown, coarsely cellular, broadly perforated above, 100 to 115  $\mu$  diameter. Sporules numerous, curved, obtuse, continuous, hyaline, 25 to 45  $\times$  2 to 3  $\mu$ .

Closely allied to *S. Sisymbrii* Ell., but that has the spots greenish at first and never becoming more than whitish and the sporules 1 to 3-septate.

**S. RUBI** West.

On living leaves *Rubus Canadensis*, Wood Co., Lockhart's Run, 1891 (*Millspaugh*). On same host July 26, 1894 (*Nuttall*, 1627).

**S. Corni-mariss** Sacc.

On *Cornus florida*, alt. 1,700 ft., July 19, 1894 (*Nuttall*, 1612).

**S. cornicola** Desmz.

On *Cornus alternifolia*, Sewell Valley, Aug. 6, 1894 (*Nuttall*, 1647).

**S. KALMIÆCOLA** (Schw.) B. & C.

On living leaves *Kalmia latifolia*, 1891, Monongalia Co., at Camp Eden (*Millspaugh*).

**S. ochroleuca** B. & C.

On leaves *Castanea dentata*, June 30, 1894. Maximum number of perithecia in a single macula 20 (*Nuttall*, 1579).

**S. microsperma** Peck.

On *Betula lenta*, alt. 1,200 ft., Oct. 24, 1894. Sporules + 18  $\times$  .25  $\mu$  (*Nuttall*, 1722).

**S. Polymniæ** E. & E., Proc. Phila. Acad., 1894, 368.

TYPE HABITAT: On leaves *Polymnia Uvedalia*, June 29, 1894 (*Nuttall*, discov. 1595, 543).

Spots, scattered, angular, limited by the veinlets, 2 to 4 mm. diameter, dirty green. Perithecia epiphyllous, minute, 75  $\mu$  diameter, scattered, innate, inconspicuous. Sporules filiform, continuous, 35 to 50  $\times$  1 to 1.25  $\mu$ .

**S. Trautvetteriæ** E. & E., Proc. Phila. Acad., 1894, 368.

TYPE HABITAT: On *Trautvetteria Carolinensis*, July 20, 1894 (*Nuttall*, discov. 1632, 564).

Spots irregular, subangular, partly limited by the veinlets of the leaf, often elongated and acute at one end, brownish-black, with an irregularly shaped white center, which is well defined, angular, 3 to 5 mm. in the longer diameter. Perithecia epiphyllous, but also visible below, small (65 to 75  $\mu$ ), scattered, dark. Sporules abundant, nearly straight or slightly curved, continuous, 22 to 30  $\times$  2  $\mu$ .

Differs from *S. Anemonis* Desm. in its broad, dark-margined white-centered spots, and rather longer and thicker sporules.

**S. hyalina** E. & E., Proc. Phila. Acad., 1894, 368.

TYPE HABITAT: On *Viola primulæfolia*, July 26, 1894 (*Nuttall*,

discov. 1641, 572). Also on *V. blanda*, Michigan (Hicks); and on *V. lanceolata*, Massachusetts (Miss Clarke).

Spots minute (.5 to 1 mm.) white, with a dark purple-shaded border. Perithecia punctiform, black, epiphyllous, subglobose, 65 to 75  $\mu$  diameter, not abundant. Sporules filiform, nearly straight, or slightly curved, hyaline, nucleate, not visibly septate, 20 to 40 (mostly 25 to 35)  $\times$  1 to 1.25  $\mu$ .

*S. Violæ* West, has yellowish-brown perithecia on pale zonate spots with a reddish-brown border.

**S. aciculosa** E. & E.

On *Fragaria* cult., Nov. 3, 1894. Sporules 15 to 25  $\times$  .75  $\mu$  (Nuttall, 1731).

**S. Helianthi** E. & K.

On *Helianthus decapetalus*, Oct. 29, 1894. Sporules 50  $\times$  2  $\mu$  (Nuttall, 1723).

**S. Leptostachyæ** E. & K.

On leaves *Phryma Leptostachya*, July 4, 1894. Sporules 20 to 25  $\times$  1  $\mu$  (Nuttall, 1584).

**S. Nolitangere** Thum.

On *Impatiens biflora*, Aug. 5, 1894. Sporules 20 to 25  $\times$  1.5 to 2  $\mu$  (Nuttall, 1649).

**S. Violæ** West.

On *Viola sagittata*, June 30, 1893. Sporules + 25  $\times$  1.25  $\mu$  (Nuttall, 1577).

**S. Saxifragæ** Pass.

On *Heuchera Americana*, June 10, 1894. Sporules 15 to 20  $\times$  2.5 to 3  $\mu$  (Nuttall, 1544, 521).

**S. Lobeliæ** Peck.

On *Lobelia cardinalis*, June 22, 1894 (Nuttall, 1559).

**S. Scrophulariæ** Peck.

On *Scrophularia Marilandica*, June 10, 1894. Sporules 40 to 50  $\mu$  long (Nuttall, 1538).

**S. VERBENÆ** Rob. & Desm.

On leaves *Verbena urticæfolia*, Jefferson Co., near Shenandoah Junction (Millsbaugh).

**S. psilostega** E. & M.

On *Galium circæzans*, July 14, 1894 (Nuttall, 1604).

**S. Nabali** B. & C?

On *Prenanthes serpentaria*, June 8, 1894 (Nuttall, 1536).

**S. atro-purpurea** Peck.

On *Aster cordifolius*, Nov. 24, 1894. Sporules + 35 to 55  $\times$  1 to 1.5  $\mu$  (Nuttall, 1757, 644).

**S. polygonorum** Desm.

On *Polygonum Hydropiper*, June 8, 1894. Sporules 25 to 40  $\times$  1.5  $\mu$  (Nuttall, 1537).

**S. Smilaciniae** E. & M.

On *Vagnera racemosa* (SMILACINA), Aug. 14, 1894. Maculae suffused with the marginal color late in season (*Nuttall*, 1252).

## RHABDOSPORA Mont.

**R. kalmiarum** (Schw.)\*

*Sphæria Kalmianum* Schw. On *Kalmia latifolia*, Aug. 26, 1894 (*Nuttall*, 1685, 590).

## PHLYCTÆNA Mont. &amp; Desm.

**P. vagabunda** Desm.

On *Vernonia noveboracensis*, *Phytolacca decandra* and *Onagra biennis*, March 18, 1894 (*Nuttall*, 1436, 408).

**P. septorioides** Sacc.

On *Phytolacca decandra*, May 25, 1894 (*Nuttall*, 1498).

**P. Ipomœæ** E. & E., Proc. Phila. Acad., 1894, 369.

TYPE HABITAT: On calyx lobes *Ipomœa pandurata*, Dec. 9, 1894 (*Nuttall*, discov. 1305, 250).

Perithecia scattered, subcuticular, 75 to 80  $\mu$  diameter, covered by the blackened, slightly raised epidermis. Sporules linear, hyaline, curved above, 15 to 20 x 1.25  $\mu$ .

**P. arcuata** Berk.

On *Onagra biennis*, March 18, 1894. Sporules 30 to 35 x .75  $\mu$  (*Nuttall*, 1433, 403).

## GELATINOSPORUM Peck.

**G. betulinum** Peck.

On *Betula lenta*, April 12, 1895. Sporules subfiliform nearly semicircular, 32 to 40 x 2.5  $\mu$  (*Nuttall*, 1797).

## SPHÆROGRAPHIUM Sacc.

**S. hystericinum** (Ell.) Sacc.

On *Azalea viscosa*, May 4, 1894. Sporules narrowly arcuate-falcate 34 x 3  $\mu$  (*Nuttall*, 1501, 481).

**S. Fraxini** (Peck) Sacc.

On dead limbs *Fraxinus Americana*, Short Creek, alt. 1,300 ft., July 16, 1894. Sporules 50 to 60 x 2.5  $\mu$  (*Nuttall*, 1609).

## NECTRIOIDACEÆ.

## ASCHERSONIA Mont.

**A.** \_\_\_\_\_ sp.

On *Cornus florida*. Spores 3 to 5 x .75  $\mu$  (*Nuttall*, 1800, 687).

\*Transferred to *Rhabdospora* by E. & E.

LEPTOSTROMACEÆ.

LEPTOTHYRIUM Kunze & Schm.

- L.** \_\_\_\_\_ sp.  
On living leaves *Ilex opaca*, Aug., 1894 (*Nuttall*, 1679, 584).  
Maculæ white.
- L. Liriodendri** Cke.  
On dead fallen leaves *Liriodendron Tulipifera*, April 17, 1895  
(*Nuttall*, 1805).
- L. dryinum** Sacc.  
On living leaves *Quercus rubra*, alt. 1,300 ft., June 29, 1894  
(*Nuttall*, 1592, 540).
- L. Castaneæ** (Spr.) Sacc.  
On fallen leaves-*Castanea dentata*, April 17, 1895 (*Nuttall*, 1806).
- L. petiolorum** \_\_\_\_\_ var.  
On dead leaves *Magnolia Fraseri*, April 11, 1895. Sporules  
5 x 1  $\mu$  (*Nuttall*, 1802, 689).
- L<sup>p</sup> foraminulatum** Sacc. & Ell.  
On fallen leaves *Ilex opaca*, Aug. 16, 1894. Sporules 25 x 3  $\mu$   
(*Nuttall*, 1657).
- L. vulgare** (Fr.) Sacc.  
On *Dioscorea villosa*, on *Aralia spinosa*, and on *Onagra biennis*.  
Sporules 5 to 7 x .75  $\mu$  on each host, Feb. 8, 1894 (*Nuttall*, 1368,  
323).

SACIDIUM Nees.

- S. Vitis** E. & E., sp. nov.  
TYPE HABITAT: On *Vitis cordifolia*, Nov. 6, 1893 (*Nuttall*, 1254,  
198). Fungi Columbiani 290, N. A. F., 3067.  
Perithecia scattered; scutellate not perforated, 180 to 220  $\mu$   
diameter. Sporules oblong-cylindrical, straight or curved, 15 to  
20 x 4 to 5  $\mu$ , filled with small nuclei.  
In the specimens of *Sacidium viticolum*, B. & C. in Rav. F. Am.  
the sporules are ovate or elliptical, 6 to 7 x 5  $\mu$ , and the peri-  
thecia are only 75 to 100  $\mu$  diameter.

DISCOSIA Lib.

- D. Artoceras** (Tode) Fr.  
On *Castanea dentata*, Oct. 20, 1893. On *Epigæa repens*, Jan. 18,  
1894. On *Betula lutea*, *Fraxinus Pennsylvanica*, *Magnolia Fraseri*,  
*Cornus florida*, *Sassafras Sassafras*, and *Cephalanthus occidentalis*  
(*Nuttall*, 1228, 173).
- D. MACULICOLA** Ger.  
On living leaves *Disporum lanuginosum*, Grant Co., near Bayard  
(*Millsbaugh*). On *Smilax rotundifolia* and *Oxydendron arboreum*,  
July 8, 1894 (*Nuttall*, 1598).

**D. rugulosa** B. & C.

Parasitic on upper surface of the maculæ of *Phyllosticta caryigena*, July 2, 1894 (*Nuttall*, 1583).

## ENTOMOSPORIUM Lév.

## E. MACULATUM Lév.

On living leaves and fruits *Pyrus communis* cult., Monongalia Co., Morgantown (*Millsbaugh*).

## LEPTOSTROMELLA Sacc.

## L. flicina (B. &amp; C.) Sacc.

On *Dryopteris spinulosa*, May 4, 1894 (*Nuttall*, 1487).

## EXCIPULACEÆ.

## DINEMASPORIUM Lév.

**D. hispidulum** (Schrad.) Sacc.

On *Asimina triloba*, March 9, 1894. Sporules 14 to 18 x 2 to 2.3  $\mu$  (*Nuttall*, 1422, 394).

## MELANCONIACEÆ.

## HAINESIA Ell. &amp; Sacc.

**H. Rubi** (West).

On leaves of *Rubus* cult. Oct. 28 1895 (*Nuttall*, 1880).

## GLÆOSPORIUM Desm. &amp; Mont.

## G. ————— sp.

On leaves of *Cercis Canadensis* (*Nuttall*, 985). Sporules 18 x 7.5, + 20 x 7.5, 23 x 7, 25 x 6  $\mu$ .

**G. Sanguinarie** E. & E., Proc. Phila. Acad., 1894, 371.

TYPE HABITAT: On leaves *Sanguinaria Canadensis*, July 8, 1894 (*Nuttall*, discov. 1601, 555).

Spots yellow, oblong or irregular, 3 to 5 mm. diameter, situated near the apex of the leaf which is more or less uniformly blackened. Acervuli epiphyllous, numerous, innate, yellow and conspicuous. Conidia oblong, hyaline, continuous, mostly a little curved, 8 to 15 x 3.5 to 5.5  $\mu$ .

**G. aridum** E. & H.

On *Fraxinus Pennsylvanica*, June 10, 1894 (*Nuttall*, 1545, 518).

**G. Rumicis** E. & E., sp. nov.

TYPE HABITAT: On leaves *Rumex obtusifolius*, Oct. 27, 1894 (*Nuttall*, discov. 1729, 617).

Spots reddish-brown, lighter in the center, more or less distinctly zonate 3 to 4 mm. diameter, margin darker. Acervuli innate, small. Sporules erumpent above, oblong, hyaline, continuous, 5 to 8 x 2 to 2.5  $\mu$ .

The spots resemble those of *Ovularia obliqua* Cke., on *Ramularia decipiens* E. & E.



**G. Alni** E. & E., sp. nov.

TYPE HABITAT: On living leaves *Alnus rugosa*, Nov. 24, 1894 (Nuttall, discov. 1737, 624).

Spots dirty brown, suborbicular, indistinctly zonate, .5 to 1 cm. diameter, paler in the center, border concolorous, rather indefinite; acervuli epiphyllous, numerous, small, 100 to 120  $\mu$  diameter, chestnut color, becoming darker; conidia oblong-elliptical, 12 to 16 x 6 to 8  $\mu$ .

Differs from *G. cylindrospermum* Bon., on the same host, in its much broader conidia, and from *G. rubicolum* E. & E. in the different character of the spots and rather broader conidia.

**G. nervisequum** (Fckl.) Sacc.

On leaves *Platanus occidentalis*, June 10, 1894. Sporules 12.5 to 18 x 4 to 6  $\mu$  (Nuttall, 1555, 526).

**G. Robergei** Desm.

On *Carpinus Caroliniana*, July 16, 1894 (Nuttall, 1608).

**G. betularum** Ell & Mart.

On *Betula nigra*, sporules 10 x 7  $\mu$ , July 16, 1894 (Nuttall, 1607).

**G. cylindrospermum** Bon.

On leaves of *Alnus rugosa*. Spores 8 to 10 x 2 to 2.5  $\mu$  (Nuttall, 1875).

**G. venetum** Speg.

On living leaves of *Rubus strigosus*, Sept. 19, 1895 (Nuttall, 1859).

**G. rubicolum** E. E., sp. nov.

TYPE HABITAT: On leaves of *Rubus strigosus* (Nuttall, discov. 747).

Spots at first numerous, angular, small and yellowish or reddish-yellow, soon confluent forming large brown, dry, dead spots of irregular shape and indefinite outline, occupying a large part of the leaf. Acervuli not numerous, scattered, epiphyllous, prominent, resembling perithecia; conidia oblong-elliptical, 12 to 16 x 6 to 7  $\mu$ .

Differs from *G. Rubi* E. & E. in its larger, fewer black acervuli and broader conidia, which also are about twice as large as in *G. Venetum* Speg.

MYXOSPORIUM Link.

**M. luteum** E. & E., Proc. Phila. Acad., 1893, 458.

TYPE HABITAT: On bark *Hicoria ovata*, April to July, 1893 (Nuttall, discov. 1015, 79). E. & E. N. A. F. 2953. Fungi Columbiani 150.

Stroma globose-conical, light yellow, .75 to 1 mm. diameter, slightly sunk in the inner bark, unilocular and opening by a single pore. The surface of the inner bark around this pore is of a pale slate color, the colored portion definitely limited by a black line

so as to form an irregular circle about 2 mm. across, but this line does not penetrate the bark. Sporules navicular-oblong, hyaline, obtuse, 10 to 11 x 4 to 5  $\mu$  with 1 to 2 large nuclei. Basidia slender-cylindrical 15 to 20 x 1.5  $\mu$ . Mass of exuded sporules flesh color.

**M. Rhois** (B. & C.) Sacc.

On fire killed *Rhus hirta*, June 20, 1895 (Nuttall, 1575, 537).

**M. platanicolum** E. & E., Proc. Phila. Acad., 1894, 372.

TYPE HABITAT: On limbs *Platanus occidentalis*, April 24, 1894 (Nuttall, discov. 1485, 467).

Acervuli subcutaneous, vesiculoid, pale, 1 mm. diameter, raising the ruptured epidermis into pustules but not erumpent. Sporules oval or oblong-ovate, hyaline, nucleate at first, 10 to 12 x 5 to 6  $\mu$ , on stout basidia.

**M. seriatum** E. & E., Proc. Phila. Acad., 1894, 372.

TYPE HABITAT: On bark of *Acer* sp., June, 1894 (Nuttall, discov. 1552, 523).

Nuclei pallid, orbicular, about 1 mm. diameter, seated on the surface of the inner bark; surrounded by a thin layer of smoky colored radiating hyphæ, from the inner extremities of which the botuliform or oblong, 6 to 8 x 2 to 2.5  $\mu$  hyaline conidia are produced. The nuclei are seriatly arranged, and the pale flesh-colored, flattish cirrhi are erumpent through narrow, longitudinal cracks in the bark.

COLLETOTRICHUM Corda.

**C. LINDEMUTHIANUM** (S. & M.) Scrib.

On pods of *Phaseolus* cult. "Wax, Butter Bean." Monongalia Co., near Morgantown, 1891 (Millsbaugh).

**C. ————** sp.

On *Sassafras* *Sassafras*, Oct. 15, 1894. Sporules 8 to 18 x 4 to 7  $\mu$  (Nuttall, 1714, 602).

**C. lineola** Corda.

On *Zea Mays*, Aug. 10, 1895 (Nuttall, 1834, 726).

CYLINDROSPORIUM Ung.

**C. Cratægi** E. & E., Proc. Phila. Acad., 1894, 372.

TYPE HABITAT: On leaves *Cratægus* sp., July 26, 1894 (Nuttall discov. 1640, 571).

Leaves more or less mottled with rusty red, at length uniformly of this same color. Acervuli innate, erumpent on both sides, and whitening the surface of the leaf with abundantly discharged conidia, which are 75 to 100 x 3 to 3.5  $\mu$ , nearly straight or more or less undulate and curved, nucleate, and faintly 3 to 5 septate.

**C. Toxicodendri** (Curtis) E. & E. Proc. Phila. Acad., 1893, 460.

On leaves *Rhus radicans*, June 29, 1894. Sporules, largest, 50 x 3  $\mu$  (Nuttall, 1568).

LIBERTELLA Desm.

**L. FAGINEA** Desm.

On bark of dead *Fagus atropunicea*, Grant Co., near Bayard (*Millsbaugh*). On dead *Prunus* sp. cult. Aug. 10, 1895. Cirrus bright red. Sporules variously curved 15 to 25 x 1  $\mu$  (*Nuttall*, 1843, 734).

**L. ———** sp.

On felled *Robinia Pseudacacia*, April 4, 1894. Sporules 15 to 20 x .75 to 1  $\mu$  (*Nuttall*, 1470, 446).

**L. acerina** Westend.

On *Acer rubrum* or *saccharinum*, April 4, 1894. Sporules hyaline, 15 to 18 x 1  $\mu$  (*Nuttall*, 1445).

MELANCONIUM Link.

**M. pallidum** Peck.

On dead limbs *Hicoria ovata*. Sporules + 20 x 5  $\mu$ , May 31, 1894 (*Nuttall*, 1514).

**M. oblongum** Berk.

On dead limbs *Juglans cinerea*. Sporules 20 x 10 to 2  $\mu$ , Dec. 5, 1893 (*Nuttall*, 1291, 236).

**M. bicolor** Nees.

On *Betula nigra*, May 2, 1894 (*Nuttall*, 1489).

THYRSIDIUM Mont.

**T. hedericolum** *Carpini* Sacc.

On dead shoots *Carpinus Caroliniana*, May 2, 1894 (*Nuttall*, 1486).

MARSONIA Fisch.

**M. Juglandis** (Lib.) Sacc.

On *Juglans cinerea*, June 22, 1894 (*Nuttall*, 1560).

**M. Martini** Sacc. & Ell.

On *Quercus alba*, *velutina*, et *Prinus*, Aug. 18, 1894 (*Nuttall*, 1674, 583).

CORYNEUM Nees.

**C. cupulatum** E. & E., Proc. Phila. Acad., 1894, 374.

TYPE HABITAT: On dead limbs *Tsuga Canadensis*, Short Creek, Dec. 16, 1893 (*Nuttall*, discov. 1317, 272).

Erumpent superficial. Acervuli tuberculiform, black, 1 to 1.5 mm. diameter, hollowed out so as to be cup-shaped above. Conidia clavate, sessile, 6 to 9 septate, brown, 60 to 80 x 12 to 15  $\mu$ .

**C. pustulatum** Peck.

On dead limbs *Castanea dentata*, Dec. 7, 1893 (*Nuttall*, 1383).

PESTALOZZIA De Not.

**P. ———** sp.

On leaves *Geum Canadense*, Oct. 24, 1894 (*Nuttall*, 1734). Sporules 20 x 6  $\mu$ . Setæ 5 to 12  $\mu$  long.

**P. funerea** Desm.

On *Ipomœa pandurata*, Sept. 19, 1895. Spores 23 to 25 x 6 to 7.5  $\mu$  (Nuttall, 1862, 750).

**P. Guepini** Desm.

On leaves *Rhododendron maximum*, *R. Catawbiense* and *Azalia viscosa*. Spores 18 to 20 x 8  $\mu$ , Aug. 2, 1894 (Nuttall, 1149).

**P. Jefferisii** Ell?

On *Opulaster opulifolius*, sporules 10 to 15 x 4 to 5 and 25 x 5  $\mu$  May 7, 1894 (Nuttall, 1509).

**P. Toxica** E. & E., Proc. Phila. Acad., 1894, 374.

TYPE HABITAT: On leaves *Rhus radicans*, Aug. 18, 1894 (Nuttall, 1670, 567).

Spots and perithecia as in *Phyllosticta rhoicola* E. & E. Sporules clavate-oblong, 4-septate, 12 to 15 x 4 to 5  $\mu$ , 3 intermediate cells pale brown, end cells short, conical, hyaline, the upper cell with a crest of 3 short, spreading hyaline bristles 6 to 7  $\mu$  long. Distance between the two extreme cells 12  $\mu$ , pedicels shorter than the spores.

## STEGANOSPORIUM Corda.

**S. Castaneæ** Lib?

On *Castanea dentata*, Nov. 14, 1893 (Nuttall, 1275, 221).

**S. piriforme** (Hoffm.) Corda.

On dying *Acer rubrum*, June 12, 1894 (Nuttall, 1553).

**S. muricatum** Bon.

On dead limbs *Betula nigra* in drift-wood along river, Oct. 21, 1895. Spores 35 to 50 x 15 to 18  $\mu$  (Nuttall, 1878).

## USTILAGINACEÆ.

## USTILAGO Pers.

**U. SEGETUM** (Bull.) Dittm.

On living heads of wheat and oats, Monongalia Co., near Morgantown; Lewis Co., near Alum Bridge; Taylor Co., near Thornton (*Millsbaugh*).

**U?** ——— sp.

On under surface of newly fallen leaves *Asimina triloba*, Nov. 20, 1893 (Nuttall discov. 1288; 233).

**U. MAYDIS** (DC.) Corda.

On living ears and tassels of sweet corn, prevalent in Monongalia Co., 1891; near Morgantown (*Millsbaugh*). Ongarden corn, July 31, 1894 (Nuttall, 1636).

**U. Avenæ lævis**, K. & S.

On oats, Sewell Mountain, alt. 2,600 ft., Aug. 23, 1894 (Nuttall, 1678).

TILLETIA Tul.

T. TRITICI (Bjerk.) Wint.

*Ustilago tritici*, C. Bauhin. On living leaves of wheat, Monongalia Co., near Morgantown (*Millsbaugh*).

UROCYSTIS Rabenh.

U. ANEMONES (Pers.) Schrœt.

On living leaves and under stem cuticle of *Actea alba*, Monongalia Co., near Morgantown (*Millsbaugh*).

UREDINACEÆ.

UROMYCES Link.

U. **Polygoni** (Pers.) Fckl.

Uredospores on *Polygonum erectum*, Aug. 2, 1894 (*Nuttall*, 1642).

U. TRIFOLII (Hedw.) Lev.

On living leaves, *Trifolium pratense*, Mason Co., near Point Pleasant (*Millsbaugh*). On *Trifolium*, July 13, 1894 (*Nuttall*, 1604).

U. APPENDICULATUS (Pers.) Link.

On living leaves of pole beans, Monongalia Co., near Morgantown (*Millsbaugh*).

U. **Caladii** (Schwein) Farl.

On *Arisæma triphyllum* (uredospores), May 28, 1894 (*Nuttall*, 1511).

U. **Hyperici** (Schwein) Curt.

Æcidiospores on *Hypericum mutilum*, at Rupert's, July 25, 1894 (*Nuttall*, 1625).

U. **Terebinthi** (DC.)

On living leaves *Rhus radicans*, Oct. 17, 1895 (*Nuttall*, 1876).

U. LESPEDEZÆ (Schwein) Peck.

On living leaves *Lespedeza violacea*, Monongalia Co., near Morgantown (*Millsbaugh*).

U. HEDYSARI-PANICULATI (Schwein) Farl.

On living leaves *Meibomia canescens*, Mason Co., near Point Pleasant (*Millsbaugh*). On leaves *M. paniculata*, July 17, 1894 (*Nuttall*, 1610).

U. **Howei** Pk.

On *Asclepias Syriaca*, Oct. 13, 1895 (*Nuttall*, 1869).

MELAMPSORA Cast.

M. **farinosa** (Pers.) Schrœt.

On *Salix nigra*, Oct. 17, 1893. Spores at first hyaline (22 x 13  $\mu$ ) but the granules finally turn yellow (22  $\mu$ ), and perfectly fill the spore (25 x 18  $\mu$ ) (*Nuttall*, 1107).

## PUCCINIA Pers.

**P. Helianthi** Sz.

On *Helianthus decapetalus*, Sept. 19, 1895 (*Nuttall*, 1851).

**P. Violæ** (Schum.) DC.

On *Viola hastata*, æcidial stage; æcidia 18 x 25  $\mu$ . A species of small reddish caterpillar found feeding on the spores, April 30, 1893. Uredo stage on *Viola rotundifolia* or *V. blanda*, April 30, 1893 (*Nuttall*, 906).

**P. Convolvuli** (Pers.) Cast.

On *Convolvulus repens*. Æcidium May 3, Uredo June 14, Teleuto. Oct. 4, 1894; all on same vine (*Nuttall*, 1516).

**P. Convolvuli** ————— var.

On dead stems of *Convolvulus repens*, Feb. 19, 1895 (*Nuttall*, 1793).

**P. PIMPINELLÆ** (Strauss) Link.

On living leaves *Osmorrhiza Claytonii*, Monongalia Co. near Morgantown (*Millsbaugh*).

**P. Menthæ** Pers.

On *Cunila origanoides*, Nov. 27, 1894. Teleutospores 28 x 20  $\mu$ , not yet verruculose (*Nuttall*, 1753).

**P. coronata** Corda.

On *Holcus lanatus*, Uredo. June 30, 1893; teleuto. Nov. 1, 1893, on same specimen. Length of pedicel 5 to 7.5  $\mu$  very thick, corona + 38  $\mu$  broad, teeth 12.5  $\mu$ , teleutospores 50 x 15  $\mu$  (*Nuttall*, 1265).

**P. RUBIGO-VERA** (DC.) Wint.

On leaves *Triticum vulgare*, Wood Co., near Kanawha Station (*Millsbaugh*).

**P. Caricis** (Schum.) Reb.

On *Carex* sp. July 10, 1894, uredospores and teleutospores (*Nuttall*, 1590).

**P. Hieracii** (Schum.) Mart.

*P. flosculosorum* Roehl. On living leaves *Carduus lanceolatus*, Mason Co., near Point Pleasant (*Millsbaugh*). On *Taraxacum Taraxacum*, Oct. 20, 1894 (*Nuttall*, 1721).

**P. SUAVEOLENS** (Pers.) Rostr.

On living leaves *Carduus lanceolatus*, Wood Co., near Kanawha Station (*Millsbaugh*).

**P. Sorghi** Schwein.

Uredo. and teleutospores on *Zea Mays*, Sept. 7, 1894 (*Nuttall*, 1676).

**P. Conoclinii** Seymour.

On *Eupatorium caelestinum*, Sept. 18, 1895 (*Nuttall*, 1849).

**P. Anemones-Virginiana** Sz.

On *Anemone Virginiana*, Sept. 18, 1895 (*Nuttall*, 1850).

**P. Heuchera** Sz.

*P. Spreta* Pk., *P. Tiarella* B. & C. On *Tiarella cordifolia*. Oct. 21, 1895 (*Nuttall*, 1886).

**P. Smilacis** Schwein.

Uredospores on *Smilax glauca*, Oct. 14, 1894 (*Nuttall*, 1708).

**P. Podophylli** Schwein.

On *Podophyllum peltatum*, May 31, 1894 (*Nuttall*, 1515).

**P. tenuis** Burrill.

On *Eupatorium ageratoides*, June 1, 1894 (*Nuttall*, 1519).

**P. Asteris** Duby.

On *Aster cordifolius*, Nov. 24, 1894 (*Nuttall*, 1758, 645).

GYMNOSPORANGIUM, Hedw.

**G. clavariiforme** (Jacq.) Rees.

Æcidiospores on *Cratægus* sp., Aug. 2, 1894 (*Nuttall*, 1637).

**G. juniperinum** (Linn.) Fr.

Teleutospores on *Juniperus Virginiana*, April 27, 1894 (*Nuttall*, 1473).

G. MACROPUS Link.

On *Juniperus Virginiana*, Mercer Co., near Princeton (*Mills-  
paugh*). Æcidia on *Pyrus coronaria*, Aug. 2, 1893 (*Nuttall*, 1157).

PHRAGMIDIUM Link.

P. POTENTILLÆ (Pers.) Karst.

On *Potentilla Canadensis*, uredospores, Monongalia Co., near Morgantown (*Mills-  
paugh*). On *P. Canadensis*, uredo. March 15, 1894, teleutospores, Oct. 5, 1894 (*Nuttall*, 1400?).

**P. subcorticium** (Schrank.) Wint.

On *Rosa humilis*, June 18, 1893 (*Nuttall*, 1008).

COLEOSPORIUM Lev.

C. SENECONIS (Pers.) Fr.

On living leaves of *Pinus echinata* (æcidia), Wood Co., near Lockhart's Run (*Mills-  
paugh*).

C. SONCHI (Pers.) Lev.

On living leaves *Vernonia noveboracense*, Mason Co., near Point Pleasant (*Mills-  
paugh*). On *Aster cordifolius*, Dec. 3, 1894 (*Nuttall*, 1760).

**C. Ipomœæ** (Schwein) Burrill.

On *Ipomœa pandurata*, Oct. 15, 1894. Spores 85 x 23  $\mu$  (*Nutt-  
all*, 1720, 608).

**C. Vernoniæ** B. & C.

On *Vernonia noveboracense* at Rupert's, Sept. 25, 1894. Spores 100 x 30  $\mu$  (*Nuttall*, 1702, 595).

**C. Solidaginis** (Schwein.) Thum.

On *Solidago Canadensis*, June 19, 1894 (*Nuttall*, 1550).

## ÆCIDIDIUM Pers.

**Æ. Impatientis** Schwein.

On *Impatiens biflora*, June 21, 1894 (*Nuttall*, 1558).

**Æ. HOUSTONIANUM** Schwein.

On *Houstonia cærulea* (spermogonia), Monongalia Co., near Morgantown (*Millsbaugh*). On leaves of *H. longifolia*, June 1, 1894 (*Nuttall*, 1518).

## PERIDERMIIUM Lev.

**P. PECKII** Thum.

On living leaves *Tsuga Canadensis*, Pocahontas Co., near Traveler's Repose (*Millsbaugh*).

**P. BALSAMEUM** Peck.

Under surface living leaves *Abies Balsamea*, Randolph Co., Shades-of-Death (*Millsbaugh*).

## UREDO Pers.

**U. Agrimoniz** (DC.) Schroet.

On *Agrimonia striata*, July 10, 1894 (*Nuttall*, 1591).

**U. CÆOMA NITENS** Schwein.

*Cæoma nitens*. On living leaves *Rubus hispidus*, Monongalia Co., near Morgantown (*Millsbaugh*). On *Rubus villosus*, May 2, 1893 (*Nuttall*, 907).

## PERISPORIACE Æ.

## PODOSPHÆRA Kunze.

**P. OXYACANTHÆ** (DC.) De By.

On living leaves of *Prunus* cult., *Cratægus oxyacantha* and *Diospyros Virginiana*, Monongalia Co., near Morgantown (*Millsbaugh*).

**P. TRIDACTYLA** (Wallr.) De By.

On living leaves of *Prunus* cult., Cabell Co., near Huntington (*Millsbaugh*).

## SPHÆROTHECA Lev.

**S. HUMULI** (DC.) Burriil.

On living leaves *Agrimonia striata*, Preston Co., near Terra Alta (*Millsbaugh*).

**S. Castagnei** Lev.

On living leaves *Erechtites hieracifolia*, Nov. 11, 1894. Asci 78 x 58  $\mu$ , spores + 20 x 12  $\mu$  (*Nuttall*, 1740).

## PHYLLACTINIA Lev.

**P. suffulta** (Reb.) Sacc.

On living leaves *Castanea dentata*, Oct. 1893 (*Nuttall*, 1315), and on *Magnolia Fraseri*.



UNCINULA Lev.

**U. AMPELOPSIDIS**, Peck.

On cultivated grapes, Monongalia Co., near Morgantown (Millsbaugh). On leaves *Parthenocissus quinquefolia*, Sept. 20, 1894 (Nuttall, 1683).

**U. Americana** How.

On leaves *Vitis cordifolia*, Sept. 20, 1894 (Nuttall, 1684).

MICROSPHÆRA Lev.

**M. Alni** (DC.)

On leaves *Castanea dentata*, alt. 1,800 ft., Oct. 20, 1893 (Nuttall, 1218, 169).

**M. erineophila** Peck.

On leaves *Fagus atropunicea*, Dec. 20, 1894 (Nuttall, 1773).

**M. Grossulariæ** Lev.

On leaves *Sambucus Canadensis*, Oct. 5, 1894 (Nuttall, 1693).

**M. elevata** Burr. and Bess.

On leaves of *Catalpa Catalpa*, alt. 2,200 ft. Oct. 2, 1894 (Nuttall, 1690).

**M. Russellii** Clint.

On *Oxalis stricta*, Aug. 9, 1894 (Nuttall, 1650).

**M. Vaccinii** Schw.

On *Epigæa repens*, Oct. 1893. Threads slender, sharply pointed, somewhat rough, asci 5-spored, 50 to 65 x 30 to 40  $\mu$ , spores 18 to 20 x 10 to 11  $\mu$  (Nuttall, 1255).

**M. quercina** Schw.

On *Quercus palustris*, at Rupert's, Aug. 26, 1894 (Nuttall, 1686, 591).

ERYSIPHE (Hedw.) DC.

**E. Cichoracearum** DC.

On *Xanthium Canadense*, Nov. 1, 1893 (Nuttall, 1223), and on *Eupatorium purpureum*.

**E. communis** (Wall.) Fr.

On *Eupatorium ageratoides*, Oct. 14, 1894 (Nuttall, 1707).

**E. GRAMINIS** DC.

On living leaves *Poa pratensis*, Preston Co., near Terra Alta (Millsbaugh).

**E. Liriodendri** Schw.

On *Liriodendron Tulipifera* in deep shade, Short Creek, alt. 1,150 ft., Oct. 9, 1894 (Nuttall, 1697).

EUROTIUM Link.

**E. herbariorum** (Wigg) Link.

On damp leaves *Liriodendron Tulipifera* neglected in plant press, Sept. 1, 1894 (Nuttall, 1681).

## ASTERINA Lev.

**A. Leemingii** E. & E.

On *Galax aphylla*, May 4, 1894 (*Nuttall*, 1493, 475).

## DIMEROSPORIUM Fckl.

**D. Galactis** E. & E.

On leaves of *Galax aphylla* in conjunction with the last species above, May 4, 1894 (*Nuttall*, 1576, 475).

**D. Collinsii** (Schw.) Thum.

On leaves *Amalanchier Canadensis*, Glade Creek, May 4, 1894 (*Nuttall*, 1490, 472).

## SCORIAS Fries.

**S. spongiosa** (Schw.) Fr.

On living *Fagus atropunicea*, at Rupert's July 26, 1894. Spores 16 to 18 x 5  $\mu$  (*Nuttall*, 1634).

## SPHÆRIACEÆ.

## VALSA Fries.

**V. ceratophora** Tul.

On fire killed *Hicoria minima*, Oct. 17, 1894. On *Acer saccharinum*, L. Asci 38 x 5  $\mu$ , spores 7.5 to 10 x 1.5 to 1.75  $\mu$  (*Nuttall*, 1711).

**V. Diospyri** E. & E. Proc. Phil., Acad., 340 (1894).

TYPE HABITAT: On dead limbs *Diospyros Virginiana*, Dec. 10, 1893 (*Nuttall*, discov. 1308, 253).

Stroma consisting of the slightly blackened substance of the bark, convex, about 2 mm. diameter, not circumscribed. Perithecia 4 to 10 in a stroma, subglobose, .25 to .33 mm. diameter, necks converging and united above in a small, black disk which perforates the pustuliform-elevated epidermis, but does not rise above it. Ostiola short, conic-cylindrical, with a smooth, round opening crowded and finally obliterating the disk. Asci clavate-lanceolate, p. sp. 25 x 5  $\mu$ , 8-spored, paraphysate. Sporidia allantoid, hyaline, 8 to 10 x 1.5 to 2  $\mu$ . Spermogonia (Cytispora), stroma multilocular gray inside, opening by a single, central pore. Spermata allantoid, hyalins, moderately curved, 4 to 5 x 1.25  $\mu$ .

**V. etherialis** E. & E. Proc. Phila. Acad., 341 (1894).

TYPE HABITAT: On dead limbs of *Acer rubrum*, Feb. 1894 (*Nuttall*, discov. 1304, 373).

Stromata cortical, thickly scattered, convex 1 to 1.5 mm. diameter. Perithecia 6 to 12 together, circinate, buried in the unaltered substance of the bark, small, 150 to 250  $\mu$  diameter their short necks terminating in an erumpent, compact fascicle of obtuse, black, slightly umbilicate ostiola closely embraced by the epidermis and scarcely rising above it. Asci (p. sp.) fusoid, 15 to 22 x 4 to 4.5  $\mu$ , stipitate, 8-spored. Sporidia biseriata, allantoid, hyaline, curved, slender, 5 to 6 x about 1  $\mu$ . When well developed, the epider-

mis is raised into subdiscoïd pustules in which the slight protuberances indicate the position of the subjacent perithecia.

*V. delicatula* C. & E. has fewer, larger perithecia and broader sporidia. *V. microspora* Cke. & Plowr. has also larger perithecia and yellowish sporidia, and the ostiola are more or less distinctly sulcate, indicating its close relationship to *Eutypella*. In *V. etherialis* the sporidia both in and out of the asci are perfectly hyaline.

**V. coronata** (Hoffm.) Fr.

On young, fire-killed *Castanea dentata*, Nov. 14, 1893 (*Nuttall*, 1263, 209).

**V. albopuncta** E. & E., sp. nov.

TYPE HABITAT: On dead limbs *Liriodendron Tulipifera*, Oct., 1894 (*Nuttall*, discov. 1712, 600).

Stromata minute (.5 to 7.5 mm.), subseriate, included in the thick epidermis and not penetrating the inner bark, white throughout, and without any distinct circumscribing line. Perithecia 3 to 6 in a stroma, pale slate-color, 110 to 130  $\mu$  diameter; necks very slender, terminating in minute, subglobose, slate-colored ostiola tardily erumpent around the margin of the minute, snow-white, pulverulent disk. Asci clavate-cylindrical, 50 x 6  $\mu$ , short-stipitate, obtuse above, paraphysate? Sporidia biseriate, allantoid, hyaline, moderately curved, 6 to 9 x 1.5 to 2  $\mu$ .

A beautiful species. The lines of snow-white disks, which barely perforate the ruptured epidermis and are closely surrounded by its torn and slightly raised margin, present a very neat appearance.

**V. Abietis** Fr.

On *Tsuga Canadensis* Short Creek, alt. 1,100 ft., Feb. 2, 1893 (*Nuttall*, 1349).

**V. Vitis** (Schw.) Fckl.

On *Vitis* sp., March 26, 1894 (*Nuttall*, 1432).

**V. præstans** B. & C.

On dead twigs *Nyssa aquatica*, Nov. 14, 1893 (*Nuttall*, 1247). Spores 10 to 12 x 3 to 4  $\mu$ .

**V. Nyssæ** Cooke.

On fire killed *Nyssa aquatica*, Nov. 14, 1893 (*Nuttall*, 1689).

**V. subclypeata** C. & Peck.

On young fire killed *Sassafras Sassafras*, Nov. 14, 1893 (*Nuttall*, 1262, 209).

**V. ambiens** (Pers.) Fr.

On *Asimina triloba*, *Cornus florida* and *Aralia spinosa*, Feb. 18, 1894 (*Nuttall*, 1383).

**V. Chionanthi** E. & E., Proc., Phila. Acad., 340 (1894).

TYPE HABITAT: On dead limbs of *Chionanthus Virginica*, March, 1894 (*Nuttall*, discov. 1456, 228).

Perithecia 4 to 10, globose, .25 to .33 mm. diameter, buried in the unchanged substance of the bark, with convergent necks, terminating in short-cylindrical, obtuse, perforated ostiola erumpent in a close fascicle perforating and slightly raising the bark. Asci clavate, p. sp. 40 to 45 x 8 to 10  $\mu$ , 8-spored, paraphysate? Sporidia allantoid, hyaline, 12 to 15 x 3.5 to 4.5  $\mu$ , biseriate above.

Spermogonia (*Cytispora Chionanthi* E. & E.) buried in the bark flask-shaped, .5 to .75  $\mu$  diameter, multilocular, the cells soon confluent, the apex erumpent and perforated by a single pore. Sporules allantoid, 4 to 6 x 1 to 1.25  $\mu$ , borne on basidia branched above, the branches erect, straight, nucleate, 7 to 10  $\mu$  long.

**V. pauperata** C. & E.

On *Acer rubrum*, March 26, 1894 (*Nuttall*, 1444, 423).

**V. leucostoma** (Pers.) Fr.

On peach tree, Dec. 2, 1894 (*Nuttall*, 1768).

**V. Linderæ** Peck.

On *Benzoin Benzoin*, April 11, 1895 (*Nuttall*, 1796).

EUTYPELLA Nits.

**E. densissima** E. & E., Proc. Phila. Acad., 341 (1894).

TYPE HABITAT: On dead limbs *Aralia spinosa*, Feb., 1894 (*Nuttall*, 363).

Stromata scattered, cortical, depressed-conical, 2 to 3 mm. diameter, not circumscribed, but staining the bark olive-gray. Perithecia numerous, often 50 to 70 in a stroma, 100 to 120  $\mu$  diameter, closely packed, their slender necks terminating in obtusely conical, 4 cleft, black, densely crowded ostiola erumpent in a brown disk surrounded by the ruptured epidermis. The disk is soon obliterated, so that only the crowded, black, subshining ostiola are seen. Asci clavate-fusoid, p. sp. 25 to 30 x 4  $\mu$ . Sporidia biseriate, allantoid, hyaline, moderately curved, 8 to 10 x 1.5 to 2  $\mu$ .

**E. rugiella** (C. & E.) Sacc.

On *Acer rubrum*, May 4, 1893 (*Nuttall*, 921).

**E. stellulata** (Fr.) Sacc.

On *Robinia Pseudacacia*, March 29, 1893 (*Nuttall*, 33).

**E. Platani** Sz.

On *Platanus occidentalis* in drift. Spores 7.5 to 8 x 1.5 to 2  $\mu$ . Oct. 21, 1895 (*Nuttall*, 1885).

EUTYPA Tul.

**E. spinosa** (Pers.) Tul.

On dry dead logs, June 29, 1893 (*Nuttall*, 1105).

**E. milliaria** (Fr.) Sacc.

On river drift wood, April 20, 1894. Asci 150 x 6  $\mu$  spores 10 x 2  $\mu$  (*Nuttall*, 1477).

DIATRYPE Fr.

**D. Stigma** (Hoffm.) Fr.

On *Castanea dentata*, April 15, 1895 (*Nuttall*, 1799).

**D. platystoma** (Schw.) Berk.

On *Magnolia Fraseri*, Feb. 8, 1894 (*Nuttall*, 1380, 342).

DIATRYPELLA C. & DeNot.

**D. verruciformis** (Ehrh.) Nits.

On *Alnus rugosa*, Nov. 20, 1893 (*Nuttall*, 1258).

ROSELLINIA DeNot.

**R. aquila** (Fr.) De Not.

On dead limbs, March 21, 1893 (*Nuttall*, 919, 19).

**R. corticium** (Schw.) Sacc.

On dead oak, alt. 1,800 ft., Short Creek. Spores 25 to 30 x 10 to 12.5  $\mu$ . Dec. 16, 1893 (*Nuttall*, 1310).

**R. subiculata** (Schw.) Sacc.

On dead dry *Robinia Pseudacacia*, July 15, 1893. Perithecia clustered or connate, each 3 to 5  $\mu$ , black, ostiola papillate. Spores 9 to 12.5 x 5 to 6  $\mu$  somewhat inequilateral, elliptical, colored. (*Nuttall*, 1129).

**R. pulveracea** (Ehrh.) Fckl.

On dead wood, Nov. 1893. Spores 10 to 15 x 7 to 9  $\mu$  (*Nuttall*, 1338).

**R. millegrana** (Schw.) Sacc.

On dead *Platanus occidentalis*, Aug. 2, 1893 (*Nuttall*, 1161).

**R. abietina trichota** C. & Ell.

On *Pinus Virginiana*, Jan. 7, 1895 (*Nuttall*, 1784, 675).

BOMBARDIA Fr.

**B. fasciculata** Fr.

On wet dead limb, *Magnolia Fraseri*, on ground, Sep. 16, 1895 (*Nuttall*, 1830).

ANTHOSTOMA Nits.

**A. microecium** E. & E., Proc. Phila. Acad., 344 (1894).

TYPE HABITAT: On dead limbs *Asimina triloba*, Feb. 12, 1894 (*Nuttall*, discov. 1377).

Stroma cortical, faintly circumscribed, 1 to 1.5 mm. diameter, orbicular, convex. Perithecia 4 to 8 in a stroma, globose, minute (200 to 250  $\mu$ ), circinate, necks slender, short, converging, with the minute papilliform ostiola erumpent in a small, black, hemispherical disk, which barely pierces the pustuliform-elevated epidermis, and is closely embraced by it. Asci cylindrical, 80 to 110 x 8 to 10  $\mu$ , paraphysate 8-spored. Sporidia uniseriate, elliptical, brown, continuous, 2-nucleate, 12 to 14 x 6 to 7  $\mu$ .

Distinguished by its very small stroma and perithecia.

**A. microplacum** (B. & C.) Sacc.

On *Sassafras Sassafras*, alt. 1,800 ft., March 21, 1893 (*Nuttall*, 915).

XYLARIA Hill.

**X. polymorpha** (Pers.) Grev.

On dead logs, April 7, 1893 (*Nuttall*, 890).

**X. corniformis** Fr.

On rotten limb, Sept. 9, 1893. On *Magnolia Fraseri*, Nov. 1895 (*Nuttall*, 1194).

**X. Hypoxylon** (Linn.) Grev.

On dead log, March 10, 1893 (*Nuttall*, 807).

**X. flabelliformis** (Schw.) B. & C.

On *Carpinus Caroliniana*, May 13, 1893 (*Nuttall*, 957).

**X. Cornu-Damæ** (Schw.) Berk.

On wet rotten log, alt. 2,000 ft., Aug. 24, 1893. Spores 20 to 21 x 4.5  $\mu$  (*Nuttall*, 1179).

USTULINA Tul.

**U. vulgaris** Tul.

On dead and rotting stumps, March 8, 1893 (*Nuttall*, 954).

HYPOXYLON Bull.

**H. atroviride** E. & E., Proc. Phila. Acad., 346 (1894).

TYPE HABITAT: On bark of dead tree, *Betula* or *Quercus*, Dec., 1893 (*Nuttall*, discov. 1320, 275).

Stroma pulvinate, 1 to 1.5 cm. across, and about 4 mm. thick, orbicular, covered above with a thin (.5 mm.) carbonaceous crust, which is soon covered by a dark green layer of the ejected spores, laterally and internally dirty-umber color. Perithecia (which constitute the entire inner substance of the stroma) ovate compressed, and including the long, stout neck, about 3 mm. long and 1 mm. broad below. Ostiola papilliform, soon covered and obscured by the ejected sporidia. Asci cylindrical 100 x 3.5 to 4  $\mu$  (p. sp. 40 to 45 long), paraphysate, 8-spored. Sporidia uniseriate, oblong-elliptical, pale brown under the microscope, 2-nucleate, 4.5 to 5.5 x 2  $\mu$ .

This is a well-marked species. The substance of the stroma, except the superficial carbonaceous layer, is friable, and in this respect as well as the color resembles *H. Petersii* B. & C., from which, however, in other respects it is very distinct.

**H. coccineum** Bull.

On *Aralia spinosa*, alt. 1,300 ft., Aug. 18, 1893 (*Nuttall*, 1173, 140).

**H. Nuttallii** E. & E., Proc. Phila. Acad., 346 (1894).

TYPE HABITAT: On bark of dead *Magnolia Fraseri* at Glade Creek, May 1894 (*Nuttall*, discov. 1494, 477).

Stromata gregarious, subconfluent, depressed-hemispherical or strongly convex, 3 to 6 mm. diameter, purplish-black, mammillose. Perithecia small (about .25 mm.) scattered irregularly through the stroma, which is rather soft and brownish within. Ostiola crowning the mammillose projections on the surface of the stroma, papilliform, soon deciduous, leaving a round perforation. The asci in the spec. examined had disappeared. Sporidia brown, oblong-elliptical, sub-inequilateral, 7 to 8 x 3 to 4  $\mu$  (exceptionally 9 x 5  $\mu$ ).

**H. Howeianum** Peck..

On decorticated *Magnolia Fraseri*, alt. 1,800 ft., Sept. 12, 1893 (*Nuttall*, 1203, 158).

**H. FUSCUM** (Pers.) Fr.

On dead and decorticated *Acer saccharum*, Grant Co., near Bayard (*Millsaugh*). On dead *Alnus rugosa*, Oct. 31, 1893. Spores 12 x 6 to 7  $\mu$  (*Nuttall*, 1221).

**H. multiforme** Fr.

On dead *Betula* sp., alt. 1,800 ft., Short Creek, Aug. 11, 1893 (*Nuttall*, 1147).

**H. perforatum** (Schw.) Fr.

On dead twig, Feb. 3, 1894. On *Ilex opaca*, Dec. 28, 1894 (*Nuttall*, 1351). Spores 10 x 5  $\mu$ . On *Magnolia Fraseri*, Nov. 1895.

**H. rubiginosum** (Pers.) Fr.

On dead hard wood of *Robinia Pseudacacia*, Feb. 25, 1893 (*Nuttall*, 949, 39).

**H. Sassafras** (Schw.) Berk.

On *Sassafras Sassafras*, March 21, 1893. Spores 8 to 12 x 3 to 4  $\mu$  (*Nuttall*, 917).

**H. stigmatum** Cooke

On living *Quercus* sp., originating under the outer bark which it pushes off, or at times remains attached at one edge and stands away like a lid. Asci, total length, 150 x 10  $\mu$ , fertile portion 120 x 10  $\mu$ , spores black, elliptical, 16 to 20 x 6 to 10  $\mu$ . Feb. 2, 1894 (*Nuttall*, 1350).

**H. colliculosum** (Schw.) Nits.

On *Rhododendron maximum*, alt. 1,800 ft., Short Creek, Dec. 16, 1893 (*Nuttall*, 1330, 289).

DALDINIA DeNot.

**D. concentrica** (Bolt.) C. & DeN.

On bark of dying *Acer* sp., April 15, 1893 (*Nuttall*, 905).

**D. vernicosa** (Schw.) C. & DeN.

On dead trees, March, 1895 (*Nuttall*, 1790).

## NUMMULARIA Tul.

**N. punctulata** (B. & Rav.) Sacc.

On dead branch of *Quercus* sp., Aug. 16, 1893 (*Nuttall*, 1151).

## GNOMONIELLA Sacc.

**G. Coryli** (Batsch.) Sacc.

On *Corylus Americana* at Rupert's, Aug. 5, 1894 (*Nuttall*, 1647).

## PHYSALOSPORA Niessl.

**P. Ilicis** (Schl.) Sacc.

On dead leaves of *Ilex opaca*, July 21, 1894. Asci 7.5 to 10  $\mu$  (*Nuttall*, 1616).

## TRICHOSPHERIA Fckl.

**T. pulchriseta** Peck.

On dry firm splinters of dead log, alt. 1,800 ft., Short Creek, March 6, 1894 (*Nuttall*, 1412, 382).

## BOTRYOSPHÆRIA Ces. &amp; DeNot.

**B. Quercuum** (Schw.) Sacc.

On *Quercus* sp., *Vitis* sp., *Pinus Virginiana*, *Opulaster opulifolius*, and *Parthenocissus quinquefolia*. Perithecia 2 to 10 (*Nuttall*, 925, 14).

**B. Quercuum** ————— var.

On cultivated *Rubus*, Feb. 25, 1895 (*Nuttall*, 1794, 685).

**B. Araliæ** Curtis.

On *Aralia spicosa*, Feb. 24, 1894. Spores 20 to 30 x 8 to 10  $\mu$  (*Nuttall*, 1395).

**B. pyriospora** (Ellis) Sacc.

On *Chionanthus Virginica*, March 28, 1894. Asci, fertile portion, 75 x 15  $\mu$  (*Nuttall*, 1438).

**B. Hibisci** ————— forma.

On *Celastrus scandens*, April 27, 1895 (*Nuttall*, 1819, 707). Spores 26 to 35 x 6 to 7.5  $\mu$ .

## CRYPTOSPORELLA Sacc.

**C. compta** (Tul.) Sacc.

On dead limbs of *Fagus atropunicea*, Nov. 17, 1893. Asci, fertile portion, 100 x 12.5 to 15  $\mu$ , spores granular 22 x 10  $\mu$ . (*Nuttall*, 1256).

## SPHÆRELLA Ces. &amp; DeNot.

**S. maculiformis** (Pers.) Awd.

On leaves of *Castanea dentata* on ground, Dec. 13, 1893 (*Nuttall*, 1314).



**S. conicola** Sacc.

On dead cones of *Tsuga Canadensis*, Dec. 16, 1893. Spores 10 x 3  $\mu$  (Nuttall, 1311).

**S. Gaultheriæ** C. & P.

On *Gaultheria procumbens*, June 8, 1894 (Nuttall, 1534).

**S. nyssæcola** Cooke?

On *Nysa aquatica*, Oct. 10, 1894 (immature) (Nuttall, 1719, 607).

DIDYMELLA Sacc.

**D. Physocarpi** E & E., Proc. Phila. Acad., 1894, 335.

TYPE HABITAT: On *Opulaster opulifolius*, May 12, 1894 (Nuttall, 1507, 488).

Perithecia gregarious, covered by the pustuliform epidermis, about .25 mm. diameter, white inside, depressed-globose, the papilliform ostiolum barely penetrating the epidermis. Asci clavate-cylindrical, short-stipitate, 85 to 90 x 10 to 12  $\mu$ , paraphysate, 8-spored. Sporidia biseriata, fusoid, 4-nucleate, uniseptate, constricted at the septum, hyaline, 19 to 22 x 4 to 5  $\mu$ , mostly a little curved.

MELANCONIS Tul.

**M. modonia** Tul.

On *Castanea dentata*, Nov. 14, 1893 (Nuttall, 1274, 221).

HERCOSPORA Tul.

**H. Tiliæ** (Fr.) Tul.

On *Tilia heterophylla*, Feb. 16, 1894. Spores 18 x 7.5  $\mu$  (Nuttall, 1834, 346).

DIAPORTHE Nits.

**D. Gladioli** E. & E. ?

On *Caulophyllum thalictroides*, July 8, 1894. Asci 38 x 5 to 7  $\mu$ , spores septate, 4-nucleate, about 10 x 3  $\mu$  (Nuttall, 1603, 553).

**D. Araliæ** E. & E., Proc. Phila. Acad., 339 (1894).

TYPE HABITAT: On *Aralia spinosa*, Feb. 2, 1894 (Nuttall, discov. 1356, 312).

Stroma buried in the wood and circumscribed by a penetrating, black line, elliptical, about 5 x 4 mm. Perithecia buried in the stroma, few (6 to 10) globose, .5 to .75 mm. diameter, their short-cylindrical ostiola projecting from a black, tubercular disk seated on the surface of the wood and perforating the pustuliform-elevated epidermis, but scarcely rising above it. Asci clavate-cylindrical, 40 to 45 x 5 to 6  $\mu$ , paraphysate, 8-spored. Sporidia biseriata, oblong, 4-nucleate, becoming uniseptate and slightly constricted, hyaline, obtuse, 12 to 13 x 2.5 to 3  $\mu$ .

Allied to and resembling *D. ocularia* C. & E.

**D. CHOROSTATE Halesiæ** E. & E., Proc. Phila. Acad., 339 (1894).

TYPE HABITAT: On dead limbs of *Mohrodendron Carolinum*, May, 1894 (Nuttall, discov. 1541).

Perithecia 4 to 8, loosely circinate. .75 mm. horn-color inside, becoming nearly black, sunk in the wood, necks converging with their obtuse, smooth, hemispherical ostiola erumpent in a close fascicle, closely surrounded by the ruptured epidermis. Asci slender, 55 to 60 x 6 to 7  $\mu$ , short-stipitate. Sporidia subbiseriate, fusoid-oblong, hyaline, 2 to 4-nucleate, becoming uniseptate and slightly constricted, straight, 12 to 15 x 2.5 to 3  $\mu$ .

There is no distinct circumscribing line around the stroma, but the inner surface of the bark is uniformly blackened.

**D. CHOROSTATE *Tetrapteræ* E. & E., Proc. Phila. Acad., 339 (1894).**

TYPE HABITAT: On dead limbs *Mohrodendron Carolinum*, May, 1894 (*Nuttall*, discov. 1542).

Stroma cortical, 1.5 to 2 mm. diameter, surrounded by a black line which does not penetrate the wood. Perithecia 4 to 12, circinate, 5 mm. diameter, sunk to the wood and leaving their impress on its surface but not penetrating it. Ostiola obtusely rounded and perforated, erumpent in a compact fascicle and closely surrounded by the ruptured epidermis. Asci clavate-cylindrical, 75 to 80 x 12  $\mu$ . Sporidia biseriate, oblong-fusoid, slightly curved, subobtuse, hyaline, uniseptate and constricted at the septum, each cell with a large nucleus, 9 to 22 x 5 to 7  $\mu$ .

Differs from *D. Halesiæ* E. & E., in its smaller perithecia, not sunk in the wood, and its much larger sporidia.

**D. CHOROSTATE *Aceris* Fckl.**

On *Acer saccharinum*, L., Jan. 26, 1894 (*Nuttall*, 1739).

**D. *Hydrangeæ* E. & E., sp. nov.**

TYPE HABITAT: On dead stems *Hydrangea arborescens*, Nov. 1894 (*Nuttall*, discov. 1756, 641).

Perithecia scattered, ovate-globose, 350 to 380  $\mu$  diameter, sunk in the wood, which is not at all discolored. Ostiola stout, erumpent, short-cylindrical, or conic-cylindrical, the apex papilliform. Asci (p. sp.) oblong-cylindrical, 50 to 60 x 8  $\mu$ , 8-spored, obscurely paraphysate. Sporidia biseriate, oblong elliptical, hyaline, uniseptate, slightly constricted, 12 to 15 x 3 to 4  $\mu$ , ends subacute.

This may be the *Sphæria* spoken of by Schweinitz as found by him on limbs of *Hydrangea*, in company with his *Sphæria* (*Teichospora*) *Hydrangeæ*.

**D. CHOROSTATE *cercophora* (Ellis.) Sacc.**

On *Ilex opaca* Dec. 23, 1893, March 15, 1894 (*Nuttall*, 1333).

**D. CHOROSTATE *oncostoma* (Du B.) Fckl.**

On *Robinia Pseudacacia*, April 7, 1894. Spores 17 to 19 x 4 to 5  $\mu$  (*Nuttall*, 1446).

**D. CHOROSTATE *sociata* (C. & E.) Sacc.**

On *Benzoin Benzoin*, April 13, 1895. Helminthospores 75 to 100 x 15  $\mu$ . 7-septate (*Nuttall*, 1798).

**D. CHOROSTATE *obscura* (Peck) Sacc.**

On *Rubus villosus*, Feb. 13, 1893 (*Nuttall*, 1313, 260).

**D. EUPORTHE aculeata** (Schw.) Sacc.

On *Phytolacca decandra*, Nov. 14, 1894. Asci 40 to 55 x 6 to 7.5  $\mu$ . Spores 12.5 to 18 x 3 to 3.5  $\mu$  (Nuttall, 1754, 639).

**D. TETRASTAGA rostellata** (Fr.) Nits.

On *Rubus odoratus*, May 31, 1894 (Nuttall, 1528, 560).

AMPHISPHERIA Ces. & DeNot.

**A. pinicola** Rehm.

On *Pinus rigida* at Glade Creek, and on *Pinus Virginiana*, May 4, 1894. Spores very variable + 35 x 12  $\mu$  (Nuttall, 1499, 480).

VALSARIA Ces. & DeNot.

**V. exasperans** (Ger.) Sacc.

On *Juglans cinerea*, Nov. 3, 1893 (Nuttall, 1260, 202).

MASSARIA DeNot.

**M. Magnoliæ** E. & E., sp. nov.

TYPE HABITAT: On bark of dead limbs *Magnolia acuminata*, April 25, 1895 (Nuttall, discov. 1818, 700, 705).

Perithecia scattered, buried in the bark, depressed-globose, .75 to 1.25 mm. diameter, the subconical ostiolum piercing but not perceptibly raising the bark. Asci broad oblong, p. sp. 150 to 200 x 55 to 65  $\mu$  with a very short, nodular stipe, 8-spored, paraphysate. Sporidia crowded-biseriate, oblong or clavate-oblong, 3 to 5-septate, mostly 3-septate, when young, hyaline at first, finally deep brown and then with only one distinct septum, though often 1 to 4 other faint septa can be seen. There is a distinct constriction at the main septum, which is a little below the middle of the spore, but none at the other faint septa which are often wanting. The sporidia measure 65 to 80 x 20 to 25  $\mu$ , larger than in *M. fœdans* (Tode), which has sporidia about 50 x 20  $\mu$ , 3-septate and constricted at all the septa, besides being more distinctly narrowed below.

LEPTOSPHERIA (Pers.) DeNot.

**L. Doliolum** (Pers.) DeNot.

On *Polymnia Uvedalia* and *Helianthus decapetalus*, Nov. 16, 1894. Perithecia covered by outer bark. Asci, fertile part 110 x 6  $\mu$ , stipitate, spores overlapping at ends. Spores 22 x 4  $\mu$ , 23 x 4.5  $\mu$ , 25 x 5  $\mu$ , 27 x 5  $\mu$ , 30 x 5  $\mu$ ; 3-septate, nucleate, constricted at septum when mature; straight, one-sided, or curved (Nuttall, 1744).

**L. vagabunda** Sacc.

On dead stems *Hydrangea arborescens*, Nov. 21, 1894. Asci 125 to 150 x 7.5  $\mu$  total length (Nuttall, 1745).

**L. ogilviensis** (B. & Br.) Ces. & DeNot.

On *Cimicifuga racemosa*, July 4, 1894 (Nuttall, 1586).

**L. clavigera** (C. & E.) Sacc.

On old stems *Phytolacca decandra*, Nov. 14, 1894 (*Nuttall*, 1741).  
Spores +  $43 \times 6 \mu$ .

**L. orthogramma** (B. & C.)

On *Zea Mays*, Aug. 10, 1895 (*Nuttall*, 1832, 722).

## MELANOMMA Nits &amp; Fckl.

**M. Pulvis-pyrius** (Pers.) Fckl.

On decorticated *Magnolia Fraseri*, alt. 1,800 ft., Short Creek,  
Sept. 10, 1893. Spores  $15 \times 3.5 \mu$  (*Nuttall*, 1200).

## TREMATOSPHERIA Fckl.

**T. vitigena** E. & E., sp. nov.

TYPE HABITAT: On old, decaying wood of *Vitis rupestris*, April,  
1894 (*Nuttall*, discov. 454).

Perithecia gregarious, sunk in the wood all except the obtuse,  
convex apex, .5 to .75 mm. diameter, depressed-globose, the  
buried part rather thin-walled, the erumpent, convex apex thick,  
solid, like the stromatic shield of *Clypeosphaeria*. Ostium papil-  
liform, soon perforated. Asci clavate-cylindrical, paraphysate,  
8 spored, 86 to 100  $\times$  10 to 12  $\mu$ . Sporida sub-biseriate, fusoid-  
oblong, 3-septate, and constricted, subacute, pale-brown, 20 to  
25  $\times$  6 to 7  $\mu$ .

## PSEUDOVALSA Ces. &amp; DeNot.

**P. sigmoidea** (C. & E.)

On dead limbs *Quercus Prinus*. Spores 50 to 88  $\times$  5 to 7  $\mu$   
(*Nuttall*, 1828, 717).

## LASIOSPHERIA Ces. &amp; DeNot.

**L. ovina** (Pers.) Ces. & DeNot.

On *Juglans cinerea*, and parasitic on *Hypoxylon*, Short Creek,  
alt. 1,800 ft., Dec. 16, 1893. Spores 50 to 60  $\times$  6 to 7.5  $\mu$ .  
Asci 135  $\times$  12.5  $\mu$  (*Nuttall*, 1322).

## ACANTHOSTIGMA DeNot.

**A. decastylum** (Cooke) Sacc.

On dead log, June 18, 1893 (*Nuttall*, 1018).

## HERPOTRICHIA Fckl.

**H. diffusa** (Schw.)

On dead twig, Sept. 10, 1893 (*Nuttall*, 1197).

## CALOSPORA Sacc.

**C. aculeans** (Schw.) Sacc.

On *Rhus hirta* and *copallina*, March 29, 1894. Spores 15 to 17  
 $\times$  3  $\mu$ , aculeate (*Nuttall*, 1440).

**C. Rhoina** (C. & Ell.) Sacc.

On *Rhus hirta*, Dec. 9, 1893. Spores 15 to 18 x 5?  $\mu$  hyaline, 4-nucleate, uniseptate constricted at septum. (*Nuttall*, 1348, 306).

PYRENOPHORA Fr.

**P. calvescens** (Fr.) Sacc.?

On *Chenopodium anthelminticum*, Aug. 10, 1895 (*Nuttall* 1845, 736).

TEICHOSPORA Fckl.

**T. nitida** E. & E.

On *Rubus villosus*, Apr. 25, 1895. Asci 75 x 12?  $\mu$ . Spores 20 to 23 x 9 to 10  $\mu$  (*Nuttall*, 1817, 706).

PLEOSPHÆRIA Speg.

**P. corticola** E. & E., Proc. Phila. Acad., 1894, 332.

TYPE HABITAT: On bark of *Pinus rigida*, Glade Creek, June 24, 1894 (*Nuttall*, discov. 1582).

Perithecia scattered, superficial, ovate, 300 to 400  $\mu$  diameter, carbonaceo-membranaceous, pilose-strigose, hairs 100 to 200 x 5 to 6  $\mu$ , soon opaque, very faintly and sparingly septate, here and there collected into closely compacted fascicles resembling stout bristles. Ostiolum papilliform, obtuse. Asci oblong-cylindrical, abruptly short-stipitate, 80 to 100 x 20 to 25  $\mu$ . Paraphysate? Sporidia crowded, acutely elliptical, nearly hyaline at first, becoming olive-brown and 5 to 7 septate and muriform, but not constricted.

Comes near *P. strigosa* Sacc., but perithecia rather smaller, not depressed, and hairs fasciculate.

**P.** ————— sp.

On *Tecoma radicans*, Feb. 20, 1894 (*Nuttall*, 1414, 385).

CUCURBITARIA Gray.

**C. Fraxini** E. & E.

On *Fraxinus Americana*, Apr. 27, 1894 (*Nuttall*, 1523).

**C. elongata** (Fr.) Grev.

On *Robinia Pseudacacia*, Jan. 15, 1894 (*Nuttall*, 1783).

OPHIOBOLUS Riess.

**O. porphyrogonus** (Tode) Sacc.

On *Vernonia noveboracensis*, Apr. 24, 1895 (*Nuttall*, 1820, 702).

**O. acuminatus** (Sow.)

On *Zea Mays*, Aug. 10, 1895 (*Nuttall*, 1839, 728).

**O. fulgidus** (C. & P.) Sacc.

On *Vernonia noveboracensis*, Apr. 24, 1895 (*Nuttall*, 1821, 703).

## CRYPTOSPORA Tul.

**C. femoralis** (Peck) Sacc.

On *Alnus rugosa*, Feb. 25, 1894. Asci 60 to 75 x 12.5 to 15  $\mu$ . Spores 50 to 65 x 3.5  $\mu$  in middle and enlarging to 5  $\mu$  at each end (*Nuttall*, 1318).

**C. cinctula** (C & P) Sacc?

On *Castanea dentata*, Dec. 8, 1893. Asci about 75 x 7.5  $\mu$ . Spores 60 to 75 x 4  $\mu$ , hyaline, nucleate, curved, often slightly thickened toward one end (*Nuttall*, 1296, 245).

**C. trichospora** (C. & P.) Sacc?

On *Quercus* sp., Feb. 1894. Spores +50?  $\mu$  (*Nuttall*, 1450).

## HYPOCREACEÆ.

## HYPOMYCES Fr.

**H. rosellus** (Alb. & Schw.) Tul?

On *Hymenocetum* on *Magnolia Fraseri*, March 6, 1894 (*Nuttall*, 1417, 387).

**H. aurantius** (Pers.) Fckl.

On *Stereum* sp. Sept. 3, 1893 (*Nuttall*, 1198). Short Creek, alt. 1,800 ft.

## NECTRIA Fr.

**N. cinnabarina** (Tode) Fr.

On *Aralia spinosa*, March 5, 1894. Asci 75 x 7.5  $\mu$ . Spores 20 to 23 x 4 to 5  $\mu$  (*Nuttall*, 1405, 374).

**N. coccinea** (Pers.) Fr?

On bark of dead *Magnolia Fraseri*, March 1894. Asci 65 x 7.5  $\mu$ . Spores 10 to 15 x 3.5 to 6  $\mu$  (*Nuttall*, 1472, 451).

**N. ditissima** Tul.

On dead and fallen *Robinia Pseudacacia*, on *Magnolia Fraseri*, Oct. 17, 1893 (*Nuttall*, 1115, 162).

**N. vulpina**, Cke.

On bark of dead *Ulmus pubescens*? Oct. 17, 1894. Spores 10 x 5  $\mu$ , 1-septate, 2-guttulate (*Nuttall*, 1216, 164).

**N. episphæria** (Tod.) Fr.

On dead fallen twigs, Aug. 15, 1893 (*Nuttall* 1175, 143).

## APONECTRIA Sacc.

**A. inaurata** (B. & Br.) Sacc.

On *Rhus copallina*, Mar. 24, 1894 (*Nuttall*, 1431). Microspores 3 x 1  $\mu$ . Spores 10 to 12.5 x 3  $\mu$ .

## HYPOCREA Fr.

**H. Virginiensis** E. & E. Proc. Phila. Acad., 442 (1893).

TYPE HABITAT: On leaves of *Rhododendron maximum*, Aug. 12, 1893 (*Nuttall*, discov. 1163).

Epiphyllous, stroma carnose, scattered, sub-hemispherical or depressed-turbinate, 1 to 2 mm. diameter, of a yellowish gray color, subtruncate above and obscurely margined (when mature). Perithecia buried in the stroma, ovate, membranaceous, 110 to 150  $\mu$  diameter. Ostiola hemispherical, prominent, black, broadly perforated and sometimes collapsed. Asci clavate-cylindrical, p. sp. 40 to 45 x 7 to 8  $\mu$ , short-stipitate, filiform-paraphysate, 8-spored. Sporidia biseriata, oblong-fusoid, yellowish-hyaline, uniseptate, slightly constricted at the septum, obtusely pointed at the ends, 10 to 12 x 3 to 3.5  $\mu$ .

*H. viridans* B. & C. seems to differ in its hidden ostiola (ostiolis latitantibus).

**H. tenerrima** E. & E., Proc. Phila. Acad., 442 (1893).

TYPE HABITAT: Incrusting moss, leaves, twigs, living *Gaultheria procumbens*, etc. July 1893 (*Nuttall*, discov. 1138, 123).

Perithecia gregarious, minute (.16 mm.), clothed (except the black apex) with a thin white tomentum, seated on a thin snow-white, tomentose-arachnoid subiculum. Asci cylindrical, 40 x 3  $\mu$ , without paraphyses, 8-spored. Sporidia uniseriate, oblong, obtuse, hyaline, 5 to 6 x 2  $\mu$ , with a single nucleus in the centre (becoming uniseptate)?

This is closely allied to *H. subcarnea* E. & E., but differs in the color of the subiculum, the black apex of the perithecia, the narrower asci, and rather longer sporidia.

HYPOCREOPSIS Karst.

**H. riccoidea** (Bolt.) Karst.

On *Kalmia latifolia*, alt. 2,000 ft., Aug. 24, 1893. Spores apparently moniliform (*Nuttall*, 1178).

GIBBERELLA Sacc.

**G. Saubinetii** (Mont.) Sacc.

On *Asparagus officinalis*, Nov. 1, 1893 (*Nuttall*, 1243, 190).

BROOMELLA Sacc.

**B. Ravenelii** (Berk.) Sacc.

On *Acer rubrum*, April 4, 1894 (*Nuttall*, 1461, 439).

PLEONECTRIA Sacc.

**P. Missouriensis** (E. & E.)

On dead stick, March 10, 1893 (*Nuttall*, 871).

CORDYCEPS Fries.

**C. militaris** (Linn.) Link.

In laurel thicket on ground, Aug. 12, 1893, Keeney's Creek (*Nuttall*, 1205, 161).

**C. ophioglossoides** (Ehrh.) Link.

On ground in wet woods, Aug. 24, 1893 (*Nuttall*, 1172).

## DOTHIDEACEÆ.

## PHYLLACHORA Nits.

**P. GRAMINIS** (Pers) Fckl.

On living leaves of *Hysterix Hysterix*, Fayette Co., near Nuttallburg (*Millspaugh*). On *Elymus Canadensis*, Oct. 28, 1893 (*Nuttall*, 1250, 192).

**P. Caricis** (Fr.) Sacc ?

On *Korycarpus diandrus*, March 1, 1894 (immature) (*Nuttall*, 1402, 238).

## DOTHIDELLA Spég.

**D. Ulmæ** (Sz.) E. & E.

On *Ulmus pubescens*, Nov. 5, 1894 (*Nuttall*, 1749, 629).

## PLOWRIGHTIA Sacc.

**P. MORBOSA** (Schw.) Sacc.

On limbs of cultivated Plum and Cherry, Monongalia Co., near Morgantown (*Millspaugh*).

## MICROTHYRIACEÆ.

## MYIOCOPORON Spég.

**M. Smilacis** (De Not) Sacc.

On *Smilax* sp. April 12, 1895 (*Nuttall*, 1807).

## LOPHIOSTOMACEÆ.

## LOPHIOTREMA Sacc.

**L. Nucula** (Fr.) Sacc.

On dead heartwood of *Liquidambar Styraciflua*, April 20, 1894 (*Nuttall*, 1479, 459).

## LOPHIDIUM Sacc.

**L. nitidum** E. & E., Proc. Phila. Acad., 333 (1894).

TYPE HABITAT: On *Vitis* cult. Feb. 2, 1894 (*Nuttall*, discov. 1336, 295).

Perithecia scattered, semi-erumpent, small (200 to 250  $\mu$  diameter) black and shining, subhemispherical, the flattened base immersed in the bark. Ostiolum compressed, thin, narrow, sometimes obsolete. Asci cylindrical, short (8 to 10  $\mu$ ), stipitate, about 100 x 10 to 12  $\mu$ , paraphysate, 8-spored. Sporidia uniseriate, oblong-elliptical, yellow-brown, 5 to 7-septate with a longitudinal septum running through two or more cells, 19 to 22 x 8 to 10  $\mu$ , ends mostly rounded and obtuse, not at all or only slightly constricted in the middle.



**L. compressum** (Pers.) Sacc.

On *Salix nigra falcata*, April 23, 1894. Asci 70 to 100 x 15  $\mu$ ,  
Spores 25 to 30 x 8 to 10  $\mu$  (*Nuttall*, 1481, 461).

HELVELLACEÆ.

MORCHELLA Dill.

**M. esculenta** (Linn.) Pers.

Along an old road among grasses and leaves under an oak,  
April 18, 1893 (*Nuttall*, 897).

**M. angusticeps** Peck.

On ground, Short Creek, April 18, 1893. Spores 25 to 28 x 15  $\mu$   
(*Nuttall*, 895).

**M.** \_\_\_\_\_ sp.

On ground, Short Creek, 5 cm. high, with smooth spores 28 x  
17  $\mu$ , April 18, 1893 (*Nuttall*, 896).

GYROMITRA Fr.

**G. curtipes** Fr.

On ground below cliffs, Short Creek. Asci, fertile part 50 x 16  
to 18  $\mu$ . Total length 350  $\mu$ . Spores 22 to 30 x 10  $\mu$ , 2-nucleate  
(*Nuttall*, 1813).

HELVELLA Linn.

**H. macropus** (Pers.) Karst?

On ground in woods at base of cliffs, July 15, 1893 (*Nuttall*,  
1097).

MITRULA Fr.

**M. phalloides** (Bull.) Chev.

In water on leaves of *Quercus Prinus*, etc., spores to 20 x 3  $\mu$ .  
Masterson's Glade, June 11, 1893 (*Nuttall*, 935).

**M. lutescens** B & C.

On ground and on dead logs in deep woods, alt. 2,000 ft.  
(*Nuttall*, 1024).

SPATHULARIA Pers.

**S. clavata** (Schæf.) Sacc?

In woods among dead leaves, stipe dark brown, spores *short*  
25  $\mu$ , July 10, 1893 (*Nuttall*, 1087).

VIBRISSEA Fr.

**V. truncorum** (A. & S.) Fr.

On wet dead sticks, alt. 1,800 ft., June 15, 1893 (*Nuttall*, 991).

## PEZIZACEÆ.

## RHIZINA Fr.

**R. inflata** (Schæf.) Quéf.

On sandstone rock in sandy loam. Blackish-brown, spreading, 2 to 9 cm. x 1 to 3 cm., convex, loosely attached to the soil by strong rootlets from all parts of the underside, margin strongly rolled inward. Asci 375? to 415 x 12.5  $\mu$ , spores hyaline, apiculate, large 40 x 10  $\mu$ . Keeney's Creek, Nov. 3, 1893 (*Nuttall*, 1224).

## GEOPYXIS Pers.

**G. nebulosa** Cooke.

On dead wet wood at low and high altitudes, Aug. 10, 1893 (*Nuttall*, 1177).

**G. carbonaria** A. & S.

On ground upon which logs had been burned the previous autumn, April 18, 1893 (*Nuttall*, 908).

## PEZIZA Dill.

**P. aurantia** Pers.

On wet and freshly crushed sandstone ballast, and on filling of sandstone cut on railroad. Spores curiously rough and apiculate at both ends. Nov. 3, 1893 (*Nuttall*, 1225).

**P. badia** Pers.

On ground, Short Creek, April 18, 1893 (*Nuttall*, 922).

## OTIDEA Pers.

**O. onotica ochracea** Fr.

On ground and on rotten *Tsuga Canadensis*, July 14, 1893. Asci 175 x 10  $\mu$  total length, spores 11 to 13 x 7.5  $\mu$ , nuclei 2 (*Nuttall*, 1098).

## HUMARIA Fr.

**H. echinosperma** Peck.

In soil of ditch, July 20, 1894 (*Nuttall*, 1630).

## SARCOSYPHA Fr.

**S. coccinea** Jacq.

On sticks, on ground in damp shady place, March 20, 1893 (*Nuttall*, 852).

**S. floccosa** Schw.

On a stick in a laurel thicket, alt. 2,000 ft., July 12, 1893 (*Nuttall*, 1112).

## LACHNEA Fr.

**L. SCUTELLATA**, Linn.

On rotting log *Fagus atropunicea* and under bark of wet log *Quercus alba*, Monongalia Co., near Morgantown (*Millsbaugh*). On rotten wood in old drift of mine, and on bare sandy ground among pebbles, Short Creek, May 11-17, 1893 (*Nuttall*, 936).

**L. umbrarum** Fr.

On open bare sandy soil, Short Creek, May, 1893 (*Nuttall*, 1107).

**L. hirta** Schum.

On floor of cave, July 25, 1893 (*Nuttall*, 1127).

**L. vitellina** Pers:

On dead log. Asci  $235 \times 15 \mu$ , Short Creek, July 2, 1893 (*Nuttall*, 1114).

**L. ERINACEUS**, Schw.

On underside of log *Quercus alba*, Monongalia Co., near Morgantown (*Millsbaugh*).

SCLEROTINIA Fuckel.

**S. trifoliorum** Eriks.

On ground in open woods, May 18, 1893 (*Nuttall*, 960).

HELOTIUM Fr.

**H. lacteum** E. & E., Proc. Phila. Acad., 145 (1893).

On bare log, Dec. 16, 1893. Pure white, stipitate, plane .4 to .6 mm. asci clavate  $95$  to  $110 \times 7.5 \mu$ . Spores curved  $16$  to  $25 \times 3$  to  $4 \mu$ , nuclei 4, paraphyses slender, Short Creek near the falls (*Nuttall*, 1318).

**H. herbarum** (Pers.) Fr.

On *Polymnia Uvedalia*, Nov. 15, 1894. Cups 6 to 12 mm. diameter, stipes 1 to 2 mm. long. Asci  $55$  to  $75 \times 7.5$  to  $10 \mu$ . Spores  $12$  to  $15 \times 2.5 \mu$  (*Nuttall*, 1755).

**H. confluens** Schw.

On dead logs, alt. 1,800 ft. Oct. 20, 1893 (*Nuttall*, 1213).

**H. citrinum** (Hedw.) Fr.

On wet dead log. Short Creek, alt. 1,800 ft. Sept. 12, 1893. (*Nuttall*, 1201).

**H. castaneum** Sacc.

On leaves of *Rhododendron maximum*. Short Creek, alt. 1,800 ft. Sept. 12, 1893 (*Nuttall*, 1206).

PHIALEA Fr.

**P. virgultorum** (Vahl.) Sacc.

On old grape twigs on wet ground. Asci  $105 \times 7.5 \mu$ . Spores  $18 \times 4 \mu$ . Nov. 1, 1894 (*Nuttall*; 1735).

PSEUDOHELOTIUM Fckl.

**P. fibrisedum** (B. & C.) Sacc.

On wet dead wood of *Juglans cinerea*, Dec. 5, 1894. More than 2 mm diameter, margin ciliate, red-maculate; asci oblong short-stipitate, total length  $65 \times 7.5 \mu$  (approx.), paraphyses filiform. Spores hyaline 1-septate (?) oblong, straight or slightly curved,  $12$  to  $18 \times 3 \mu$  (*Nuttall*, 1764).

## CYATHICULA De Not.

**C. quisquiliaris** E. & E., Proc. Phila. Acad., 451 (1893).

TYPE HABITAT: On dead racemes of *Quercus Prinus*, on dead leaves, on rock in woods, April 2, 1893 (Nuttall, discov. 941).

Gregarious, sessile, 1 to 1.5 mm. diameter, cup-shaped, disk pale with a tinge of brick color when dry, dull, dirty-white and purpuraceo-squammulose outside, margin distinctly cleft-toothed and incurved when dry. Asci cylindrical, subsessile, 55 to 65 x 6 to 7  $\mu$ , with filiform paraphyses scarcely thickened above. Sporidia uniseriate, or sub-biseriate above, elliptical, hyaline, continuous, 6 to 8 x 3 to 4  $\mu$ .

## CHLOROSPLENIUM Fr.

**C. æruginosum** (Eder) De N.

On loose core inside dead tree, Nov. 6, 1893. Disk more brilliant than margin in light and when young, in shadow varying to pure white. Asci over 105 x 7.5  $\mu$  total length, spores guttate 12.5 x 3  $\mu$  (Nuttall, 1235).

**C. versiforme** (Pers.) Karst.

On rotten mossy log, Short Creek, Oct. 28, 1893 (Nuttall, 1238).

**C. Schweinitzii** Fr.

Top of oak log, Dec. 10, 1893. Yellow in spring, green in autumn and winter (Nuttall, 1276).

## MOLLISIA Fr.

**M. Dehnii** (Rabenh) Karst.

On *Potentilla Monspeliensis*, July 21, 1894 (Nuttall, 1618).

**M. pinastri** (C. & P.) Sacc.

On *Pinus Virginiana*, Dec. 3, 1894 (Nuttall, 1763).

**M. CINEREA** (Batsch.) Karst.

*Peziza cinerea* Batsch. On decayed log, Monongalia Co., near Morgantown (Millspaugh).

## PYRENOPEZIZA Fuckel.

**P. lacerata** (C. & P.) Sacc.

On *Rubus odoratus*, May 31, 1894. Asci 45 to 50 x 5 to 6  $\mu$ . Spores 8 to 12.5 x 2 to 2.5  $\mu$  (Nuttall, 1529).

## TAPESIA Pers.

**T. sanguinea** (Pers.) Fckl.

On dead log, and on *Magnolia Fraseri* and *Juglans cinerea*, June 10, 1894 (Nuttall, 1546).

**T. fusca** (Pers.) Fckl.

On rotten burned log *Tsuga Canadensis*; on dead rotting log *Juglans cinerea*, alt. 1,300 ft. Asci 45 to 92 x 495  $\mu$ . Spores 8 to 10 x 2  $\mu$ . Short Creek, Feb. 2, 1894 (Nuttall, 1365).

**T. subiculata** (Schw.) Sacc.

On wet sticks, June 15 and Oct. 25, 1893. Spores large, 10 x 3  $\mu$  (*Nuttall*, 992).

LACHNELLA Fr.

**L. Virginia** E. & E., Proc. Phila. Acad. 349 (1894).

TYPE HABITAT: On decaying wood of *Magnolia Fraseri*, Feb., 1894 (*Nuttall*, discov. 1410, 380).

Gregarious, sessile, .75 to 1.2 mm. diameter. Disk pallid, concave, nearly closed when dry, margin fringed with brown septate hairs, 100 to 125 x 3  $\mu$ , outside clothed with shorter reddish-brown hairs. Asci about 50 x 5  $\mu$ , clavate-cylindrical, short stipitate, paraphysate, 8-spored. Sporidia sub-biseriate, oblong, 6 to 8 x 1.5 to 2.5  $\mu$ .

TRICHOPEZIZA Fockel.

**T.** ——— sp.

Torn end of firm dead tree, Short Creek, alt. 1,800 ft. Dec. 16, 1893 (*Nuttall*, 1326).

**T. albo-lutea** (Pers.) Sacc.?

On rotten log, Short Creek, June 18, 1893, stipitate (*Nuttall*, 1013).

**T. capitata** (Peck) Sacc.

On leaves *Quercus palustris* on ground in woods, alt. 2,000 ft., June 18, 1893. At Rupert's, July 26, 1894 (*Nuttall*, 1011).

DASYSCYPHA Fr.

**D. virginea** (Batsch) Fockl.

On dead sticks on wet cliff, and on wet dead wood of *Juglans cinerea*, June 12, 1893 (*Nuttall*, 841).

**D. Ellisiana** (Rehm.) Sacc.

On *Pinus rigida*, top of Cavendish cliff, April 8, 1894 (*Nuttall*, 1447).

HELOTIELLA Sacc.

**H. Nuttallii** E. & E., Proc. Phila. Acad., 351 (1894).

TYPE HABITAT: On wet shady side of rotten log, March 15, 1894 (*Nuttall*, discov. 1425, 397).

Ascomata gregarious, sessile, flat-hemispherical, .5 to 1 mm. diameter, cup-shaped when fresh, contracted and sub-spherical when dry with only a small round apical opening, orange color throughout, disk watery-orange, outside and fringed margin paler. Substance fibrous, the fibers prolonged above so as to make a marginal fringe of pale yellow, roughish straight hairs. Asci clavate-cylindrical, sessile, curved, 50 to 60 x 5 to 6  $\mu$ . Paraphyses filiform, hardly thickened above, with a series of orange-colored, globose nuclei. Sporidia biseriate, oblong, hya-

line, obtuse, uniseptate, but not constricted, 8 to 10 x 2.5 to 3  $\mu$ . The outside of the ascomata is slightly hairy so that the species might be referred to *Solenopeziza* if that is really worthy of generic distinction.

#### DERMATELLA Karst.

##### D. viticola E. & E., Proc. Phil. Acad., 352 (1894).

TYPE HABITAT: On dead shoots of *Vitis* (cult.) Nov. 1893 (*Nuttall*, discov. 1337, 295).

Sessile, convex-discoid and pale when moist, concave and flesh-color when dry, and then scarcely projecting above the surface of the bark. .50 to .75 mm. diameter. Asci clavate-cylindrical, about 70 x 12  $\mu$ , with filiform paraphyses, 8-spored, sub-sessile. Sporidia subbiserial, oblong elliptical, hyaline, 1 to 3-septate (3 to 4-nucleate at first), 15 to 18 x 6 to 6.5  $\mu$ .

#### BELONIDIUM Mont.

##### B. AURELIA (Pers.) DeNot.

*Peziza aurelia*, Pers. On a dead leaf in rotten log, Monongalia Co., at Little Falls (*Millsbaugh*).

#### ERINELLA Sacc.

##### E. miniopsis (Ell.) Sacc.

On wet rotten wood, May 1893 (*Nuttall*, 934).

#### ASCOBOLACEÆ

##### ASCOBOLUS, Pers.

##### A. furfuraceus Pers.

On old dung of cow, June 8, 1893 (*Nuttall*, 974).

##### A. carbonarius Karst.

On bare spots where logs were burned by forest fire the previous winter, associated with *Geopyxis carbonaria*. It appears that only the prolonged burning of large logs makes proper habitat for this species, as it does not grow where the fire was light. May 20, 1893 (*Nuttall*, 926).

##### ASCOPHANUS Boud.

##### A. granuliformis (Cr.) Boud. ?

On dung of cow, Dec. 1893 (*Nuttall*, 1340, 298).

##### A. carneus (Pers.) Boud.

On dung of cow Oct. 10, 1893 (*Nuttall*, 1249, 182).

#### DERMATEACEÆ

##### URNULA Fr.

##### U. Craterium (Schw.) Fr.

On dead twigs on ground, Short Creek, April 15, 1893 (*Nuttall*, 898).

DERMATEA Fr.

**D. Chionanthi** E. & E?

Macrostylospores of this species? on *Chionanthus Virginica*, March, 25, 1894 (*Nuttall*, 1454, 426).

**D. lobata** Ellis.

On the maculæ of *Pestalozzia Guepinii* on leaves of *Rhododendron maximum*. Asci 130 x 12.5  $\mu$  spores 15 x 8  $\mu$ . Aug. 15, 1895 (*Nuttall* 1162, 132).

TYMPANIS Tode.

**T. Oxydendri** E. & E., Proc. Phila. Acad., 352 (1894).

TYPE HABITAT: On dead limbs of *Oxydendron arboreum*, Dec. 12, 1893 (*Nuttall* discov., 1296, 258).

Cespitose, 3 to 5 together, erumpent, at length deciduous, black .5 to .75 mm. diameter, contracted below into a short thick stipe, margin subundulate. Asci cylindrical 110 to 130 x 15 to 18  $\mu$  including the short stipe. Paraphyses slender, abundant, Sporidia numerous, minute, oblong, hyaline or yellowish 2.5 to 3 x 1  $\mu$ , Allied to *T. fasciculata*, but asci larger and sporidia smaller.

CENANGELLA Sacc.

**C. Aceris** (Hazzl.) Sacc.

On bark of dying *Acer rubrum*, June 10, 1894. Sporidia 15 x 6  $\mu$  (*Nuttall*, 1554).

SCLERODERRIS Fr.

**S. rhabarbarina** (Berk) E. & E.

On *Rubus Baileyanus*, Dec. 8, 1893, Short Creek. Asci total length 63 to 75 x 17 to 21  $\mu$ , 8-spored, clavate, very short stipitate. Spores 20 to 27 x 6 to 7.5  $\mu$ , nucleate, becoming clearly 3-septate and rather one-sided (*Nuttall*, 1294, 242).

**S. pallidula** (Cooke) Sacc.

On dead branches and bark of *Acer saccharinum*, L., Feb. 26, 1894. On *Oxydendron arboreum*. Sporidia 20 to 22 x 7  $\mu$  (*Nuttall*, 1408, 665).

BULGARIACEÆ.

LEOTIA Hill.

**L. lubrica** (Scop.) Pers.

On ground, alt. 1,800 ft., Short Creek, Aug. 14, 1893 (*Nuttall*, 1150).

ORBILIA Fr.

**O. rubella** (Pers.)

On decorticated log, *Liriodendron Tulipifera*, Sept. 19, 1895 (*Nuttall*, 1858, 748).

**O. vinosa** (A. & S.) Karst.

On *Poria spissa*, Nov. 6, 1893 (*Nuttall*, 1261).

**O. occulta** (Rehm) Sacc.

On dead log, Aug. 20, 1893 (*Nuttall*, 1202).

**O. coccinella** (Somm.) Karst.

On dead log *Juglans cinerea*, Short Creek, July 25, 1893 (*Nuttall*, 1134, 663).

**O. Xanthostigma** Fr.

On decorticated limbs *Magnolia Fraseri*, Sept. 1895 (*Nuttall*, 1855, 742, 743).

## BULGARIA Fr.

**B. inquinans** (Pers.) Fr.

In clefts of bark of trees and railroad ties, Oct. 10, 1893 (*Nuttall*, 1210).

## CORYNE Tul.

**C. URNALIS** (Nyl.) Sacc.

On rotting log *Fagus atropunicea*; under bark of *Quercus alba* log, Monongalia Co., near Morgantown (*Millsbaugh*). On wet mossy logs, alt. 1,800 ft. and over, Oct. 10, 1893 (*Nuttall*, 1212, 167).

## HOLWAYA Sacc.

**H. ophiobolus** (Ellis) Sacc.

On bark of dead *Magnolia Fraseri*, alt. 1,800 ft., Dec. 16, 1893, Short Creek (*Nuttall*, 1323, 279).

## STICTACEÆ.

## PROPOLIS Fr.

**P. faginea** (Schrad.) Karst.

On *Juglans cinerea* and on old board of *Liriodendron Tulipifera*, Dec. 5, 1894 (*Nuttall*, 1765, 660). On *Liquidambar styraciflua*.

**P. Leonis** (Tul.) Rehm.

On bark of *Pinus rigida* and *Virginiana*, June 31, 1894 (*Nuttall*, 1581).

## STICTIS Pers.

**S. fimbriata** Schw.

On strobiles of *Pinus Virginiana*, alt. 2,000 ft., March, 1894 (*Nuttall*, 1468, 443). Sporidia 5 to 7 x 1.5  $\mu$ .

## PHACIDIACEÆ.

## RHYTISMA (Pers.) Fr.

**R. punctatum** (Pers.) Fr.

On leaves of *Acer Pennsylvanicum*, alt., 2,000 ft., Oct. 22, 1893 (*Nuttall*, 1217, 168).

**R. decolorans** Fr.

On *Xolisma ligustrina*, at Rupert's, July 26, 1894 (*Nuttall*, 1638, 569).



**R. Prini** Schwein.

On leaves of *Ilex verticillata*, at Rupert's, Sept. 7, 1894 (*Nuttall*, 1677).

PATELLARIACEÆ.

DURELLA Tul.

**D. commutata** Fckl.

On dead log, Short Creek, Dec. 16, 1893 (*Nuttall*, 1328).

LECANIDION Rabenh.

**L. atratum** (Hedw.) Rabenh.

On *Aralia spinosa*, Feb. 17, 1894. Sporidia 50 x 12.5  $\mu$  (*Nuttall*, 1390, 359).

**L. Hamamelidis** (Peck) Sacc.

On bark of *Hamamelis Virginica*. Sporidia 1 to 3 septate when mature. Dec. 18, 1894 (*Nuttall*, 1772).

CALICIACEÆ.

CALICIUM Pers.

**C. tigillare** (B. & Br.) Sacc.

On *Polyporus pergamenus*, Aug. 23, 1893 (*Nuttall*, 1159).

GYMNOASCACEÆ.

EXOASCUS Fckl.

**E. DEFORMANS** (Berk.) Fckl.

*Taphrina deformans* Tul. On leaves of *Amygdalus Persica*, Jefferson Co., near Charlestown (*Millsbaugh*).

**E. PRUNI** Fckl.

*Taphrina Pruni* Tul. On fruits of *Prunus domestica*, Monongalia Co., at Morgantown (*Millsbaugh*).

HYSTERIACEÆ.

GLONIUM Mühl.

**G. stellatum** Mühl.

On rotten oak plank, and on *Sassafras Sassafras*, July 28, 1893 (*Nuttall*, 1128).

**G. parvulum** (Ger.) Sacc.

On fence rails *Castanea dentata*, April 20, 1894 (*Nuttall*, 1496).

**G. simulans** Ger.

On old fence rails, alt. 2,000 ft., March 16, 1894. Asci 75 x 7.5 to 10  $\mu$ . Spores 13 to 18 x 5  $\mu$ . On *Juglans cinerea*, asci 65 x 10  $\mu$ . Spores 12 to 15 x 4 to 5  $\mu$  (*Nuttall*, 1415).

HYSTERIUM Tod.

**H. Kalmiæ** Schw.?

On dead *Kalmia latifolia*, Nov., 1893 (*Nuttall*, 1287, 232).

## HYSTEROGRAPHIUM Corda.

**H. insidens** (Schw.) Sacc.

On old fence rails, April 6, 1894. Largest spores  $43 \times 7.5 \mu$ , 11-septate (Nuttall, 1452).

**H. subrugosum** (C. & E.) Sacc.

On old fence rails, May 26, 1894. Spores 33 to  $35 \times 12.5 \mu$  (Nuttall, 1522).

**H. Mori** (Schw.) Rehm.

On old fence rails, April 6, 1894 (Nuttall, 1451).

## HYPODERMA DC.

**H. DESMAZIERI** DuBy.

On living leaves or *Pinus echinata*. Wood Co., near Lockhart's Run (Millspaugh).

**H. virgultorum** DC.

On *Rubus Baileyanus*, Dec. 5, 1893 (Nuttall, 1293).

**H. commune** (Fr.) Duby.

On *Hydrangea arborescens*, Nov. 3, 1893 (Nuttall, 1259). On old petioles *Aralia spinosa* (Nuttall, 1830, 719).

## LOPHODERMIUM Chev.

**L. Rhododendri** (Schw.)

On *Rhododendron maximum*, Sept. 4, 1893 (Nuttall, 1192).

**L. Pinastri** (Schrad.) Chev.

On leaves of *Pinus rigida*, Short Creek, Aug. 2, 1893 (Nuttall, 1158).

**L. culmigenum** (Fr.) Karst.

On straw brought in from Ohio, May 24, 1894. Measurements of asci and spores prove to be the same as those of *L. arundinaceum* (Schrad.) Chev. (Nuttall, 1496).

## OSTROPA Fr.

**O. cinerea** (Pers.) Fr.?

Issuing from cracks in a decorticated log, May 31, 1894 (Nuttall, 1526).

## AGARICACEÆ.

## AMANITA Pers.

**A. verna** Fr.

On ground in open woods, July 11, 1893. Spores round  $7.5 \mu$  (Nuttall, 964).

**A. MUSCARIA** Linn.

Rooted on buried limb of *Betula lenta*, Grant Co., near Bayard (Millspaugh).

An ecarunculate form, apparently of this species, on leaf mold in deep woods, same locality.

**A. muscaria** alba Peck.

On ground in laurel thicket, Short Creek, alt. 2,000 ft. Has the odor of buttermilk. Spores  $9 \times 6 \mu$  (Nuttall, 1131).

**A. flavo-rubens** Berk.

On ground in woods, Short Creek, July 6, 1893 (Nuttall, 1078).

**A. polypyraxis** B. & C.

Among bushes, alt. 2,400 ft., July 6, 1893. Largest stipes 15 cm. long, thick, heavy and deep in ground. Pileus 15 cm. broad. Spores  $10 \times 6 \mu$  (Nuttall, 1101).

AMANITOPSIS Roze.

**A. nivalis** (Grev.) Sacc.

In loam on rocks in laurel tangle, June 26, 1893 (Nuttall, 1004).

TRICHOLOMA Fr.

**T. fumoso-luteum** Peck?

On unknown substance, June 15, 1893, Short Creek (Nuttall 990).

CLITOCYBE Fr.

**C. illudens** Schwein.

On old stumps in clusters of about eight, dark and deep rich orange color throughout,  $+20 \text{ cm.} \times +12 \text{ cm.}$  broad, stipe  $+25 \text{ mm.}$  thick. Spores  $+6 \mu$  diameter, uneven. July 14, 1893 (Nuttall, 1093).

**C. laccata** Scop.

Common in wet sand, bank of Short Creek, alt. 2,000 ft., July 28, 1893. Spores  $10 \mu$  diameter, echinate (Nuttall, 1124).

COLLYBIA Fr.

**C. RADICATA** Relh.

On leaf mold in rich woods, Monongalia Co., near Morgantown (Millsbaugh). May 17, 1893 (Nuttall, 980).

**C. platyphylla** Fr.

On mold and moss on trunk of tree, Short Creek, June 15, 1893 (Nuttall, 981).

**C. velutipes** Curt.

About roots of rotting stumps, Jan. 31, 1895 (Nuttall, 830).

**C. conigenoides** Ellis.

On last year's cones *Magnolia Fraseri*, Sept. 18, 1895 (Nuttall, 1848).

**C. DRYOPHILA** Bull.

In moss on log in deep woods, Grant Co., near Bayard (Millsbaugh).

MYCENA Fr.

**M.** \_\_\_\_\_ sp.

White. On mossy stump, July 4, 1893 (Nuttall, 1019).

**M. GALERICULATA** Scop.

In rich woods under oak twig, and oak bark on log, Monongalia Co., near Morgantown (*Millspaugh*). On damp dead logs, May 7, 1893 (*Nuttall*, 930).

**M. epipterygia** Scop.

On moss-covered log and in clefts of bark, Feb. 28, 1893 (*Nuttall*, 948).

## OMPHALIA Fr.

**O. CAMPANELLA** Batsch.

On leaf mold at base of *Castanea dentata*, Preston Co., near Terra Alta (*Millspaugh*). On rotten log in thicket, June 18, 1893 (*Nuttall*, 1009).

## PLEUROTUS Fr.

**P. corticatus** Fr.?

On fallen *Acer* sp. Gills not anastomosing, July 28, 1893 (*Nuttall*, 1123).

**P. sapidus** Kalchbr.

On dead log, Feb. 2, 1893 (*Nuttall*, 850).

**P. serotinus** Schrad.

On dead trunks, Nov. 20, 1893 (*Nuttall*, 1267).

**P. cyphelliformis** Berk.

On dead stems of *Polymnia Uvedata*, Nov. 15, 1894 (*Nuttall*, 1742).

**P. niger** Fr.

On bark of dead tree, May 13, 1893 (*Nuttall*, 953).

## HYGROPHORUS Fr.

**H. nitidus** B. & C.

On decayed log, May 14, 1893 (*Nuttall*, 968).

## LACTARIUS Fr.

**L. cilicioides** Fr.

On border of woods and swamp, July 4, 1893 (*Nuttall*, 1020).

## RUSSULA Pers.

**R. emetica** Fr.?

In woods, July 4, 1893 (*Nuttall*, 1021).

## CANTHARELLUS Adans.

**C. minor** Peck?

On ground in woods, alt. 2,000 ft., July 4, 1893 (*Nuttall*, 1023).

**C. Wrightii** Berk?

Wet sand of river bank in shade, July 18, 1893. Whole plant light red except gills, which are little deeper red than salmon-color, gills decurrent, veined between, pileus more convex than

plane, and depressed edge wavy, rather tough. Largest 4 cm. broad, stem 5 cm. high, 5 mm. thick. Spores 7.5 to 8.5 x 3.5 to 5  $\mu$ , apparently rough and guttulate inward (*Nuttall*, 1104).

**C. tubæformis** Fr.?

On rotten wood in woods, alt. 2,000 ft., July 4, 1893 (*Nuttall*, 1022).

**C. floccosus** Schw.

On open bank of stream in laurel thicket, alt. 2,000 ft., June 20, 1893. Largest squamules .75 cm. long, thick and triangular, forming a cone which closes the funnel below the top an inch or more. Spores 15 x 7.5  $\mu$  (*Nuttall*, 992).

NYCTALIS Fr.

**N. asterophora** Fr.

On dead Agaric in deep damp woods, July 14, 1893. Stars 20  $\mu$  diameter (*Nuttall*, 1096).

MARASMIUS Fr.

**M. semihirtipes** Peck.

On dead leaves, sticks, etc., Short Creek, May 14, 1893 (*Nuttall*, 967).

**M. ramealis** (Bull.) Fr.

Very common on twigs and leaves in laurel thickets, June 15, 1893 (*Nuttall*, 994).

**M. OPACUS** B. & C.

Common on dead branches *Rhododendron maximum*, Grant Co., near Bayard (*Millsbaugh*).

**M. campanulatus** Peck.

On dead leaves and sticks in damp woods, Short Creek, alt. 1,300 ft., Aug. 16, 1893 (*Nuttall*, 1153).

**M. ROTULA** (Scop.) Fr.

On dead limb of *Betula lenta*, Grant Co., near Bayard (*Millsbaugh*). On stick in open woods, June 8, 1893 (*Nuttall*, 972).

LENTINUS Fr.

**L. STRIGOSUS** Fr.

On dead log *Betula lutea*, Grant Co., near Bayard (*Millsbaugh*). On dead logs, common, May 6, 1893 (*Nuttall*, 904).

PANUS Fr.

**P. stipticus** (Bull.) Fr.

On fallen dead limb, Jan. 29, 1893 (*Nuttall*, 950).

**P. dorsalis** Bosc.

On *Tsuga Canadensis*, Sept., 1893 (*Nuttall* 1253, 196).

TROGIA Fr.

**T. crispa** (Pers.) Fr.

On various dead dry logs and limbs, March 21, 1893 (*Nuttall*, 875).

## LENZITES.

**L. betulina** (Linn.) Fr.

On logs, Sept. 6, 1893 (*Nuttall*, 837).

**L. SEPIARIA** Fr.

On decorticated stumps *Picea Mariana*, Tucker Co., near Falls of Blackwater (*Millspaugh*).

A resupinate form on dead logs *Tsuga Canadensis* Grant Co.

**L. abietina** (Bull.) Fr.

On dead branches *Tsuga Canadensis*, alt. 1,600 ft., Dec. 15, 1893 (*Nuttall*, 1301).

**L. corrugata** Klot.

On *Castanea dentata*, alt. 1,800 ft., Dec. 19, 1893 (*Nuttall*, 1302).

## SCHIZOPHYLLUM Fr.

**S. COMMUNE** Fr.

On bark log *Quercus alba*, Grant Co., near Bayard. On dead twig *Pyrus Malus*, Monongalia Co., near Morgantown (*Millspaugh*)  
On dead logs, Jan. 30, 1893 (*Nuttall*, 827).

## VOLVARIA Fr.

**V. BOMBYCINA** (Pers.) Fr.

On dead insect sp? Monongalia Co., near Morgantown (*Millspaugh*).

## CLITOPILUS Fr.

**C. abortivus** Fr.

Wet woods, alt. 1,800 ft, aborted specimens in quantity, Sept. 4, 1893 (*Nuttall*, 1190).

## AGARICUS Linn.

**A. Rodmani** Peck.

On ground, June 28, 1893 (*Nuttall*, 1005).

**A. campester** Linn.

On lawn, June 24, 1893 (*Nuttall*, 1002).

**A. silvicola** Vitt.

In woods, Short Creek, alt. 1,150 ft. (*Nuttall*, 1699). Spores 5 to 7 x 3 to 4  $\mu$ .

**A. silvaticus** Schæff.

In woods, Short Creek, July 2, 1893 (*Nuttall*, 1117).

## STROPHARIA Fr.

**S. STERCORARIA** Fr.

On decaying vegetable matter, Preston Co., near Terra Alta (*Millspaugh*).

## HYPHOLOMA Fr.

**H. SUBLATERITIUM** Schæff.

Under bark log *Fraxinus Americana*, Monongalia Co., near Morgantown (*Millspaugh*).

**H. elæodes** Fr?

On sticks on ground, May 18, 1893 (*Nuttall*, 858).

COPRINUS Pers.

**C. variegatus** Peck?

On stick in shady place, Aug. 11, 1893 (*Nuttall*, 1143).

**C. micaceus** (Bull) Fr.

On bases of stumps in woods, July 8, 1893 (*Nuttall*, 1081).

**C. domesticus** (Pers.) Fr.

In woods on mixture of manure and leaf mold. June 29, 1893 (*Nuttall*, 1007).

**C. semilanatus** Peck?

In heavily manured asparagus bed in garden, Apr. 29, 1893 (*Nuttall*, 903).

**C. ephemerus** Fr.

On horse dung, July 2, 1893 (*Nuttall*, 1113).

**C. angulatus** Peck.

On baked soil where log had been burned, with *Peziza carbonaria*, May 8, 1893 (*Nuttall*, 932).

PANÆOLUS Fr.

**P. CAMPANULATUS** Linn.

On cow dung in deep coniferous woods, Grant Co., near Bayard (*Millsbaugh*). Plentiful on manured lawn, May 8, 1893, (*Nuttall*, 933).

PSATHYRELLA Karst.

**P. disseminata** Pers.?

In rotten mold about the base of a stump, a mass about 1 meter containing perhaps thousands of specimens. Specimens 3 to 5 cm. high, thin, fleshless. Pileus 1 cm. broad, 8 mm. high, slightly umbonate, sulcate striate, mealy, gray. Stems white, smooth 1 mm thick, at first slightly furfuraceous. Largest stems 6 cm., broadest pileus 18 mm. Spores 6 to 10 x 3 to 5  $\mu$  (*Nuttall*, 1189). Sept. 3. 1893.

POLYPORACEÆ.

BOLETUS Dill.

**B. speciosus** Frost.

On ground in woods, alt. 2,000 ft. July 16, 1893 (*Nuttall*, 1099).

**B. chrysenteron** Fr.

In deep woods, June 18, 1893 (*Nuttall*, 987).

**B. parasiticus** Bull.

On *Scleroderma vulgare* wet woods, alt. 1,800 ft., Sept. 4, 1893. (*Nuttall*, 1191).

**B. edulis** Bull.

Old field under *Rhus copallina*, June 20, 1893. Pileus 15 x 10 cm. stipe 4.5 cm. Spores 10 to 14 x 3 to 4  $\mu$  (*Nuttall*, 990).

## STROBILOMYCES Berk.

**S. strobilaceus** (Scop.) Berk.

Shady fence row, alt. 2,300 ft., Sept. 16, 1893 (*Nuttall*, 1100). Spores about 10  $\mu$ , round.

## FISTULINA Bull.

**F. hepatica** Fr.

On dead stumps, alt. 2,000 ft., July 16, 1893 (*Nuttall*, 1102).

## POLYPORUS Mich.

**P. brumalis** (Pers.) Fr.

On dead logs, Mar. 6, 1893 (*Nuttall*, 1835).

**P. pocula** (Schw.) B. & C.

On dead *Rhus copallina*, Glade Creek, May 4, 1894 (*Nuttall*, 1495, 478).

**P. arcularius** (Batsch) Fr.

On rotten logs, Short Creek, May 6, 1893. Largest 4 cm. broad, stipe 2.3 cm. (*Nuttall*, 929).

**P. varius** Fr.

On dead logs, Aug. 18, 1893. Thin, tough, 20 cm. broad (*Nuttall*, 840).

**P. UMBELLATUS** Fr.

In dry exposed hollow of oak stump, Monongalia Co., near Morgantown (*Millspaugh*).

**P. lactifluus** Peck.

On ground in open woods, June, 1893; Aug. 24, 1893. Height 30 cm., greatest breadth 60 cm. (*Nuttall*, 1171).

**P. BERKELEYI** Fr.

In dry exposed hollow of oak stump, Monongalia Co., near Morgantown (*Millspaugh*).

**P. SULPHUREUS** (Bull.) Fr.

On decaying oak stump, Monongalia Co., near Morgantown (*Millspaugh*).

**P. nidulans** Fr.

*P. niveus*, Fr. On dead trunk *Prunus serotina*, Short Creek, alt. 1,000 ft., July 8, 1894 (*Nuttall*, 1599, 548).

**P. ADUSTUS** (Willd.) Fr.

On dead branch *Rhus hirta*, and in decayed stump *Hicoria ovata*, Monongalia Co., near Morgantown (*Millspaugh*). On sides of dead logs, Feb. 2, 1893 (*Nuttall*, 851).



**P. hypococcineus** Berk?

A resupinate form of this species, or new (E. & E). Under-  
side of dead log, Oct. 15, 1893 (*Nuttall*, 1214, 166).

**P. pubescens** (Schum.) Fr.

On dead *Betula* sp. July 27, 1894 (*Nuttall*, 1673).

**P. resinusus** (Schrad.) Fr.

On dead log, March 24, 1893 (*Nuttall*, 879).

**P. Pilotæ** Schw.

On dead rotten log, Sewell Valley, Aug. 23, 1894 (*Nuttall*,  
1661).

FOMES Fr.

**F. LUCIDUS** (Leys.) Fr.

On dead logs *Tsuga Canadensis*, Preston Co., near Terra Alta  
(*Millsbaugh*). On old stumps in deep woods, April 10, 1893,  
full grown in June (*Nuttall*, 892).

**F. volvatus** Peck.

On west side *Pinus Virginiana*, March 8, 1893 (*Nuttall*, 870).

**F. pinicola** Fr.

On fallen trunks *Tsuga Canadensis*, Short Creek, May 1, 1893  
(*Nuttall*, 924).

**F. APPLANATUS** (Pers.) Wallr.

On dead *Acer saccharum* and *Quercus palustris*, *alba* and *rubra*  
Wood Co., near Kanawha Station, and McDowell Co., near Elk-  
horn. On dead *Fraxinus Americana*, Grant Co., near Bayard,  
and Monongalia Co., near Morgantown (*Millsbaugh*). On dead  
logs, Jan., 1893 (*Nuttall*, 828).

**F. FOMENTARIUS** (Linn.) Fr.

On dead *Betula lutea*, Grant Co., near Bayard (*Millsbaugh*).  
On dead log, at Rupert's, July 26, 1894 (*Nuttall*, 1635).

**F. RIMOSUS** Berk.

On dead *Robinia Pseudacacia*, Monongalia Co., near Morgan-  
town (*Millsbaugh*). On same sp. and on oak? alt. 2,000 ft.,  
March 21, 1893 (*Nuttall*, 914).

**F. CARNEUS** Nees.

On dead decorticated *Picea Mariana*, Tucker Co., near Falls of  
Blackwater (*Millsbaugh*). On *Pinus Virginiana*, March, 1893  
(*Nuttall*, 865).

POLYSTICTUS Fr.

**P. PERGAMENUS** Fr.

On dry exposed oak railroad tie, Monongalia Co., near Morgan-  
town (*Millsbaugh*). On various dead trees, Feb. 17, 1893 (*Nutt-*  
*all*, 847).

**P. CINNABARINUS** (Jacq.) Fr.

On dead limbs of cultivated *Prunus*, Monongalia Co., near  
Morgantown (*Millsbaugh*). On various dead deciduous trees,  
Jan. 31. 1893 (*Nuttall*, 829).

**P. VERSICOLOR** (Linn.) Fr.

On decorticated stump *Picea Mariana*, Tucker Co., near Falls of Blackwater. Under bark of log *Quercus alba*, Monongalia Co., near Morgantown (*Millspaugh*). On various dead trees, Feb., 1893 (*Nuttall*, 834).

**P. HIRSUTUS** Fr.

On dead twig cultivated *Pyrus Malus*, Monongalia Co., near Morgantown. On dead *Betula lutea*, and on log *Liriodendron Tulipifera*. A form with white spores, on roots fallen *Betula lenta*, Grant Co., near Bayard (*Millspaugh*). On *Fagus atropunicea*, Keeney's Creek, Oct. 20, 1893 (*Nuttall*, 1211).

**P. ABIETINUS** Fr.

On fallen *Tsuga Canadensis*, Grant Co., near Bayard (*Millspaugh*). On *Pinus Virginiana*, March 4, 1894 (*Nuttall*, 1420, 390).

## MUCRONOPORUS E. &amp; E.

**M. Everhartii** E. & Gall.

On *Ulmus pubescens*, Nov. 6, 1893 (*Nuttall*, 1598).

**M. Gilvus** (Sz) E. & E.

On *Castanea dentata*, July 20, 1894 (*Nuttall*, 1169, 558).

**M. ferruginosus** (Schrad.) E. & E.

On *Juglans cinerea*, Aug. 29, 1893 (*Nuttall*, 1183).

## PORIA Pers.

**P. vulgaris** Fr.

On various dead limbs, Feb. 2 and Nov. 20, 1893 (*Nuttall*, 833). A nodular form on oak ties, Nov. 5, 1894 (*Nuttall*).

**P. obducens** Pers.

On underside of dead log, Feb. 8, 1893 (*Nuttall*, 1297, 246).

**P. nigra** Berk.

On side and beneath dead *Hicoria?* log. Effused, long and narrow, perennial (*Nuttall*, 1268).

**P. tomento-cincta** B. & R.

Effused on bark underside of log *Betula lenta*, 60 x 30 cm. One layer taken off on Sept. 16, the same surface of the bark was found to be again covered with the species on Oct. 5, 1895 (*Nuttall*, 1865, 752). On *Magnolia Fraseri*, Nov. 1895.

**P. vaporaria** Fr.

On *Prunus scrotina*, Dec. 8, 1893 (*Nuttall*, 1295, 244).

**P. Tulipiferæ** Schw.

On *Liriodendron Tulipifera*, April 13, 1893 (*Nuttall*, 893).

**P. xantha** Fr.

On *Pinus Virginiana*, March 21, 1893 (*Nuttall*, 913).

**P. micans** (Ehrenb.) Fr.

On burned log, Short Creek, alt. 1,300 ft., Nov. 24, 1893  
(*Nuttall*, 1277, 223).

**P. corticola** Fr.

Underside of young dead *Liriodendron Tulipifera*, diffused through clefts of bark, incrusting other polypori, etc. Alt. 925 ft., Nov. 20, 1893 (*Nuttall*, 1280).

**P. sinuosa** Fr.?

Appears to be between this and *Irpex obliquus*, preferably this (E. & E.). On *Kalmia latifolia*. Short Creek, alt. 1,800 ft., Aug. 12, 1893 (*Nuttall*, 1166, 138).

**P. farinella** Fr.

On dead log, Short Creek, alt. 1,800 ft., Sept. 12, 1893 (*Nuttall*, 1204, 160).

TRAMETES Fr.

**T. Ohioensis** Berk.

On dead standing sapling, Short Creek, alt. 1,800 ft., April 10, 1894 (*Nuttall*, 1467).

**T. SEPIUM** Berk.

On dry railroad ties, Monongalia Co., near Morgantown (*Mills-  
paugh*). Resupinate form on dead oak beams over a boiler, Dec. 15, 1893 (*Nuttall*, 1300).

**T. mollis** Fr.?

On drift wood, March 1893 (undeveloped) (*Nuttall*, 1597, 550).

**T. odorata** Wulf.

On dead limbs in a drain, July 20, 1894 (*Nuttall*, 1628, 557).  
*Ozonium auricomum*, Link., a curious tawny mycelium, supposed to be intimately connected with this species was found here with this specimen.

**D. ambigua** Berk.

*Trametes ambigua* (Berk.) Fr. On dead *Acer* sp., Feb. 2, 1893  
(*Nuttall*, 846).

DÆDALIA Pers.

**D. confragosa** Pers.

On dead *Cornus florida*, Feb. 20, 1894 (*Nuttall*, 1774).

**D. unicolor** (Bull.) Fr.

On dead *Hicoria ovata*, Dec. 8, 1893 (*Nuttall*, 1298, 247).

MYRIADOPORUS Peck.

**M. INDURATUS** Peck.

Top of decayed oak stump, Monongalia Co., near Morgantown (*Mills-  
paugh*). Professor Peck remarks, in a letter, on receipt of the specimens: "This is probably only an imperfect condition of *Poria abducens*."

## FAVOLUS Fr.

**F. Europæus** Fr.

On dead branch, Feb. 12, 1894 (*Nuttall*, 1360). Spores 12.5 x 3  $\mu$  somewhat curved, obliquely apiculate. Feb. 17, 1895 (*Nuttall*, 1788).

## GLÆOPORUS Mont.

**G. CONCHOIDES** Mont.

On oak chips, Monongalia Co., near Morgantown (*Millsbaugh*).

## MERULIUS Hall.

**M. TREMELLOSUS** Schrad.

Under bark *Quercus alba* log, Monongalia Co., near Morgantown (*Millsbaugh*). On *Alnus rugosa*, Oct. 31, 1893 (*Nuttall*, 1220).

**M. rubellus** Peck.

On dead *Quercus alba*, Short Creek, Nov. 9, 1893 (*Nuttall*, 1244).

Also found parasitic on *Stereum versicolor*. In some cases the hymenium simply covers the under surface of the Stereum. In nearly all specimens observed it grows at the base of the host to which it is at least partly adherent. Largest pileus 5 x 5 cm., sometimes laterally connected for 15 cm.

**M. Corium** Fr.

On branches, April 27, 1895 (*Nuttall*, 1822, 699).

**M. molluscus** Fr.

On dead limb *Tsuga Canadensis*, Short Creek, alt. 1,800 ft., Feb. 8, 1894 (*Nuttall*, 1399).

**M. ambiguus** Berk.

On dead *Prunus serotina*, Short Creek, alt. 975 ft., Jan. 1894 (*Nuttall*, 1602).

**M. lachrymans** (Jacq.) Fr.

On bottom of box in cellar, Feb. 14, 1894 (*Nuttall*, 1379, 341).

**M. pulverulentus** Fr.

Underside of old logs, Oct. 19, 1893. Spores orange colored, nucleate, 10 x 6  $\mu$  (*Nuttall*, 1208).

## POROTHELIUM Fr.

**P. lacerum** Fr.

On dead trunk, Feb. 11, 1894 (*Nuttall*, 1381, 343).

## SOLENIA Hoffm.

**S. ochracea** Hoffm.

On dry dead log, Short Creek, alt. 1,800 ft., Dec. 16, 1893 (*Nuttall*, 1327; 285).

**S. villosa** Pers.

On fallen dead limb of *Quercus* sp., Sept. 1895 (*Nuttall*, 1854, 741).

HYDNACEÆ.

HYDNUM Linn.

**H. imbricatum** Linn.

In oak woods, June 18, 1893 (*Nuttall*, 988).

**H. adustum** Schwein.

On dead limbs on ground, Aug. 11, 1893. Nuclei at first cream-colored gradually changing from periphery to center through reddish purple to dark brown (*Nuttall*, 863).

**H. coralloides** Scop.

In decayed cleft on living *Fagus atropunicea*, Feb. 13, 1893 (*Nuttall*, 845).

**H. erinaceus** Bull.

On top dead log and in hollow of living *Fagus atropunicea*, alt. 1,800 ft., Nov. 27, 1894 (*Nuttall*, 1752).

**H. ochraceum** Pers.

Underside of dead log, April 7, 1893. *Stegonotis fusca* found growing upon this species (*Nuttall*, 891).

**H. croceum** (Schw.) Fr.

Effused in crevices of dead log, Short Creek, alt. 1,800 ft., March 6, 1894 (*Nuttall*, 1426, 398).

**H. cinnabarinum** (Schw.) Fr.

Effused under bark of dry dead limb, June 20, 1893 (*Nuttall*, 997).

**H.** ————— sp.

Underside old decayed log, Feb. 13, 1893. Effused 36 x 12 cm. or more (*Nuttall*, 844).

TREMELLODON Pers.

**T. gelatinosum** (Scop.) Pers.

On dead *Tsuga Canadensis* Aug. 13, 1893 (*Nuttall*, 1148).

IRPEX Fr.

**I. LACTEUS** Fr.

On dead *Rhus hirta*, Monongalia Co., near Morgantown (*Mills-  
paugh*).

**I. sinuosus** Fr.

On dead sticks, Dec. 10, 1893 (*Nuttall*, 1309, 255).

**I. obliquus** (Schrad.) Fr.?

On *Magnolia Fraseri*, Feb. 8, 1894 (*Nuttall*, 1385, 347).

RADULUM Fr.

**R. orbiculare** Fr.

On *Carpinus Caroliniana?* Nov. 27, 1893 (*Nuttall*, 1270).

**R. Magnoliæ** B. & C.

On *Magnolia Fraseri*, Feb. 9, 1894, Short Creek, alt. 1,800 ft. (*Nuttall*, 1352).

## PHLEBIA Fr.

**P. merismoides** Fr.

On mossy bark of dead log, Dec. 1, 1893 (*Nuttall*, 1289). On *Magnolia Fraseri*, Nov. 1895.

**P. radiata** Fr.

On *Alnus rugosa*, Oct. 31, 1893 (*Nuttall*, 1219).

## GRANDINIA Fr.

**G. granulosa** Fr.

On underside dead log, May 20, 1893 (*Nuttall*, 1304).

**G. crustosa** (Pers.) Fr.

On *Rhododendron maximum*, May 20, 1893. On *Liquidambar styraciflua*, Oct. 2, 1894 (*Nuttall*, 909 ident. Masee).

## ODONTIA Pers.

**O. farinacea** Fr.

Underside dead *Tsuga Canadensis*, Nov. 9, 1893, Short Creek, alt. 1,300 ft. (*Nuttall*, 1245).

## KNEIFFIA Fr.\*

**K. setigera** Fr.

In interstices dead log, Dec., 1893 (*Nuttall*, 1290, 231).

## THELEPHORACEÆ.

## THELEPHORA Ehrh.

**T. sebacea** Pers.

On stem of *Aster* sp., damp shady woods, July 25, 1893 (*Nuttall*, 1120).

**T. pedicellata** Schwein.

On *Ostrya Virginiana*, Feb. 8, 1894 (*Nuttall*, 1370, 328).

## STEREUM Pers.

**S. versicolor** (Sw.) Fr.

On *Quercus* sp. and *Betula lenta*, March 7, 1893 (*Nuttall*, 856).

**S. fasciatum** Sz.


In dead stumps, alt. 2,000 ft., March 8, 1893 (*Nuttall*, 860).

**S. VERSICOLOR** var. (Peck).

On dry oak railroad ties, Monongalia Co., near Morgantown (*Millsbaugh*).

\* A section of the genus *Enothera* has been elevated to generic rank by Raimann, in Engler & Prantl's *Natur. Pflanzenf.*, and Spach's *Kneiffia* (1835) re-instated to designate it. Fries' genus *Kneiffia* (1836) has, however, been so long established in mycologic literature, that we are loth to make any change at this time, feeling that the re-instating of Spach's *Kneiffia* is hardly proper, with Fries' genus well established.











- S. purpureum** Pers.  
On stump in pine woods (*Liriodendron?*) Dec. 9, 1893 (*Nuttall*, 1281, 391).
- S. hirsutum** (W.) Fr.  
On log, Nov. 24, 1893. Short Creek, alt. 1,300 ft. (*Nuttall*, 1277, 224).
- S. sanguinolentum** (A. & S.) Fr.  
On *Tsuga Canadensis*, Feb. 8, 1894 (*Nuttall*, 1386, 251).
- S. SULPHURATUM** B. & Rav.  
On log *Quercus alba*, Monongalia Co., near Morgantown (*Mills-  
paugh*).
- S. FRUSTULOSUM** (Pers.) Fr.  
On log *Quercus* sp., Monongalia Co., near Morgantown (*Mills-  
paugh*). On dead wood, March 7, 1893 (*Nuttall*, 857).
- S. Pini** Fr.  
On *Pinus Virginiana*, April 6, 1894 (*Nuttall*, 1469, 445).
- S. ochraceo-flavum** Schw.  
On small dead twigs, March 21, 1894 (*Nuttall*, 1416).
- S. albo-badium** Schwein.  
On dead sticks and twigs of various trees, April 27, 1893 (*Nutt-  
all*, 1269).
- S. SERICEUM** Schw.  
On dead twig *Betula lenta*, Grant Co., near Bayard (*Millspaugh*).  
On *Rhus copallina*, March 6, 1894 (*Nuttall*, 1419, 389).
- S. COMPLICATUM** Fr.  
On roots of overturned *Betula lenta*, Grant Co., near Bayard;  
on dry oak railroad ties, Monongalia Co., near Morgantown  
(*Millspaugh*). On dead trees and on *Quercus Prinus*, Dec. 8, 1893  
(*Nuttall*, 867, 235).
- S. versiforme** B. & C.?  
On *Castanea dentata*, Jan. 18, 1893 (*Nuttall*, 1347, 305).
- S. triste** B. & C.  
On *Rhododendron maximum* in thicket, alt. 1,800 ft. Feb. 8, 1894  
(*Nuttall*, 966, 350).
- S. acerinum** Pers.  
On living tree trunks, March 7, 1893 (*Nuttall*, 858).

HYMENOCHÆTE Lev.

- H. rubiginosa** (Schr.) Lev.  
On dead log, March, 1893 (*Nuttall*, 911).
- H. tabacina** (Sow.) Lev.  
On dead twigs, March 1893 (*Nuttall*, 910).

**H. CORRUGATA** (Fr.) Lev.

On decorticated limb *Betula lenta*, Grant Co., near Bayard (Millspaugh). On *Magnolia Fraseri*, Feb. 8, 1894. Setæ 38 x 7.5  $\mu$  (Nuttall, 1364).

**H. unicolor** B. & C.?

On *Betula lenta*, on dead *Kalmia latifolia*? July 1, 1893 (Nuttall, IIII, 96).

## CORTICIUM Fr.

**C. salicinum** Fr.

On dead *Salix nigra*, March 15, 1894 (Nuttall, 1401).

**C. Oakesii** B. & C.

On living *Ostrya Virginica*, March 21, 1893 (Nuttall, 877).

**C. globiferum** E. & E.

Underside of dead log in deep laurel thicket, Short Creek, March 25, 1893 (Nuttall, 940).

**C. arachnoideum** Berk.

On dead damp twigs in dense pile, Jan. 18, 1894. Diffused for 30 x 8 cm. Spores 4 to 5 x 2 to 3  $\mu$ . The species when fresh has a wrinkled or crusted appearance looking very like a *Merulius*, but soon becomes flat and plane (Nuttall, 1344, 303).

**C. radiosum** Fr.

On dead trunk, partly beneath loose bark, Short Creek, May 18, 1893 (Nuttall, 965).

**C. leve** Pers.

On *Hamamelis Virginica*, Dec. 20, 1894 (Nuttall, 1500).

**C. albo-flavescens** E. & E., sp. nov.

TYPE HABITAT: On bark of dead *Tsuga Canadensis*, Feb. 24, 1894 (Nuttall, discov. 1394).

At first orbicular, subpezizoid, 1 to 2 mm. diameter, becoming more irregular in shape and 1 cm. or more diameter, lower stratum milk-white, consisting of loosely interwoven, branching threads, and extending out beyond the pale-olivaceous or yellowish, membranaceous hymenium so as to form a thin subfimbriate, white margin. Spores abundant, globose, with a single large nucleus, 4 to 6  $\mu$  diameter, slightly colored.

The hymenium has only a slight tinge of olive and is perhaps better described as light-drab color.

**C. leptaleum** E. & E., sp. nov.

TYPE HABITAT: On under side dead sapling *Magnolia Fraseri*, alt. 1,800 ft. April 11, 1895 (Nuttall, discov. 1803, 690).

Thin, closely adnate, grayish-white, pulverulent, soon cracking into areas 2 to 3 mm across, with margins incurved so as to resemble the shallow cups of some *Peziza*. Spores oblong, hyaline, continuous, 10 to 12 x 4  $\mu$ , borne on cylindrical basidia 30 to 40 x 6  $\mu$ , with two stout sporophores. There are also other smaller

(immature)? sporidia, globose (4 to 6  $\mu$ ) or elliptical, 5 to 7 x 3 to 3.5  $\mu$ . The membrane on which the hymenium stands where exposed on the incurved margin of the pezizoid areas is pale brown.

**C. Petersii** B. & C.

On *Betula* sp. and *Vitis* sp., March 21, 1893 (*Nuttall*, 912).

**C. colliculosum** B. & C.

On dead logs in laurel thicket, Short Creek, May 1893 (*Nuttall*, 1088).

**C. prasinum** B. & C.

On dead burned log, Feb. 2, 1894 (*Nuttall*, 1382, 344).

**C. lilacino-fuscum** B. & C.

On dead limb *Hicoria* sp., Short Creek, alt. 1,800 ft., March 5, 1894 (*Nuttall*, 1403).

**C. ochraceum** Fr.

On underside dead log *Tsuga Canadensis*, May 8, 1893 (*Nuttall*, 931).

**C. incarnatum** (Pers.) Fr.

On underside dead log and on *Ulmus pubescens*, April 18, 1893 (*Nuttall*, 923).

**C. lilacinum** Schroet.

On firm wood of dead log, March 6, 1894, Short Creek, alt. 1,800 ft. (*Nuttall*, 1427, 399).

**C. subgiganteum** Berk.

On *Hicoria ovata*, Dec. 14, 1893. Spores 16 x 5  $\mu$  (*Nuttall*, 1312, 262).

**C. martianum** B. & C.

On underside of wet log, Dec. 16, 1893. When fresh this species is the color of fresh blood. Short Creek, alt. 1,800 ft. (*Nuttall*, 1324, 281).

**C. scutellare** B. & C.

On *Magnolia Fraseri*, *Sambucus Canadensis* and *Pinus Virginiana*, Aug. 18, 1893 (*Nuttall*, 1176, 144).

ASTEROSTOMA Masee.

**A. corticolum** Masee.

On wild vine *Vitis* sp., in pine woods, alt. 2,000 ft., Feb. 1893 (*Nuttall*, 1000).

**A. cervicolor** (B. & C.) Masee.

On bark of living *Quercus* sp., Short Creek, alt. 1,800 ft. (*Nuttall*, 1325, 282).

PENIOPHORA Cooke.

**P. quercina** (Fr.) Cooke.

On *Castanea dentata*, Jan. 18, 1893 (*Nuttall*, 1346, 305).

**P. cinerea** (Fr.) Cooke.

On dead limb, March 1893 (*Nuttall*, 945, 170).

## CONIOPHORA DC.

**C.** ————— sp.

Near *C. submembranacea* (B. & Br.). On *Zea Mays*, Aug. 10, 1895 (*Nuttall*, 1846, 737).

**C. olivacea** (Fr.) Karst.

In dark place underside old railroad ties, Nov. 5, 1894 (*Nuttall*, 1750, 631).

## EXOBASIDIUM Woron.

**E. RHODODENDRI** Cram.

Forming "cups" near the tips or margins of living leaves of *Rhododendron maximum*. Common in Grant and Tucker Counties. I understand from Prof. Peck that this is his first knowledge of the occurring of this species in North America (*Mills-paugh*).

## CLAVARIACEÆ.

## CLAVARIA Vaill.

**C. amethystina** Bull.

On ground, June 20, 1893 (*Nuttall*, 993).

**C. pyxidata** Pers.

On rotten log, June 10, 1893 (*Nuttall*, 975).

**C. formosa** Pers.

On ground in laurel thicket, and deep woods, alt. 2,000 ft., July 9, 1893. Spores  $12.5 \times 5 \mu$  (*Nuttall*, 1085).

**C. FLACCIDA** Fr.

On leaf mold, in deep woods, Grant Co., near Bayard (*Mills-paugh*).

**C. fusiformis** Sowerb.

On low ground in open woods, alt. 2,000 ft., July 9, 1893 (*Nuttall*, 1086).

**C. juncea** Fr.

On leaf mold in bottom of dead hollow stump. Oct. 28, 1893 (*Nuttall*, 1232, 179).

**C. mucida** Pers.

On upper side dead wet logs, Oct. 20, 1893 (*Nuttall*, 1231, 178).

## CALOCERA Fr.

**C. viscosa** (Pers.) Fr.

On bark of *Magnolia Fraseri*, Short Creek, alt. 1,800 ft., Feb. 8, 1894. Spores  $12.5 \times 4 \mu$  (*Nuttall*, 1387, 354).

LACHNOCLADIUM Lev.

**L. semivestitum** B. & C.

On dead wet limbs, alt. 1,800 ft., Sept. 18, 1895. Three to five inches high (*Nuttall*, 1867, 753).

TREMELLINACEÆ.

HIRNEOLA Fr.

**H. Auricula-Judæ** (Linn.) Berk.

On damp sticks, June 15, 1893, Short Creek (*Nuttall*, 993).

EXIDIA Fr.

**E. glandulosa** (Bull) Fr.

On bark *Juglans cinerea*, alt. 2,000 ft., Feb. 20, 1893 (*Nuttall*, 946).

**E. alba** (Huds.) Bref.

On dead log, June 23, 1893 (*Nuttall*, 999).

ULOCOLLA Bref.

**U. foliacea** (Pers.) Bref.

On dead twig, Nov. 10, 1893. Spores + 12.5 x 6 to 7  $\mu$  (*Nuttall*, 1281).

TREMELLA Dill.

**T. frondosa** Fr.

On *Rhododendron maximum*, Dec. 16, 1893 (*Nuttall*, 1321).

**T. aurantia** Fr.

On bark *Pinus Virginiana*, alt. 2,000 ft., March 20, 1893 (*Nuttall*, 947).

**T. intumescens** Sm.?

On dead limbs *Hicoria* sp., Short Creek, Dec. 16, 1893. Spores 10 to 13 x 3 to 4  $\mu$ , curved (*Nuttall*, 1332).

**T. virens** Schwein.

On dead branch *Cornus florida*, Aug. 17, 1893 (*Nuttall*, 1155). Spores mostly 7 x 5  $\mu$  or round 7  $\mu$  diameter.

NÆMATELIA Fr.

**N. nucleata** (Schw.) Fr.

On *Aralia spinosa*, alt. 1,300 ft., Aug. 18, 1893 (*Nuttall*, 1174, 142). On *Magnolia Fraseri*.

DACRYOMYCES Nees.

**D.** \_\_\_\_\_ sp.

Of this form Mr. J. B. Ellis says: "It seems to be a *Dacryomyces*. I find the same thing here at Newfield, N. J., but have never been able to decide upon it, and though I have sent it to

various European mycologists, I have never received a definite opinion from them." The form occurs on oak limbs on ground, Dec. 1893; is of a bright orange-red color, with curved spores 7.5 to 10 x 4  $\mu$  (Nuttall, 1271, 217).

**D. deliquescens** (Bull.) Dub.

On underside dead rotten limb *Pinus Strobus*, Dec. 13, 1894 (Nuttall, 1770).

**D. Syringæ** (Schum.)?

On dead limbs of *Platanus occidentalis*, Nov. 10, 1895 (Nuttall, 1889).

**D. stellatus** Nees.

On clean dead wood *Tsuga Canadensis*, Dec. 20, 1893 (Nuttall, 1331).

**D. corticioides** Ell.

On decayed *Tsuga Canadensis*, Short Creek, alt. 1,750 ft., Dec. 16, 1893. Lemon yellow .25 cm. to 3.5 x 1.5 cm. Spores hyaline, nucleate 9 to 12.5 x 5 to 7  $\mu$  obliquely apiculate at one end (Nuttall, 1316, 271).

**D. minor** Peck?

On *Kalmia latifolia*, Short Creek, alt. 1,800 ft., March, 1894. Spores 10 to 12 x 4  $\mu$  (Nuttall, 1532).

GUEPINIA Fr.

**G. spathularia** (Schw.) Fr.

On *Quercus* sp., Jan. 1, 1894. Spores uniseptate 10 x 4  $\mu$  (Nuttall, 1782).

HORMOMYCES Bon.

**H. FRAGIFORMIS** Cooke.

On bark dead *Fagus atropunicea*, Grant Co., near Bayard (Mills-paugh). On dead wood, May 18, 1893 (Nuttall, 963). *Hypsilo-phora fragiformis* (Fr.) Lev.



# Lichenes.

## USNACEÆ.

### RAMALINA Ach.

**R. rigida** (Pers.) Tuck.

On trees, alt. 2,000 ft. (*Nuttall*, 1060).

**R. calcicaris fastigiata** Fr.

On trees, alt. 2,000 ft. (*Nuttall*, 869).

**R. miniuscula** Nyl.

On *Acer saccharinum*, alt. 2,000 ft. (*Nuttall*, 1696).

### CETRARIA Ach.

**C. ciliaris** (Ach.) Tuck.

On *Pinus Virginiana*, alt. 1,800 ft. (*Nuttall*, 861).

**C. lacunosa** Ach.

On *Pinus Virginiana*, alt. 1,800 ft. (*Nuttall*, 1053).

**C. aurescens** Tuck.

On *Rhododendron maximum*, alt. 1,850 ft. (*Nuttall*, 1050).

### USNEA Ach.

**U. BARBATA** (L.) Fr.

Wirt Co., on old trees, common, Burning Springs. Randolph Co., on *Rhododendron maximum*, common, Cheat Bridge. Mercer Co., on oak twigs, Bluefield; and elsewhere about State, common on trees, rocks and old fence rails (*Millsbaugh*). On trees, alt. 2,000 ft. (*Nuttall*, 819).

**U. BARBATA FLORIDA** Fr.

Mercer Co., on oak chips and twigs among dead leaves at Bluefield (*Millsbaugh*). On trees, alt. 2,000 ft. (*Nuttall*, 859).

### THELOSCHISTES Norm.

**T. parietinus** (Linn.) Norm.

On *Robinia Pseudacacia*, alt. 2,000 ft. (*Nuttall*, 1145).

**T. concolor** (Dicks.) Tuck

On *Robinia Pseudacacia*, alt. 2,000 ft. (*Nuttall*, 1144).

## PARMELIÆ.

## PARMELIA Ach.

**P. perlata** (Linn.) Ach.

On *Acer rubrum*, and on sandstone boulders (*Nuttall*, 848).

**P. Borreri** Turn.

On oak, alt. 1,850 ft. (*Nuttall*, 1032).

## P. BORRERI RUDECTA Tuck.

On *Robinia Pseudacacia*, Monongalia Co., along Falling Run (*Millspaugh*, 1531).

**P. colpodes** (Ach.) Nyl.

On oak, alt. 1,850 ft. (*Nuttall*, 1141).

## P. OLIVACEA (L.) Ach.

On *Liriodendron* log newly felled, Monongalia Co., Falling Run (*Millspaugh*, 1343).

## P. CAPERATA (L.) Ach.

Monongalia Co., on sandstone rocks, and base of beech, Falling Run (*Millspaugh*, 1283).

## PHYSCIA DC.

## P. LUCOMELA (L.) Michx.

Mercer Co., with moss on wet limestone ledge, Beaver Spring (*Millspaugh*, 1539).

**P. ciliaris** (Linn.) DC.

On mossy sandstone rocks (*Nuttall*, 1063).

**P. ciliaris crinalis** Schær.

On moss, alt. 1,850 ft. (*Nuttall*, 1042).

**P. aquila** (Ach.) Nyl.

On rocks and on moss (*Nuttall*, 1035).

**P. stellaris** (Linn.) Tuck.

On sandstone rock (*Nuttall*, 872).

**P. stellaris aipolia** Nyl.

On sandstone rock (*Nuttall*, 1778).

## UMBILICARIÆ.

## UMBILICARIA, Hoffm.

**U. Dillenii** Tuck.

On sandstone rocks (*Nuttall*, 823).

**U. Pennsylvanica** Hoffm.

On sandstone rocks (*Nuttall*, 849).

**U. pustulata** (Linn.) Hoffm.

On faces of sandstone cliffs (*Nuttall*, 855).

PELTIGERIEÆ.

STICTA Schreb.

- S. HERBACEA** (Huds.) Ach.  
 McDowell Co., on oaks, Elkorn (*Millsbaugh*).
- S. erosa** (Eschw.) Tuck.  
 On trees, alt. 1,850 ft. (*Nuttall*, 982).
- S. PULMONARIA** (L.) Ach.  
 Wirt Co., on trunks of oaks, near Burning Springs (*Millsbaugh*, 327). On trees, alt. 1,875 ft. (*Nuttall*, 824).
- S. quercizans** (Michx.) Ach.  
 On trees and rocks (*Nuttall*, 1057).

PELTIGERA Willd.

- P. APHTHOSA** (K.) Hoffm.  
 Monongalia Co., on rock ledge, near Cassville (*Millsbaugh*).  
 On rocks among mosses, alt. 1,800 ft. (*Nuttall*, 1090).
- P.** ————— sp. *non fruct.*  
 On *Rhododendron maximum* (*Nuttall*, 938).
- P. horizontalis** (Linn.) Hoffm.  
 On rocks among mosses, alt. 1,875 ft. (*Nuttall*, 1091).
- P. canina** (Linn.) Hoffm.  
 On earth, rocks and trees (*Nuttall*, 1061).

PANNARIEÆ.

PANNARIA Delis.

- P. rubiginosa** (Thunb.) Delis.  
 On mossy logs (*Nuttall*, 1055).
- P. leucosticta** Tuck.  
 On mossy logs and rocks (*Nuttall*, 1064).
- P. nigra** (Huds.) Nyl.  
 On sandstone rocks (*Nuttall*, 1077).

COLLEMEÆ.

COLLEMA Hoffm.

- C. myochroum** Nyl.  
 On *Rhododendron maximum*, alt. 1,875 ft. (*Nuttall*, 1058).

LEPTOGIUM Fr.

- L. pulchellum** (Ach.) Nyl.  
 On oak (*Nuttall*, 1067).

- L. tremelloides** (Linn. f.) Fr.  
On sandstone rock (*Nuttall*, 1093).
- L. chloromelum stellans** Tuck.  
On *Quercus coccinea*, alt. 2,400 ft. (*Nuttall*, 1146).

## LECANOREÆ.

## PLACODIUM DC.

- P. elegans** (Link.) DC.  
On sandstone boulders at base of cliffs, alt. 1,800 ft. (*Nuttall*, 862).
- P. CERINUM** (Hedw.) Naeg. & Hepp.  
Monongalia Co., on bark of beech, Falling Run (*Millspaugh*, 1357). On *Robinia Pseudacacia*, alt. 2,000 ft. (*Nuttall*, 1030).
- P. ferrugineum** (Huds.) Hepp.  
On top of wet sandstone rock (*Nuttall*, 1092).
- P. camptidium**, Tuck.  
On trees (*Nuttall*, 1786).

## LECANORA, Ach.

- L. coarctata** Ach.  
On pieces of sandstone under drip of wet cliff, alt. 1,800 ft. (*Nuttall*, 1780).
- L. subfusca** (Linn.) Ach.  
On trees and rocks (*Nuttall*, 1038).
- L. ATRA** (Huds.) Ach.  
Monongalia Co., on bark *Liriodendron* log, newly felled, Falling Run (*Millspaugh*, 1342). On flat exposed surface sandstone rock, same locality (1287).
- L. deplanans** Nyl.  
On rocks in bed of creek, alt. 1,300 ft., Fayette Co. in Short Creek (*Nuttall*, 1126).
- L. varia** (Ehrh.) Nyl.  
On trees (*Nuttall*, 1036).
- L. pallescens** (Linn.) Schær.  
On trees and rails (*Nuttall*, 937).
- L. tartarea** (Linn.) Ach.  
On sandstone rocks (*Nuttall*, 886).
- L. cervina** (Pers.) Nyl.  
The light colored form on wet rocks; the dark form on dry rocks (*Nuttall*, 854).
- L. privigna** (Ach.) Nyl.  
On sandstone rocks (*Nuttall*, 1028).

**Lecanora deplanans** Nyl., sp. nov.—Thallus glauco-cinerascens tenuis areolato-rimosus determinatus; apothecia badio-rufescens (satis diluta) innata subconcaviuscula, latit. 0.5–0.7mm; spora ellipsoidea, long. 0.015–16, crass. 0.009–1.010mm; epithecium inspersum. In toto gelatina hymenialis fulvo-rubescens.

Videtur species e stirpe *Lecanora cervina*, spermatiis ellipsoideis.

On rocks in bed of creek. West Virginia, Short Creek, alt. 1300 feet, L. W. Nuttall coll. no. 1126.

(INSERT OPPOSITE PAGE 178).



RINODINA Mass.

**R. oreina** (Ach.) Mass.

On sandstone rocks at base of cliff, alt. 1,800 ft. (*Nuttall*, 952).

**R. sophodes** (Ach.) Nyl.

On small sandstone rocks (*Nuttall*, 1072).

PERTUSARIA DC.

**P. velata** (Turn.) Nyl.

On *Magnolia Fraseri* and other trees (*Nuttall*, 1044).

**P. multipuncta** (Turn.) Nyl.

On sandstone rocks in bed of creek (*Nuttall*, 1043).

**P. leioplaca** (Ach.) Schaer.

On oak (*Nuttall*, 1041).

**P. pustulata** (Ach.) Nyl.

On trees (*Nuttall*, 853).

THELOTREMA Ach.

**T. subtile** Tuck.

On trees (*Nuttall*, 1056).

MYRIANGIUM Mont. & Berk.

**M. Duriæi** (Mont. & Berk.) Tuck.

On dead bush (*Nuttall*, 1062).

CLADONIEÆ.

CLADONIA Hoffm.

**C. MITRULA** Tuck.

Monongalia Co., on an old beech log, Falling Run (*Millsbaugh*, 1346).

**C. cariosa** (Ach.) Spreng.

On earth, Fayette Co. (*Nuttall*, 1066).

**C. PYXIDATA** (L.) Fr.

Monongalia Co., along Falling Run, on bare sandstone rocks (*Millsbaugh*, 1281); on moss in clay soil (*Millsbaugh*, 1285); on decayed log (*Millsbaugh*, 1338); base of beech in soil (*Millsbaugh*, 1282); on earth (*Nuttall*, 1031).

**C. fimbriata** (Linn.) Fr.

On an old log, Fayette Co. (*Nuttall*, 1059).

**C. fimbriata tubæformis** Fr.

On an old log (*Nuttall*, 1059).

**C. GRACILIS** (L.) Nyl.

Monongalia Co., along Falling Run, on decayed log (*Millsbaugh*, 1337); among mosses on clay soil (*Millsbaugh*, 1286).

- C. gracilis verticillata** Fr.  
On earth (*Nuttall*, 885).
- C. sobolescens** Nyl.  
On earth (*Nuttall*, 839).
- C. papillaria** (Ehrh.) Hoffm.  
On earth (*Nuttall*, 1068).
- C. delicata** (Ehrh.) Fl.  
On top of stump (*Nuttall*, 1065).
- C. cæspiticia** (Pers.) Fl.  
On sandstone rocks (*Nuttall*, 1075).
- C. FURCATA RACEMOSA** Floerk.  
Monongalia Co., large patches on ground under chestnuts, at Dille's; Mercer Co., same growth under oaks, near Beaver Spring (*Millsbaugh*) on earth (*Nuttall*, 1045).
- C. RANGIFERINA** (L.) Hoffm.  
Monongalia Co., on moss, Falling Run (*Millsbaugh*, 1361); on earth, rocks and rails (*Nuttall*, 825).
- C. macillenta** (Ehrh.) Hoffm.  
On rotten logs (*Nuttall*, 1070).
- C. pulchella** Schwein.  
On dead wood (*Nuttall*, 838).
- C. CRISTATELLA** Tuck.  
Monongalia Co., on an old decayed log, Falling Run (*Millsbaugh*, 1336).

## LECIDEÆ.

## BÆOMYCES Pers.

- B. roseus** Pers.  
On ground (*Nuttall*, 873).

## BIATORA Fr.

- B. russula** (Ach.) Mont.  
On bark of tree (*Nuttall*, 1046).
- B. varians** (Ach.) Tuck.  
On bark of tree (*Nuttall*, 978).
- B. uliginosa** (Schrad.) Fr.  
On rotten stumps (*Nuttall*, 1054).
- B. fuliginosa** Ach.  
On ground (*Nuttall*, 1089).
- B. Schweinitzii** Tuck.  
On *Rhododendron maximum* (*Nuttall*, 1052).
- B. rubella** (Ehrh.) Rab.  
On mossy trunk of tree (*Nuttall*, 1029).





**Lecidea Nuttallii** Calk. & Nyl., sp. nov.—Apothecia nigra parva; epithecium impressum; spora fusca oviformis 1-septata, long 0.014–16, crass. 0.005–6mm; hypothecium fuscum. In toto gelatina hymenialis vinose rubescens.

Super thallum *Ricasolia sublævis* Nyl. West Virginia, near Nuttallburg, L. W. Nuttall coll. no. 1781.

**Lecidea Virginiensis** Calk. & Nyl., sp. nov.—Thallus glaucescens tenuis laevigatus rimulosus citriño-flavus; apothecia fusca aut nigra convexiuscula immarginata latit. circ. 0.5mm, intus medio sordida; spora oblonga incoloris, long. 0.009–0.012, crass. 0.004–6mm; epithecium et pars media hypothecii dilute fuscescens. In toto gelatina hymenialis coerulescens, dein theca vinose rubescens.

E stirpe videtur *Lecidea sanguineo-atra*, prope *Lecideam delineatam* Nyl.

On sandstone rock under the drip of a wet cliff. West Virginia, near Nuttallburg, L. W. Nuttall coll. no. 1779.

(INSERT OPPOSITE PAGE 181).

**B. inundata** Fr.

On sandstone rock (*Nuttall*, 1074).

**B. suffusa** Fr.

On bark of tree (*Nuttall*, 1095).

LECIDEA Ach.

**L. Tennesseeensis** Nyl.

On sandstone rock (*Nuttall*, 1073).

**L. fuliginea** Ach.

On upper side of dead log (*Nuttall*, 1787).

**L. Nuttallii** Calkins. mss.

Parasitic on *Sticta erosa* (*Nuttall*, 1781).

**L. Virginiensis** Calkins. mss.

On sandstone rock under drip of wet cliff (*Nuttall*, 1779).

**L. ericina** Nyl.

Bark of trees (*Nuttall*, 1047).

BUELLIA DeNot.

**B. myriocarpa** (DC.) Mudd.

On bark of trees (*Nuttall*, 1185).

**B. petræa** (Fl.) Tuck.

On sandstone rocks (*Nuttall*, 1071).

GRAPHIDACEÆ.

OPEGRAPHA Humb.

**O. varia** (Pers.) Fr.

Bark of trees (*Nuttall*, 1034).

**O. atra** Pers.

Bark of trees (*Nuttall*, 1033).

GRAPHIS Ach.

**G. scripta** (Linn.) Ach.

On bark of trees (*Nuttall*, 1051).

**G. scripta recta** Schær.

On bark of *Betula lutea* and of *B. lenta* (*Nuttall*, 1026).

**G. elegans** (Sm.) Ach.

On bark of trees, Fayette Co. (*Nuttall*, 1039).

**G. dendritica** Ach.

On bark of *Betula lenta*, etc. (*Nuttall*, 1037).

ARTHONIA Ach.

**A. cinnabarina** Wallr.

Bark of tree (*Nuttall*, 939).

- A. aleuromela** Nyl. mss.  
On bark of oak, alt. 2,000 ft. (*Nuttall*, discov. 1182).
- A. dispersa** (Schrad.) Nyl.  
On bark of young *Tsuga Canadensis*, etc. (*Nuttall*, 1142).
- A. astroidea** Ach.  
On bark of *Liriodendron Tulipifera* (*Nuttall*, 1816).
- A. punctiformis** Ach.  
On bark (*Nuttall*, 1076).
- A. tædiosa** (Nyl.) Fr.  
On bark (*Nuttall*, 1069).
- A. spectabilis** Fl.  
On bark (*Nuttall*, 1025).

## AGYRIUM Fr.

- A. rufum** (Pers.) Fr.  
On smooth dead wood (*Nuttall*, 1814).

## CALICIACEÆ.

## CALICIUM Pers.

- C. subtile** Fr.  
On sticks, dead leaves, etc., under cliff (*Nuttall*, 1226).

## VERRUCARIACEÆ.

## ENDOCARPON Hedw.

- E. miniatum** (Linn.) Schær.  
On wet cliffs (*Nuttall*, 1049).
- E. miniatum complicatum** Schær.  
On wet cliffs (*Nuttall*, 1823).

## TRYPETHELIUM Spreng.

- T. virens** Tuck.  
On *Ilex opaca* (*Nuttall*, 1040).
- T. exocanthum** Tuck.  
On *Fagus atropunicea* (*Nuttall*, 1181).

## PYRENULA Ach.


- P. rhyponia** Ach.  
On bark of *Hamamelis Virginica* (*Nuttall*, 1184).
- P. nitida** Ach.  
On *Betula lenta* bark (*Nuttall*, 1027).
- P. pachycheila** Tuck.  
On bark of *Fagus atropunicea* (*Nuttall*, 1048).

**Arthonia aleuromela** Nyl., sp. nov.—Thallus albus subfarinaceus chrysogonicus tenuissimus; apothecia subrotundata vel oblonga, prominula, latit. 0.4–0.5mm; spora oblongo-oviformis parte inferiore attenuata, 1-septata, long. 0.010–11, crass. 0.003mm. In toto gelatina hymenialis cœrulescens, dein obscurata.

Thallus detritus subaureus, CaCl vix reagens. Gonidia chroolepoidea fulvescentia.

On bark of *Quercus* sp. West Virginia, near Nuttallburg, alt. 2000 feet, L. W. Nuttall coll. no. 1182.

(INSERT OPPOSITE PAGE 182).



## Hepaticae.

---

### JUNGERMANNIACEÆ.

FRULLANIA ASA-GRAYANA Mont.

**Frullania plana** Sull.  
(*Nuttall*).

JUBULA HUTCHINSIÆ SULLIVANTII Spruce.

### ARCHILEJUNEA Spruce.

A. CALYPEATA (Schw.) Spruce.  
*Lejunea calypeata* Schw. (*Nuttall*).

RADULA XALAPENSIS Mont.

RADULA TENAX Lindb.

### BELLINCINIA Raddi.

B. PLATYPHYLLA (L.) O. K.  
*Porella platyphylla* Lindb.

B. PORELLA (Dicks.) O. K.  
*Porella pinnata* Schw.

TRICHOCOLEA TOMENTELLA (Ehrh.) Dumort.

HERBERTIA ADUNCA (Dicks.) S. F. Gray.

BAZZANIA TRILOBATA (L.) S. F. Gray.

BAZZANIA DEFLEXA (Mart.) Underw.

**Cephalozia media** Lindb.  
*Cephalozia multiflora* Spruce (*Nuttall*).

CEPHALOZIA CURVIFOLIA (Dicks.) Dumort.  
(*Nuttall*).

ODONTOCHISMA SPHAGNI (Dicks.) Dumort.  
(*Nuttall*).

BLÉPHAROSTOMA TRICHOPHYLLUM (L.) Dumort.

KANTIA TRICHOMANIS (L.) S. F. Gray.

GEOCALYX GRAVEOLENS (Schrad.) Nees.

**Scapania undulata** (L.) Dumort.

Randolph Co., at Pickens, on clay at the banks of a spring  
(*Millsbaugh*).

SCAPANIA NEMOROSA (L.) Dumort.

DIPLOPHYLLUM TAXIFOLIUM (Wahlenb.) Dumort.

PLAGIOCHILA VIRGINICA, A. W. Evans, Fl. W. Va., 497 (1892) Plate.

Growing in wide, depressed, and intricate tufts; stems ascending from a prostrate caudex, simple or sparingly branched, sometimes geniculate and rooting at the joints, otherwise eradicate; leaves contiguous or somewhat imbricated, widely patent, ovate or rhomboid-ovate, the dorsal margin decurrent, slightly reflexed, entire, the ventral margin plane or reflexed at base, mostly entire, the apex broad, rounded or truncate, sharply and irregularly spinulose; amphigastria none.

Stems 1 to 3 cm. long, with the leaves 1 to 2 mm. wide; leaves 1.2 mm. long, 0.7 mm. wide; spines short, acute, separated by rounded sinuses, varying in number from 2 to 8 on each leaf, usually 4 or 5; leaf-cells averaging 0.023 mm. in diameter in middle of leaf, thin-walled and scarcely thickened at the angles.

Mercer: on walls of dry limestone cave, Beaver Spring (1550).

PLAGIOCHILA PORELLOIDES Lindenb.

HARPANTHUS SCUTATUS (Web. & Mohr.) Spruce.

JUNGERMANNIA EXSECTA Schmid.

#### MARSUPELLA Dumort.

**M. emarginata** (Ehrh.) Dumort.

Randolph Co., at Pickens, on clay bank of a small spring.  
(*Millsbaugh*)

PELLIA EPIPHYLLA (L.) Dumort.

METZGERIA CONJUGATA Lindb.

(*Nuttall*).

ANEURA MULTIFIDA (L.) Dumort.

**Aneura sessilis** Spreng.

(*Nuttall*).

#### MARCHANTIACEÆ.

MARCHANTIA POLYMORPHA Linn (*Nuttall*).

CONOCYPHALUM CONICUM (L.) Dumort.



## Musci.

### POLYTRICHACEÆ.

- POLYTRICHUM COMMUNE Linn.  
POLYTRICHUM OHIENSE Ren. & Card.  
POLYTRICHUM PILIFERUM Schreb.  
POLYTRICHUM TENUE (Menz.) E. G. Britton.

### DIPHYSCIUM, Mohr.

- D. foliosum** Mohr (*Nuttall*).

### BUXBAUMIA Haller

- B. aphylla** Linn (*Nuttall*).

- CATHARINEA ANGUSTATA Brid.  
CATHARINEA UNDULATA Web. & Mohr.

### GEORGIACEÆ.

- GEORGIA PELLUCIDA (L.) Rabenh.

### FISSIDENTACEÆ.

- FISSIDENS ADIANTOIDES (L.) Hedw.  
FISSIDENS DECIPIENS De Not.

### MNIACEÆ.

- ASTROPHYLLUM SYLVATICUM Lindb.  
ASTROPHYLLUM ROSTRATUM Schrad.  
ASTROPHYLLUM PUNCTATUM (L.) Lindb.  
ASTROPHYLLUM HORNUM (L.) Lindb.  
SPHÆROCEPHALUS HETEROSTICHUS (Brid.) E. G. Britton.

### BARTRAMIACEÆ.

- BARTRAMIA POMIFORMIS (L.) Hedw.  
On ground, Greenbrier Co., Kate's Mountain, alt. 3,300 ft.  
(*Small & Vail*).  
BARTRAMIA POMIFORMIS CRISPA (Sw.) Schimp.  
BARTRAMIA FONTANA (L.) Brid.

## BRYACEÆ.

- BRYUM BIMUM Schreb.  
 BRYUM ARGENTEUM Linn.  
 BRYUM PROLIFERUM (L.) Sibth.  
 LEPTOBRYUM PYRIFORME (L.) Wils.

## FUNARIACEÆ.

- FUNARIA HYGROMETICA (L.) Sibth.  
 FUNARIA HYGROMETICA PATULA Br. & Sch.  
 FUNARIA FLAVICANS Michx.  
 PHYSCOMITRIUM PYRIFORME (L.) Brid.

## TORTULACEÆ.

- LEERSIA STREPTOCARPA (Hedw.) Lindb.  
 TORTULA MURALIS (L.) Hedw.  
 BARBULA HUMILIS Hedw.  
 BARBULA TORTUOSA (L.) Web. & Mohr.  
 MOLLIA VIRIDULA (L.) Lindb.

## DICRANACEÆ.

- LEUCOBRYUM GLAUCUM (L.) Br. & Sch.  
 DICRANODONTIUM VIRGINICUS, E. G. Britton, Fl. W. Va., 488 (1892)  
 Plate:

Monongalia: On sandstone boulder along a woodland path, Tibb's Run (*Millspaugh*, 1635).

Plants bright glossy green, stems matted below by a red tomentum, leafy nearly to apex, denudate roughened above, with a few leaves at summit; leaves erect or secund, straight or curled and twisted, often 5 mm. long, narrowly subulate, from a short, thick base, caducous ones with a long, slender, smooth point; persistent ones serrate, blade inflexed, cells densely chlorophyllose, filled with oil globules, those of the basal angles clear. Dioecious, the antheridia terminal in conspicuous heads, bracts brown at base, apex subulate, serrate; perichætal bracts 3-4 mm. long, from a short base, suddenly subulate, dentate at apex; pedicels lateral by the growth of innovations, 1½-2 cm. long, pale, glossy yellow, twisted in two directions, very slender, arcuate when young, becoming erect before capsules mature. Capsule cylindrical, ribbed only at the mouth, 1½-2 mm. long, beak straight or curved, shorter than the capsule, peristome bright red, *not deep set*, teeth split unequally to middle, striolate at base, pale and granulose above, annulus none, spores small, calyptra cucullate, 2 mm. long, beaked, entire. Maturing in summer.

Differs from European specimens of *D. longirostre* collected by Seringe; in the longer, paler, more slender, scarcely arcuate pedicels, longer capsules, peristome not deep set, and teeth split only to the middle, more united than figured in the *Bryologia*

Europea, Table 88. It may be distinguished from *Campylopus Virginicus*, also remarkable for its caducous leaves, by the longer, more slender subulate point, which is entire or minutely serrate and smooth on the back, by the thick base, with inflexed blades, and by the shape of the basal cells at the angles.

**DICRANODONTIUM MILLSPAUGHII**, E. G. Britton, Fl. W. Va., 488 (1892) Plate.

Monongalia: on sandstone boulder, deep woods, Tibb's Run (*Millsbaugh*, 1596).

Plant slight yellowish green, silky, caespitose; stems matted with rufous tomentum at base, 1-3 cm. long, a few denudate, roughened by the fragments of the slightly caducous leaves. Leaves secund or erect-spreading, 4-5 mm. long, narrowly subulate from a broad base 1-1½ mm. long, becoming tubular above with inrolled margins, basal angles not auricled, filled by large hyaline cells to the base of the broad, brown vein, those of the blade oblong or square next the vein, becoming spindle-shaped and prosenchymatous toward the margin, vein thick, excurrent into a dentate slender tip, rough on back. Dioecious, perichæcium 5-7 mm. long, bracts sheathing half their length, tapering to a long, slender, obscurely serrate tip, outer shorter, abruptly subulate, more sharply serrate; pedicels recurved, burying the capsules among the leaves, becoming erect when old, 5-8 mm. long, stout and twisted in two directions; capsules pyriform-cylindric with a distinct neck, length about 1 mm. without the lid, which is as long as the rest of the capsule, with a straight beak; calyptra cucullate, entire; peristome red, connivent, teeth deep set, slender, split to middle, or perforate to base, striolate below, granulose above; annulus none, mouth bordered by a dense, dark rim. Maturing in summer, old capsules persistent, not sulcate.

Differs from European specimens of *D. longirostre* in the structure of the base of the leaf, lacking the suddenly inflated basal auricles; differing also in the cells above the base, teeth not split to base, occasionally only perforate. From *D. Virginicus* it may be distinguished by the less caducous leaves, shorter, stouter, more arcuate pedicels, smaller capsules, and longer sheathing perichæcium.

Through the kindness of Dr. Robinson I have been able to compare these specimens with those collected by Sullivant on Grandfather Mt. in 1843. His also are fruiting, and an excellent drawing is preserved, hence I am able to assert that the specimens are almost identical. Sullivant's showing no naked stems, but many of the leaves are caducous. Dr. Braithwaite kindly compared the West Virginia specimens with *Campylopus pyriformis*, sending me specimens of this and the variety *Mulleri*, and sketches of the bases of the leaves. It is evident that Sullivant was mistaken in referring his specimens to *C. flexuosus*, as they are more closely allied to *Dicranodontium longirostre*, var. *alpinus*.

**DICRANUM FLAGILLARE** Hedw.

**DICRANUM SCOPARIUM** (L.) Hedw.

**Dicranum longifolium** Hedw. (*Nuttall*).

DICRANUM FULVUM Hook.

DICRANELA HETEROMALLA (L.) Schimp.

DITRICHUM PALLIDUM Schreb.

CERATODON PURPUREUS (L.) Brid.

## GRIMMIACEÆ.

WEISSIA AMERICANA (P. Beauv.) Lindb.

**Weissia ulophylla** Ehrh. (*Nuttall*).

ORTHOTRICHUM BRAUNII Br. &amp; Sch.

**Orthotrichum Ohicense** S. and L. (*Nuttall*).

## HYPNACEÆ.

THUIDIUM RECOGNITUM (Hedw.) Lindb.

ANOMODON ROSTRATUS (Hedw.) Schimp.

ANOMODON ATTENUATUS (Schreb.) Hedw.

AMBLYSTEGIUM ADNATUM Hedw.

AMBLYSTEGIUM SERPENS (L.) Br. &amp; Sch.

AMBLYSTEGIUM ORTHOCLADON (Beauv.) Aust. *Amblystegium serpens orthocladon* (Beauv.) Aust.

AMBLYSTEGIUM VARIUM (Hedw.) Lindb.

AMBLYSTEGIUM IRRIGUUM (Hook. &amp; Wils.) Br. &amp; Sch.

AMBLYSTEGIUM RIPARIUM (L.) Br. &amp; Sch.

AMBLYSTEGIUM CHRYSOPHYLLUM (Brid.) De Not.

HYPNUM DENTICULATUM Linn.

HYPNUM PALUSTRE ?

HYPNUM MOLLE Dicks.

HYPNUM PROLIFERUM Linn.

HYPNUM RUTABULUM Linn.

HYPNUM RECURVANS Schwaeger.

HYPNUM MICROCARPUM C. Muell.

HYPNUM HIANIS Hedw.

HYPNUM DEMISSUM Wils.

HYPNUM PLUMOSUM Huds. *Brachythecium salebrosum*, Br. & Sch.

## STEREODONTACEÆ.

THELIA HIRTELLA (Hedw.) Sull.

THELIA ASPRELLA (Schim.) Sull.

HYLOCOMIUM PARIETINUM (L.) Lindb.

**Hylocomium rugosum** (L.) De Not.Greenbrier Co., Kate's Mountain, alt. 3,300 ft., on ground  
(*Small & Vail*).

HYLOCOMIUM TRIQUETRUM (L.) Br. & Sch.

CAMPYLUM HISPIDULUM (Brid.) Mitt.

CAMPYLUM CHRYSOPHYLLUM (Brid.) E. G. Britton.

Greenbrier Co., Kate's Mountain, alt. 3,300 ft., on ground  
(*Small & Vail*).

CAMPYLUM CHRYSOPHYLLUM TENELLUM L. & J.

PTILIUM CRISTA-CASTRENSE (L.) De Not.

STEREODON IMPONENS (Hedw.) Brid.

STEREODON CUPRESSIFORME (L.) Brid.

STEREODON CURVIFOLIUS (Hedw.) Brid.

PYLAISIA VELUTINA Br. & Sch.

PLAGIOTHECIUM DENTICULATUM Br. & Sch.

PLAGIOTHECIUM DENTICULATUM DENSUM Br. & Sch.

PLAGIOTHECIUM SULLIVANTIÆ Schimp.

CYLINDROTHECIUM SEDUCTRIX (Hedw.) Sull.

CYLINDROTHECIUM CLADORHIZANS (Hedw.) Schimp.

ENTODON PALATINUS (Neck.) Lindb.

#### NECKERACEÆ.

NECKERA PENNATA (L.) Hedw. (*Nuttall*).

CLIMACIUM AMERICANUM Brid.

LEUCODON JULACEUS (Hedw.) Sull.

LEUCODON BRACHYPUS Brid.

HEDWEGIA CILIATA Ehrh. (*Nuttall*)

### Sphagna.

#### SPHAGNACEÆ.

SPHAGNUM CYMBIFOLIUM Ehrh.

## Petríodophyta.

### OPHIOGLOSSACEÆ.

- OPHIOGLOSSUM VULGATUM Linn.  
BOTRYCHIUM TERNATUM (Thunb.) Sw.  
BOTRYCHIUM TERNATUM AUSTRALE (R. Br.) Eaton.  
BOTRYCHIUM TERNATUM RUTÆFOLIUM Man.  
BOTRYCHIUM TERNATUM OBLIQUUM (Muhl.) Milde.  
BOTRYCHIUM TERNATUM DISSECTUM (Spreng.) Milde.  
BOTRYCHIUM VIRGINIANUM (Linn.) Sw.

### FILICES.

- POLYPODIUM VULGARE Linn.  
POLYPODIUM VULGARE *forma* BISSEKATA Millsp. Fl. W. Va., 479 (1892).  
“A form with regularly doubly-serrate thinnish fronds. On mossy boulders along the Blackwater fork of Cheat river.”  
POLYPODIUM POLYPODIOIDES (Linn.) Hitch. *Polypodium incanum* Sw.  
CHEILANTHES LANOSA (Michx.) Watt. *C. vestita* (Spreng.) Sw.  
CHEILANTHES GRACILIS (Fee.) Met. *C. lanuginosa*, Nutt.  
PELLÆA ATROPURPUREA (Linn.) Link.  
PTERIS AQUILINA Linn.  
ADIANTUM PEDATUM Linn.  
ASPLENIUM PINNATIFIDUM Nutt.  
ASPLENIUM TRICHOMANES Linn.  
ASPLENIUM PLATYNEURON (Linn.) Oakes.  
ASPLENIUM MONTANUM Willd.  
ASPLENIUM ANGUSTIFOLIUM Michx.  
ASPLENIUM ACROSTICHOIDES Sw.  
ASPLENIUM FILIX-FÆMINA (Linn.) Bernh.  
CAMPTOSORUS RHIZOPHYLLUS (Linn.) Link.  
PHEGOPTERIS PHEGOPTERIS (Linn.) Underw.  
PHEGOPTERIS HEXAGONOPTERA (Michx.) Fee.  
PHEGOPTERIS DRYOPTERIS (Linn.) Fee.  
DRYOPTERIS THELYPTERIS (Linn.) A. Gray. *Aspidium Thelypteris* (L.) Sw.  
DRYOPTERIS NOVEBORACENSE (Linn.) A. Gray. *Aspidium Noveboracense* (L.) Sw.  
DRYOPTERIS FRAGRANS (Linn.) Schott. *Aspidium fragrans* (L.) Sw.

- DRYOPTERIS SPINULOSA (Retz.) O. K. *Aspidium spinulosum* (Retz.) Sw.  
 DRYOPTERIS SPINULOSA INTERMEDIA (Muhl.) Underw. *Aspidium spinulosum* var. *intermedium* Eaton.  
 DRYOPTERIS SPINULOSA DILATATA (Hoffm.) Underw. *Aspidium spinulosum* var. *dilatatum* Hook.  
 DRYOPTERIS SIMULATA Davenport, Bot. Gaz. 29, 495 (1894).  
*Aspidium spinulosum* var. Comparing my specimens from the deep, wet woods of Randolph Co. with specimens of *D. simulata* recently sent me by Professor Davenport, I find them identical. In publishing the species Professor Davenport includes it under Dryopteris as well as Aspidium, Nephrodium and Lastrea.  
 DRYOPTERIS CRISTATA (Linn.) A. Gray. *Aspidium cristatum* (L.) Sw.  
 DRYOPTERIS GOLDIEANA (Hook) A. Gray. *Aspidium Goldieanum* Hook.  
 DRYOPTERIS FILIX-MAS (Linn.) Schott. *Aspidium Filix-mas* Sw.  
 DRYOPTERIS MARGINALIS (Linn.) A. Gray. *Aspidium marginale* Sw.  
 DRYOPTERIS ACROSTICHOIDES (Michx.) O. K. *Aspidium achrostichoides* Sw.  
 DRYOPTERIS ACROSTICHOIDES SCHWEINITZII (Beck). *Aspidium achrostichoides Schweinitzii* (Beck.) B. S. P.  
 CYSTOPTERIS BULBIFERA (Linn.) Bernh.  
 CYSTOPTERIS FRAGILIS (Linn.) Bernh.  
 ONOCLEA SENSIBILIS Linn.  
 WOODSIA OBTUSA (Spreng.) Torr.  
 DICKSONIA PUNCTILOBULA (Michx.) A. Gray.  
 OSMUNDA REGALIS Linn.  
 OSMUNDA CLAYTONIANA Linn.  
 OSMUNDA CINNAMOMEA Linn.

#### EQUISETACEÆ.

- EQUISETUM ARVENSE Linn.  
 EQUISETUM SYLVATICUM Linn.  
 EQUISETUM HYEMALE Linn.  
 EQUISETUM LÆVIGATUM A. Br.

#### LYCOPODIACEÆ.

- LYCOPODIUM LUCIDULUM Michx.  
 LYCOPODIUM ANNOTINUM Linn.  
 LYCOPODIUM OBSCURUM Linn (*Nuttall*).  
 LYCOPODIUM CLAVATUM Linn.  
 LYCOPODIUM COMPLANATUM Linn.

#### SELAGINELLACEÆ.

- SELAGINELLA RUPESTRIS (Linn.) Spring.

## Anthophyta.

### CONIFERACEÆ.

THUJA OCCIDENTALIS Linn.

JUNIPERUS COMMUNIS Linn.

JUNIPERUS VIRGINIANA Linn.

Jefferson Co., plentiful near Harper's Ferry, trunks 7 to 9 inches in diameter.

TAXUS MINOR (Michx.) Britton. *Taxus baccata* var. *Canadensis* A. Gray.

PINUS STROBUS Linn.

PINUS TÆDA Linn.

PINUS RIGIDA Mill.

PINUS PUNGENS Michx. f.

PINUS VIRGINIANA Mill.

Greenbrier Co., slopes of Kate's Mountain, alt. 3,300 ft. (*Small & Vail*).

PINUS ECHINATA Mill.

PICEA MARIANA (Mill.) B. S. P.

TSUGA CANADENSIS (Linn.) Carr.

ABIES BALSAMEA (Linn.) Miller.

### TYPHACEÆ.

TYPHA LATIFOLIA Linn.

### SPARGANIACEÆ.

SPARGANIUM EURYCARPUM Engelm.

### NAIDACEÆ.

POTAMOGETON LONCHITES Tuck. *P. fluitans* Roth.

### ALISMACEÆ.

ALISMA PLANTAGO-AQUATICA Linn.

SAGITTARIA LATIFOLIA Willd. *S. sagittifolia* Linn. forma *hastata* et *angustifolia*.

SAGITTARIA GRAMINEA Michx.

### HYDROCHARITACEÆ.

UDORA CANADENSIS (Michx.) Nutt. *Elodea Canadensis* Michx.



GRAMINACEÆ.

SPARTINA CYNOSUROIDES (Linn.) Willd.

PASPALUM SETACEUM Michx. *P. setaceum ciliatifolium* (Michx.) is included.

PASPALUM LÆVE Michx.

PANICUM SANGUINALE Linn.

PANICUM SANGUINALE *forma* DEPAUPERATA, Vasey.

This form from dry sterile fields along Falling Run, Monongalia Co., was given the prominence of a form name by Dr. Vasey (1892).

PANICUM PROLIFERUM Lam. *P. proliferum geniculatum* (Eu.) Vasey, included.

PANICUM CAPILLARE Linn.

PANICUM CAPILLARE AGRESTRE Gatt.

PANICUM ANCEPS Michx.

PANICUM AGROSTIDIFORME Lam. *P. agrostioides* Trin.

PANICUM VIRGATUM Linn.

PANICUM WALTERI Poir. *P. latifolium* Linn.

PANICUM CLANDESTINUM Linn.

PANICUM DEPAUPERATUM Muhl.

PANICUM DICHOTOMUM Linn.

*forma* COMMUNE Man.

*forma* FASCICULATUM Man.

*forma* GRACILE Man.

PANICUM DICHOTOMUM ELATUM.

PANICUM PUBESCENS Lam.

PANICUM CRUS-GALLI Linn.

PANICUM CRUS-GALLI HISPIDUM (Muhl.) Torr.

CHAMÆRAPHIS GLAUCA (Linn.) O. K.

CHAMÆRAPHIS VIRIDIS (Linn.) Millsp. Fl. W. Va. (1892).

CHAMÆRAPHIS ITALICA (Linn.) O. K.

CENCHRUS TRIBULOIDES Linn.

HOMALOCENCHRUS VIRGINICUS (Willd.) Britton.

HOMALOCENCHRUS ORYZOIDES (Linn.) Poll.

TRIPSACUM DACTYLOIDES Linn.

ANDROPOGON PROVINCIALIS Lam.

ANDROPOGON SCOPARIUS Michx.

ANDROPOGON VIRGINICUS Linn.

ANDROPOGON NUTANS LINNÆNUS Hack.

PHALARIS ARUNDINACEA Linn.

ARISTIDA DICHOTOMA Linn.

ARISTIDA OLIGANTHA Michx.

ARISTIDA LANATA Poir.

MUHLENBERGIA SOBOLIFERA (Muhl.) Trin.

MUHLENBERGIA MEXICANA (Linn.) Trin.

MUHLENBERGIA SYLVATICA (Torr.) A. Gray.

MUHLENBERGIA DIFFUSA Schreb.

BRACHYELYTRUM ERECTUM (Schreb.) Beauv. *Brachyelytrum aristosum* (Michx.) B. S. P.

BRACHYELYTRUM ERECTUM GLABRATUM (Vasey.)

*Brachyelytrum aristosum glabratum* Vasey, mss. In Millsp. Flora W. Va., 469 (1892).

A new variety discovered by Mr. L. W. Nuttall. It agrees with the species except that it is perfectly smooth, and has an awn pointed second glume which is about one-half the length of the flowering glume. Its most striking peculiarity is that it has invariably two culms from each rootstock.

High, rocky woods, rare.

PHLEUM PRATENSE Linn.

AGROSTIS ALBA Linn.

AGROSTIS ALBA VULGARIS (With.) Thurb.

*Forma* ARISTATA Millsp. in Fl. W. Va., 469 (1892).

AGROSTIS PERENNANS (Walt.) Tuck.

AGROSTIS HIEMALIS (Walt.) B. S. P.

**Calamagrostis cinnoides** (Muhl.) Scribn. (*Nuttall*).

CINNA ARUNDINACEA Linn.

ARRHENATHERUM ELATIUS (Linn.) Beauv.

HOLCUS LANATUS Linn.

DANTHONIA SPICATA (Linn.) Beauv.

DANTHONIA COMPRESSA Austin.

ELEUSINE INDICA (Linn.) Gærtn.

DACTYLOCTENIUM ÆGYPTICUM (Linn.) Willd. *Eleusine Ægyptica* Pers.

SIEGLINGIA SESLERIOIDES (Michx.) Scribn. *Sieglingia cuprea* (Michx.)

EATONIA OBTUSATA (Michx.) A. Gray.

EATONIA PENNSYLVANICA (DC.) A. Gray.

ERAGROSTIS HYPNOIDES (Lam.) B. S. P.

ERAGROSTIS ERAGROSTIS (Linn.) Karst. *Eragrostis minor* Host,

ERAGROSTIS MAJOR Host.

ERAGROSTIS PILOSA (Linn.) Beauv.

ERAGROSTIS CAROLINIANA (Spreng.) Scribn. *Eragrostis Purshii* Schrad.

ERAGROSTIS CAPILLARIS (Linn.) Nees.

ERAGROSTIS FRANKII Meyer.

MELICA MUTICA Walt.

KORYCARPUS DIANDRUS (Michx.) O. K. *Corycarpus Americanus* (Beauv.)

UNIOLA LATIFOLIA Michx.

DACTYLIS GLOMERATA Linn.

POA ANNUA Linn.

POA COMPRESSA Linn.

*Forma* DEPAUPERATA Fl. W. Va., 472 (1892).

POA PRATENSIS Linn.

POA TRIVALIS Linn.

POA SYLVESTRIS A. Gray.

POA ALSODES A. Gray.

POA AUTUMNALIS Muhl. *Poa flexuosa* Muhl. not Wahl.

POA BREVIFOLIA Muhl.

PANICULARIA CANADENSIS (Michx.) O. K.

PANICULARIA ELONGATA (Torr.) O. K.

PANICULARIA NERVATA (Willd.) O. K.

*Forma* MAJOR Fl. W. Va., 473 (1892).

FESTUCA OCTOFLORA Walt.

FESTUCA ELATIOR Linn.

FESTUCA ELATIOR PRATENSIS (Huds.) Hack. *Festuca pratensis* Huds.

FESTUCA NUTANS Willd.

BROMUS HORDEACEUS Linn. *Bromus mollis* Linn.

BROMUS SECALINUS Linn.

BROMUS RACEMOSUS Linn.

BROMUS CILIATUS Linn.

LOLIUM PERENNE Linn.

ELYMUS VIRGINICUS Linn.

ELYMUS CANADENSIS GLAUCIFOLIUS (Willd.) Torr.

ELYMUS STRIATUS Willd.

ELYMUS STRIATUS VILLOSUS (Muhl.) A. Gray.

HYSTRIX HYSTRIX (Linn.) Millsp. Fl. W. Va., 474 (1892).

#### CYPERACEÆ.

CYPERUS FLAVESCENS Linn.

CYPERUS DIANDRUS Torr.

CYPERUS ESCULENTUS Linn.

CYPERUS STRIGOSUS Linn.

CYPERUS REFRACTUS Engelm.

CYPERUS LANCASTRIENSIS Porter.

KYLLINGA PUMILA Michx.

DULICHIMUM ARUNDINACEUM (Linn.) Britton. *Dulichium spathaceum* Pers.

ELEOCHARIS TUBERCULOSA (Michx.) R. & S.

ELEOCHARIS OVATA (Roth.) R. & S.

ELEOCHARIS PALUSTRIS (Linn.) R. & S.

ELEOCHARIS PALUSTRIS GLAUDESCENS (Willd.) A. Gray.

ELEOCHARIS TENUIS (Willd.) Schult.

ELEOCHARIS ACICULARIS (Linn.) R. & S.

FIMBRISTYLIS AUTUMNALIS (Linn.) R. & S.

- SCIRPUS AMERICANUS Pers. *Scirpus pungens* Vahl.  
 SCIRPUS LACUSTRIS Linn.  
 SCIRPUS SYLVATICUS Linn.  
 SCIRPUS ATROVIRENS Muhl.  
 SCIRPUS POLYPHYLLUS Vahl.  
 SCIRPUS LINEATUS Michx. *Eriophorum lineatum* (Michx.) Benth & Hook.  
 SCIRPUS CYPERINUS (Linn.) Kunth. *Eriophorum cyperinum* Linn.  
 ERIOPHORUM VIRGINICUM Linn.  
 RYNCHOSPORA GLOMERATA (Linn.) Vahl.  
 CAREX FOLLICULATA Linn.  
 CAREX INTUMESCENS Rudge.  
 CAREX ASA-GRAYI Bailey. *Carex Grayii* Carey.  
 CAREX LUPULINA Muhl.  
 CAREX LURIDA Vahl.  
 CAREX LURIDA GRACILIS (Boott.) Bailey.  
 CAREX FRANKII (Kunth.) *Carex stenolepis* Torr.  
 CAREX SQUARROSA Dewey.  
 CAREX STRICTA Lam.  
 CAREX TORTA Boott.  
 CAREX PRASINA Vahl.  
 CAREX CRINITA Lam.  
 CAREX CRINITA × PRASINA Bailey, Flor. W. Va., 460 (1892).  
 CAREX VIRESCENS Muhl.  
 CAREX VIRESCENS COSTATA (Schw.) Dewey.  
 CAREX TRICEPS HIRSUTA (Willd.) Bailey.  
 CAREX DEBILIS RUDGEI Bailey.  
 CAREX VENUSTA MINOR Bœckl.  
 CAREX GRACILLIMA Schw.  
 CAREX AMPHIBOLA Steud. *Carex grisea angustifolia* Boott.  
 CAREX GLANCODEA Tuck.  
 CAREX LAXIFLORA Lam.  
 CAREX LAXIFLORA PATULIFOLIA (Dewey.) Carey.  
 CAREX ALBURSINA Sheldon. *Carex laxiflora latifolia* Boott.  
 CAREX DIGITALIS Willd.  
 CAREX LAXICULMIS Schw.  
**Carex platyphylla** Carey (*Nuttali*).  
 CAREX PLANTAGINEA Lam. (*Nuttall*).  
**Carex varia** Muhl. (*Nuttall*).  
 CAREX VARIA COLORATA Bailèy.  
 CAREX PENNSYLVANICA Lam.  
 CAREX PEDICILLATA (Dewey.) Britton. *Carex communis* Bailey.  
 CAREX JAMESII Schw.  
 CAREX LEPTALEA Vahl. *Carex polytrichioides* Muhl.

CAREX FRASERI And.

The following remarks of Prof. T. C. Porter render it evident that this rare and odd sedge came originally from Randolph or Barbour County, each of which lies between the headwaters of the two Kanawhas:

“Muhlenberg, in his *Descriptio uberior Graminum*, etc., p. 265, under *C. lagopus?* which is *C. Fraseri*, Andrews adds these words: ‘Habitat in Tyger-Valley, Pennsylvania, unde siccam habeo et vivam.’ Kin, the German gardener who collected in Pennsylvania brought it home, and his label reads thus: ‘Deigher Walli in der Wilternus.’ Dr. Gray has shrewdly conjectured that by ‘Deigher Walli,’ or Tyger Valley, is meant Tygart’s Valley, which lies further south in Virginia.” In a foot-note, Professor Porter adds, “A box containing the Carices of Muhlenberg has just been discovered (1877) in the herbarium of the Academy, Philadelphia, and the label attached to the specimens of Kin’s collection places Tyger Valley ‘prope amnem Kenahway.’”

Mr. Nuttall has found a plentiful station for this species near Nuttallburg in the Great Kanawha region.

CAREX STIPATA Muhl.

CAREX VULPINOIDEA Michx.

CAREX ROSEA Schk.

CAREX ROSEA RADIATA Dewey.

CAREX RETROFLEXA Muhl. *Carex rosea retroflexa* Torr.

CAREX SPARGANIOIDES Muhl.

CAREX MUHLENBERGII var. Bailey.

CAREX MUHLENBERGII XALAPENSIS (Kth.) Britton. *Carex Muhlenbergii enervis* Boott.

CAREX CEPHALOPHORA Muhl.

CAREX CANESCENS VULGARIS Bailey.

CAREX TRIBULOIDES Wahl.

CAREX TRIBULOIDES TURBATA Bailey.

CAREX SCOPARIA Schk.

CAREX STRAMINEA Willd.

ARACEÆ.

ARISÆMA TRIPHYLLUM (Linn.) Torr.

ARISÆMA DRACONTIUM (Linn.) Schott.

SPATHYEMA FÆTIDA (Linn.) Raf.

ACORUS CALAMUS Linn. (*Nuttall*).

LEMNACEÆ.

LEMNA MINOR Linn.

COMMELINACEÆ.

COMMELINA VIRGINICA Linn.

TRADESCANTIA VIRGINIANA Linn.

TRADESCANTIA PILOSA Lehm.

## JUNCACEÆ.

- JUNCUS EFFUSUS Linn.  
 JUNCUS SETACEUS Rostk.  
 JUNCUS MARGINATUS Rostk.  
 JUNCUS MARGINATUS PAUCICAPITATUS Engelm.  
 JUNCUS TENUIS Willd.  
 JUNCUS ACUMINATUS Michx.  
 JUNCUS NODOSUS Linn.  
 JUNCUS CANADENSIS J. Gay.  
 JUNCOIDES PILOSUM (Linn.) O. K. *Luzula pilosa* Willd.  
 JUNCOIDES CAMPESTRE (Linn.) O. K. *Luzula cumpestris* DC.

## LILIACEÆ.

- ASPARAGUS OFFICINALIS Linn.  
 POLYGONATUM BIFLORUM (Walt.) Ell.  
 POLYGONATUM BIFLORUM COMMUTATUM (R. & S.) Morong. *Polygonatum commutatum* Dietr.  
 STREPTOPUS ROSEUS Michx.  
 DISPORUM LANUGINOSUM (Michx.) Britton.  
 VAGNERA RACEMOSA (Linn.) Morong. *Unifolium racemosum* Britton.  
 UNIFOLIUM CANADENSE (Desf.) Greene. *Maianthemum Canadense* Desf.  
 HEMEROCALLIS FULVA Linn.  
 ALLIUM VINEALE Linn.  
 ALLIUM TRICOCCUM Aiton.  
 ALLIUM CERNUUM Roth.  
 ALLIUM CANADENSE Kalm.  
 CAMASSIA FRASERI (A. Gray) Torr.  
 MUSCARI BOTRYOIDES (Linn.) Mill.  
 ORINTHOGALLUM UMBELLATUM Linn.  
 LILIUM PHILADELPHICUM Linn.  
 LILIUM SUPURBUM Linn. (*Nuttall*).  
 LILIUM CANADENSE Linn.  
 LILIUM TIGRINUM Andr.  
 ERÛTHRONIUM AMERICANUM Ker.  
**Erythronium albidum** Nutt.  
 Rich soil, Kanawha region, along Coal and Len's Creeks, Ohio Co., near Mt. de Chantal (*Rev. A. Boutlou*).  
 CHAMÆLIRIUM LEUTEUM (Linn.) A. Gray.  
 UVULARIA PERFOLIATA Linn.  
**Uvularia puberula** Michx.  
 Kate's Mountain, alt. 3,300 ft. (*Small & Vail*).  
 UVULARIA SESSILIFOLIA Linn.  
 CLINTONIA BOREALIS (Ait.) Raf.  
 CLINTONIA UMBELLULATA (Michx.) Torr.

- MEDIOLA VIRGINIANA Linn.  
 TRILLIUM SESSILE Linn.  
 TRILLIUM ERECTUM Linn.  
 TRILLIUM ERECTUM ALBUM Pursh.  
 TRILLIUM ERECTUM DECLINATUM Pursh.  
 TRILLIUM GRANDIFLORUM (Michx.) Salisb.  
 TRILLIUM CERNUUM Linn.  
 TRILLIUM NIVALE Riddell.  
 TRILLIUM UNDULATUM Willd. *Trillium erythrocarpum* Michx.  
 ALETRIS FARINOSA Linn.  
 MELANTHIUM VIRGINICUM Linn.  
 MELANTHIUM PARVIFLORUM (Michx.) S. Wats.  
 VERATRUM VIRIDE Ait. (*Nuttall*).  
 CHROSPHERMA MUSCÆTOXICUM (Walt.) O. K.

SMILACEÆ.

- SMILAX HERBACEA Linn.  
 SMILAX ROTUNDIFOLIA Linn.  
 SMILAX GLAUCA Walt.  
 SMILAX PSEUDO-CHINA Linn.  
 SMILAX HISPIDA Muhl. (*Nuttall*).

AMARYLLIDACEÆ.

- HYPOXIS HIRSUTA (Linn.) Coville. *Hypoxis erecta* Linn.

DIOSCOREACEÆ.

- DIOSCOREA VILLOSA Linn.

IRIDACEÆ.

- IRIS VERSICOLOR Linn.  
 IRIS VERNA Linn.  
 IRIS CRISTATA Ait.  
 SISYRINCHIUM BERMUDIANA Linn. *S. angustifolium* Mill. *S. anceps* Cav.

ORCHIDACEÆ.

- ACHROANTHES UNIFOLIA (Michx.) Raf. *Malaxis unifolia* Michx.  
 LEPTORCHIS LILIFOLIA (Linn.) O. K. (*Nuttall*).  
 APLECTRUM SPICATUM (Walt.) B. S. P.  
 CORALLORHIZA CORALLORHIZA (Linn.) Karst. *C. innata* R. Br.  
 CORALLORHIZA ODONTORHIZA (Willd.) Nutt.  
 CORALLORHIZA MULTIFLORA Nutt.  
 GYROSTACHYS CERNUA (Linn.) O. K.  
 GYROSTACHYS GRACILIS (Biegl.) O. K.  
 PERAMIUM REPENS (Linn.) Salisb. *Goodyera repens* R. Br.

- PERAMIUM PUBESCENS (Willd.) C. C. Curtis. *Goodyera pubescens* R. Br.  
 LIMODORUM TUBEROSUM Linn. *Calopogon tuberosus* (Linn.) B. S. P.  
 POGONIA OPHIOGLOSSOIDES (Linn.) Ker. *Pogonia trianthophora* (Sw.)  
 B. S. P. (*Nuttall*).  
 ORCHIS SPECTABILIS Linn.  
 HABENARIA CLAVELLATA (Michx.) Spreng. *H. tridentata* Hook.  
 HABENARIA FLAVA (Linn.) A. Gray.  
 HABENARIA ORBICULATA (Pursh) Torr.  
 HABENARIA CILIARIS (Linn.) R. Br.  
 HABENARIA LACERA (Michx.) R. Br. *Habenaria peramæna* A. Gray.  
 (*Nuttall*).  
 HABENARIA PSYCODES (Linn.) A. Gray.  
 CYPRIPEDIUM PARVIFLORUM Salisb.  
 CYPRIPEDIUM HIRSUTUM Mill. *C. pubescens* Willd.  
 Slope of Kate's Mountain, alt. 3,300 ft. (*Small & Vail*).  
 CYPRIPEDIUM ACAULE Ait.

## SAURURACEÆ.

- SAURURUS CERNUUS Linn.

## JUGLANDACEÆ.

- JUGLANS CINEREA Linn.  
 JUGLANS NIGRA Linn.  
 HICORIA OVATA (Mill.) Britton.  
 HICORIA SULCATA (Willd.) Britton.  
 HICORIA ALBA (Linn.) Britton.  
 HICORIA GLABRA (Mill.) Britton.  
 HICORIA MICROCARPA (Nutt.) Britton.  
 HICORIA MINIMA (Marsh.) Britton. (*Nuttall*.)

## SALICACEÆ.

- SALIX NIGRA Marsh.  
 SALIX NIGRA FALCATA Torr.  
 SALIX AMYGDALOIDES And.  
 SALIX ALBA VITELLINA Koch.  
 SALIX BABYLONICA Tourn.  
 SALIX DISCOLOR Muhl.  
 SALIX HUMILIS Marsh.  
 SALIX SERICEA Marsh.  
 SALIX CORDATA Muhl.  
 POPULUS ALBA Linn.  
 POPULUS TREMULOIDES Michx.  
 POPULUS GRANDIDENTATA Michx.  
 POPULUS BALSAMIFERA CANDICANS (Ait.) A. Gray.  
 POPULUS MONILIFERA Ait.



BETULACEÆ.

BETULA LENTA Linn.

**Betula lenta** × **lutea**.

A number of specimens of this hybrid birch were found in 1892 at Pickens, Randolph Co. In all, the bark forms of each species were intermixed. All the specimens were small saplings about 2 in. in diameter (*Millsbaugh*).

BETULA LUTEA Michx. f. (*Nuttall*).

BETULA POPULIFOLIA Marsh.

BETULA NIGRA Linn.

ALNUS VIRIDIS (Chaix.) DC.

ALNUS RUGOSA (Ehrh.) Koch.

CARPINUS CAROLINIANA Walt.

OSTRYA VIRGINIANA (Mill.) Willd.

CORYLUS AMERICANA Walt. (*Nuttall*).

CORYLUS ROSTRATA Ait.

FAGACEÆ.

QUERCUS ALBA Linn.

QUERCUS MINOR (Marsh). Sarg.

QUERCUS MACROCARPA Michx.

QUERCUS PRINUS Linn.

QUERCUS MUHLENBERGII Engelm.

QUERCUS RUBRA Linn.

QUERCUS COCCINEA Wang.

QUERCUS VELUTINA Lam. *Q. tinctoria* Bartr.

QUERCUS PALUSTRIS DuRoi. (*Nuttall*).

QUERCUS DIGITATA (Marsh) Sudw. *Q. cuneata* Wang.

QUERCUS NIGRA Linn.

QUERCUS ILICIFOLIA Wang.

QUERCUS IMBRICARIA Michx.

CASTANEA DENTATA (Marsh.) Sudw. *C. sativa Americana* Wats. & Coult.

CASTANEA PUMILA Mill.

FAGUS ATROPUNICEA (Marsh.) Sudw. *F. ferruginea* Ait.

ULMACEÆ.

ULMUS PUBESCENS Walt. *U. fulva* Michx.

ULMUS AMERICANA Linn.

ULMUS RACEMOSA Thomas.

CELTIS OCCIDENTALIS Linn.

MORACEÆ.

CANNABIS SATIVA Linn.

HUMULUS LUPULUS Linn.

MORUS RUBRA Linn.

MORUS ALBA Linn.

- Toxylon pomiferum** Raf. *Maclura aurantiaca* Nutt.  
Banks of the Ohio River, Mason Co., near Point Pleasant.  
(*Millspaugh*).  
PAPYRIUS PAPYRIFERA (Vent.) O. K. *Broussonetia papyrifera* Vent.

## URTICACEÆ.

- URTICA GRACILIS Ait.  
URTICA URENS Linn.  
URTICASTRUM DIVARICATUM (Linn.) O. K. *Laportea Canadensis* Gaud.  
ADICEA PUMILA (Linn.) Raf. *Pilea pumila* A. Gray.  
BOEHMERIA CYLINDRICA (Linn.) Willd.

## LORANTHACEÆ.

- PHORADENDRON FLAVESCENS (Pursh) Nutt.

## SANTALACEÆ.

- PYRULARIA PUBERA Michx.  
Also found in Nicholas Co., along Peter Creek. (*Rev. A. Boutlou.*)

## ARISTOLOCHIACEÆ.

- ASARUM CANADENSE Linn.  
ASARUM VIRGINICUM Linn.  
ARISTOLOCHIA SERPENTARIA Linn.  
ARISTOLOCHIA MACROPHYLLA Lam. *A. Siphon* L'Her.

## POLYGONACEÆ.

- ERIGONUM ALLENII Watson, in A. Gray, Man. Ed. 6, 734 (1890).  
Perennial, white tomentose throughout, the tall scape-like stem repeatedly dichotomous above; radical leaves lanceolate, long-petiolate, the upper in whorls of 4 or 5, ovate to ovate-oblong, very shortly petiolate, much reduced above; involucre mostly sessile; flowers glabrous, yellow, the segments elliptical.  
Near White Sulphur Springs, Greenbrier Co. (*Dr. T. F. Allen*).  
POLYGONUM ORIENTALE Linn.  
POLYGONUM PENNSYLVANICUM Linn.  
POLYGONUM PERSICARIA Linn.  
POLYGONUM PERSICARIA *forma* ALBIFLORA Millsp. Fl. W. Va. 332 (1892).  
POLYGONUM HYDROPIPER Linn.  
POLYGONUM PUNCTATUM Ell. *P. acre* H. B. K.  
POLYGONUM HYDROPIPEROIDES Michx.  
POLYGONUM VIRGINIANUM Linn.  
POLYGONUM AVICULARE Linn.  
POLYGONUM ERECTUM Linn.  
POLYGONUM TENUE Michx.

- POLYGONUM SAGITTATUM Linn.  
 POLYGONUM ARIFOLIUM Linn.  
 POLYGONUM CONVULVULUS Linn.  
 POLYGONUM SCANDENS Linn. *P. dumetorum scandens* A. Gray.  
 FAGOPYRUM FAGOPYRUM (Linn.) Karst. *F. esculentum* Mœnch.  
 RUMEX BRITANICA Linn.  
 RUMEX CRISPUS Linn.  
 RUMEX OBTUSIFOLIUS Linn.  
 RUMEX CRISPUS × OBTUSIFOLIUS (Meisn.)  
 RUMEX SANGUINEUS Linn.  
 RUMEX CONGLOMERATUS Murr.  
 RUMEX ACETOSELLA Linn.

#### CHENOPODIACEÆ.

- CHENOPODIUM ALBUM Linn.  
 CHENOPODIUM ALBUM VIRIDE (Linn.) Moq.  
 CHENOPODIUM HYBRIDUM Linn.  
 CHENOPODIUM URBICUM Linn.  
 CHENOPODIUM GLAUCUM Linn.  
 CHENOPODIUM BOTRYS Linn.  
 CHENOPODIUM AMBROSIODES Linn.  
 CHENOPODIUM ANTHELMINTICUM Linn. *C. ambrosioides anthelminticum*  
 A. Gray.

#### AMARANTHACEÆ.

- AMARANTHUS HYBRIDUS Linn. *A. hypochondriacus* Linn. *A. chlorostachys* Willd. (*Nuttall*).  
 AMARANTHUS HYBRIDUS PANICULATUS (Linn.) Uline & Bray. *A. paniculatus* Linn.  
 AMARANTHUS RETROFLEXUS Linn. (*Nuttall*).  
 AMARANTHUS ALBUS Linn.  
 AMARANTHUS SPINOSUS Linn.

#### PHYTOLACCACEÆ.

- PHYTOLACCA DECANDRA Linn.

#### AIZOACEÆ.

- MOLLUGO VERTICILLATA Linn.

#### PORTULACACEÆ.

- PORTULACA OLERACEA Linn.  
 CLAYTONIA VIRGINICA Linn.  
 CLAYTONIA CAROLINIANA Michx.

## CARYOPHYLLACEÆ.

DIANTHUS ARMERIA Linn.

SAPONARIA OFFICINALIS Linn.

SILENE STELLATA (Linn.) Ait.

SILENE VIRGINICA (Linn.)

SILENE ALBA Muhl. *S. nivea* Otth.SILENE CAROLINIANA Walt. *S. Pennsylvanica* Michx.Greenbrier Co., on Kate's Mountain, alt. 3,300 ft. (*Small & Vail*).

SILENE ANTERRHINA Linn.

AGROSTEMMA GITHAGO Linn. *Lychnis Githago* Lam.

CERASTIUM VULGATUM Linn.

CERASTIUM VISCOSUM Linn.

CERASTIUM ARVENSE Linn.

CERASTIUM LONGIPEDUNCULATUM Muhl. *C. nutans* Raf.ALSINE MEDIA Linn. *Stellaria media* Smith.ALSINE PUBERA (Michx.) Britton. *Stellaria pubera* Michx.ALSINE LONGIFOLIA (Muhl.) Britton. *Stellaria longifolia* Muhl.

ARENARIA SERPYLLIFOLIA Linn.

SPERGULA ARVENSIS Linn. (*Nuttall*).

ANYCHIA CANADENSIS (Linn.) B. S. P.

**Anychia dichotoma** Michx.

This species first appeared in this locality in 1895, at the bottom of a newly excavated railroad cut. Had the seeds been buried and dormant? (*Nuttall*, 1837).

PARONYCHIA DICHOTOMA (Linn.) Nutt.

## NYMPHACEÆ.

CASTALIA ODORATA (Dryand) Woodv. &amp; Wood.

NYMPHÆA ADVENA Soland.

## MAGNOLIACEÆ.

MAGNOLIA ACUMINATA Linn.

MAGNOLIA TRIPETALA Linn.

MAGNOLIA FRASERI Walt., Fl. Car., 159 (1788).

Leaves oblong-obovate or spatulate, auriculate at the base, glabrous, 8 to 20 in. long; petals obovate-spatulate, with narrow claws, 4 in. long. Flower more graceful and cone of fruit smaller than in the preceding species.

Type locality doubtless in the Alleghanies of West Virginia, where it is abundant and in typical form.

LIRIODENDRON TULIPIFERA Linn.

## ANONACEÆ.

ASIMINA TRILOBA (Linn.) Dunal.

RANUNCULACEÆ.

CLEMATIS VIRGINIANA Linn.

CLEMATIS VIORNA Linn.

**Clematis ovata** Pursh.

Greenbrier Co., in rocky soil high up on Kate's Mountain, near White Sulphur Springs (*N. L. Britton*).

ASTRAGENE AMERICANA Sims. *Clematis verticillaris* DC.

ANEMONE CYLINDRICA A. Gray.

ANEMONE VIRGINIANA Linn.

ANEMONE CANADENSIS Linn.

ANEMONE QUINQUEFOLIA Linn.

ANEMONE TRIFOLIA Linn.

Greenbrier Co., in a dry ravine at White Sulphur Springs, alt. about 2,300 ft. (*A. A. Heller*).

HEPATICACHE HEPATICACHE (Linn.) Karst.

HEPATICACHE ACUTA (Pursh.) Britton.

SYNDESMON THALICTROIDES (Linn.) Hoffmg.

SYNDESMON THALICTROIDES *forma* ROSEA Millsp. Fl. W. Va., 319 (1892).

THALICTRUM DIOICUM Linn.

THALICTRUM POLYGAMUM Muhl.

THALICTRUM PURPURASCENS Linn.

THALICTRUM CLAVATUM DC.

TRAUTVETTERIA CAROLINENSIS (Walt.) Vail? *T. palmata* F. & M.

RANUNCULUS OBTUSIUSCULUS Raf. *R. ambigenus* Watson.

RANUNCULUS ABORTIVUS Linn.

RANUNCULUS SCELERATUS Linn.

RANUNCULUS RECURVATUS Poir.

RANUNCULUS FASCICULATUS Muhl.

RANUNCULUS SEPTENTRIONALIS Poir.

RANUNCULUS PENNSYLVANICUS Linn f.

RANUNCULUS REPENS Linn.

RANUNCULUS ACRIS Linn.

HELLEBORUS VIRIDIS Linn.

CALTHA PALUSTRIS Linn.

ISOPYRUM TRIFOLIUM (Linn.) Britt.

AQUILEGIA CANADENSIS Linn.

DELPHINIUM TRICORNE Michx. (*Nuttall*).

DELPHINIUM TRICORNE *forma* ALBIFLORA Millsp. Fl. W. Va., 322 (1892).

DELPHINIUM CONSOLIDA Linn.

ACONITUM UNCINATUM Linn.

CIMICIFUGA AMERICANA Michx.

CIMICIFUGA RACEMOSA (Linn.) Nutt.

ACTÆA RUBRA (Ait.) Willd. *A. spicata rubra* Ait.

ACTÆA ALBA (Linn.) Mill.

HYDRASTIS CANADENSIS Linn.

XANTHORRHIZA APIIFOLIA L'Her.

#### BERBERIDACEÆ.

BERBERIS CANADENSIS Mill.

BERBERIS VULGARIS Linn.

CAULOPHYLLUM THALICTROIDES (Linn.) Michx.

JEFFERSONIA DIPHYLLO (Linn.) Pers.

PODOPHYLLUM PELTATUM Linn.

#### MENISPERMACEÆ.

MENISPERMUM CANADENSE Linn.

#### CALYCANTHACEÆ.

CALYCANTHUS FLORIDUS Linn. *Beurera florida* Ehret.

CALYCANTHUS FERTILIS Walt. *Beurera levigatus* (Willd.) O.K.

#### LAURACEÆ.

SASSAFRAS SASSAFRAS (Linn.) Karst. *Sassafras officinale* Nees.

BENZOIN BENZOIN (Linn.) Coult. *Lindera Benzoïn* Meisn.

#### PAPAVERACEÆ.

SANGUINARIA CANADENSIS Linn.

STYLOPHORUM DIPHYLLUM (Michx.) Nutt. (*Nuttall*).

CHELIDONIUM MAJUS Linn.

**Glaucium Glaucium** (L.) Karst.

Jefferson Co., near Charlestown, Shenandoah Junction, Spring Mill, and Milltown (*Millsbaugh*).

PAPAVER DUBIUM Linn.

ADLUMIA FUNGOSA (Ait.) Greene.

BICUCULLA CUCULLARIA (Linn.) Millsp. Fl. W. Va., 327 (1892).

BICUCULLA CANADENSIS (Goldie) Millsp. loc. cit.

BICUCULLA EXIMINA (Ker) Millsp. loc. cit.

CAPNOIDES SEMPERVIRENS (Linn.) Boerk. (*Nuttall*).

CAPNOIDES FLAVULUM (Raf.) O.K.

CAPNOIDES AUREUM (Willd.) O.K.

FUMARIA OFFICINALIS Linn.

#### CRUCIFERACEÆ.

RORIPA NASTURTIUM (Linn.) Rusby. *Nasturtium officinale* R. Br.

RORIPA SYLVESTRIS (Linn.) Bess. *Nasturtium sylvestre* R. Br.

RORIPA OBTUSA (Nutt.) Britton. *Nasturtium obtusum* Nutt.

RORIPA PALUSTRIS (Linn.) Bess. *Nasturtium palustre* DC.

- RORIPA HISPIDA (Desv.) Britton. *Nasturtium hispidum* DC.  
 RORIPA ARMORACIA (Linn.) Hitch. *Nasturtium Armoracia* Fries.  
 BARBAREA BARBAREA (Linn.) MacM. *B. vulgaris* R. Br.  
 BARBAREA PRAECOX (J. E. Smith) R. Br.  
 ARABIS PATENS Sulliv.  
 ARABIS LAEVIGATA (Muhl.) Poir.  
 ARABIS CANADENSIS Linn.  
 ARABIS LYRATA Linn.  
 DENTARIA DIPHYLLO Michx. *Cardamine diphylla* Wood.  
 DENTARIA HETEROPHYLLA Nutt. *Cardamine heterophylla* Wood.  
 DENTARIA LACINIATA Muhl. *Cardamine laciniata* Wood.  
 DENTARIA LACINIATA MULTIFIDA Muhl.  
 CARDAMINE HIRSUTA Linn.  
 CARDAMINE BULBOSA (Schreb.) B. S. P.  
 CARDAMINE DOUGLASSII (Torr.) Britton.  
 CARDAMINE ROTUNDIFOLIA Michx.  
 DRABA RAMOSISSIMA Desv.  
 DRABA VERNA Linn.  
 Greenbrier Co., on Kate's Mountain, alt. 3,300 ft. (*Small & Vail*).  
 HESPERIS MATRONALIS Linn.  
 SISYMBRIUM OFFICINALE (Linn.) Scop.  
 STENOPHRAGMA THALIANA (Linn.) Celak. *Sisymbrium Thalianum* (L. Gay).  
 ERYSIMUM CHEIRANTHOIDES Linn.  
 CAMELINA SATIVA (Linn.) Crantz.  
 BRASSICA NIGRA (Linn.) Koch.  
 BRASSICA SINAPISTRUM Boiss.  
 BURSA BURSA-PASTORIS (Linn.) Weber.  
 LEPIDIUM VIRGINICUM Linn.  
 LEPIDIUM CAMPESTRE (Linn.) R. Br.  
 RAPHANUS SATIVUS Linn.

CAPPARIDACEÆ.

- CLEOME SPINOSA Linn.

DROSERACEÆ.

- DROSELA ROTUNDIFOLIA Linn.

CRASSULACEÆ.

- SEDUM PULCHELLUM Michx.  
 SEDUM NEVII A. Gray.  
 SEDUM TERNATUM (Haw) Michx.  
 SEDUM TELEPHIOIDES Michx.  
 SEDUM TELEPHIUM Linn.  
 PENTHORUM SEDOIDES Linn.

## SAXIFRAGACEÆ.

ASTILBE BITERNATA (Vent.) Britton. *A. decandra* Don.

SAXIFRAGA VIRGINIENSIS Michx.

SAXIFRAGA PENNSYLVANICA Linn.

SAXIFRAGA MICRANTHIDIFOLIA (Haw) B. S. P.

THEROFON ACONITIFOLIUM (Nutt.) Millsp. Fl. W. Va. 361 (1892).

TIARELLA CORDIFOLIA Linn.

MITELLA DIPHYLLA Linn.

HEUCHERA VILLOSA Michx.

HEUCHERA AMERICANA Linn.

HEUCHERA RUGELII Shuttlw.

PARNASSIA GRANDIFOLIA DC.

HYDRANGEA ARBORESCENS Linn.

HYDRANGEA ARBORESCENS KANAWHANA Millsp. Fl. W. Va. 363 (1892).

Low straggling bush, leaves small, paler beneath, acuminate, somewhat cordate at the base; cymes very open and branching, marginal radiant flowers many, 1 in. broad, fertile flowers nearly glabrous, smaller than in the species. Along the Little Kanawha River from Kanawha Station to Glenville. (*Millsbaugh*).

RIBES CYNOSBATI Linn.

RIBES ROTUNDIFOLIUM Michx.

RIBES PROSTRATUM L'Her.

RIBES FLORIDUM L'Her.

## HAMAMELIDACEÆ.

HAMAMELIS VIRGINICA Linn.

LIQUIDAMBAR STYRACIFLUA Linn.

## PLATANACEÆ.

PLATANUS OCCIDENTALIS Linn.

## ROSACEÆ.

PRUNUS AMERICANA Marsh.

Randolph Co., near Pickens (*Millsbaugh*). (*Nuttall*).

PRUNUS ANGUSTIFOLIA Marsh. *P. Chicasa* Michx.

PRUNUS PENNSYLVANICA Linn. f.

PRUNUS VIRGINIANA Linn. (*Nuttall*).

PRUNUS SEROTINA Ehrh.

OPULASTER OPULIFOLIUS (Linn.) O.K.

SPIRÆA CORYMBOSA Raf. *S. betulæfolia* Pall.

SPIRÆA VIRGINIANA Britton, Bull. Torr. Club, 17, 314 (1890).

"A glabrous shrub, the branches forming long wands, erect or reclining, 1 to 4 ft. long. Leaves oblong or slightly oblanceolate, thin, obtuse or short-pointed at the apex, rounded or cuneate at the base, 1.5 to 2 in. long, 5 to 8 lines wide, green above, pale



beneath, entire or with a few low serrations in the upper half; petioles 2 lines long; pedicels and peduncles pale and glaucous; flowers about 2 lines broad, in terminal compound corymbs 1 to 3 in. across; calyx teeth 5, triangular, blunt, about the length of the short-campanulate tube, distinctly glaucous; petals 5, white ovate-orbicular, obtuse; stamens 15 to 20, persistent; styles 5 to 6; follicles, in the specimens examined, 5 to 6, apparently sterile, included in the persistent calyx."

"On damp rocks along the Monongahela river, Morgantown, West Virginia, collected by Dr. C. F. Millspaugh in flower, June 20, 1890, and in apparently imperfect fruit late in September. Collected also by Mr. G. R. Vasey in the mountains of North Carolina, 1878."

"*Spiræa betulæfolia*, Pall, and *S. corymbosa*, Raf., have much longer follicles exerted beyond the calyx; broader, thicker and dentate leaves, and are different in habit. Rafinesque published a number of species in his New Flora, but none of them can apply to this one."

***Spiræa salicifolia* Linn. (Nuttall).**

*SPIRÆA TOMENTOSA* Linn. (Nuttall).

*ULMARIA RUBRA* Hill. *Spiræa rubra* Britton.

*ARUNCUS ARUNCUS* (Linn.) Karst. *Spiræa Aruncus* Linn.

*PORTERANTHUS TRIFOLIATUS* (Linn.) Britton. *Gillenia trifoliata* Linn.  
Greenbrier Co., on Kate's Mountain, alt. 3,300 ft. (*Small & Vail*).

*PORTERANTHUS STIPULATUS* (Muhl.) Britton. *Gillenia stipulacea* Nutt.

*RUBUS ODORATUS* Linn.

*RUBUS ODORATUS COLUMBIANUS* Millsp. Fl. W. Va., 356 (1892).

Leaves thin, ample, 5 to 7 incised almost to the petiole, divisions oblong-lanceolate, long and taper pointed, sharply and mostly double-serrate. Inflorescence smaller and more compact than in the species. Fruit also smaller and of a more decided musky flavor. Monongalia Co., cool woods of Tibb's Run (*Millspaugh*).

*RUBUS STRIGOSUS* Michx.

*RUBUS OCCIDENTALIS* Linn.

*RUBUS VILLOSUS* Ait.

*RUBUS VILLOSUS FRONDOSUS* Torr.

*RUBUS BAILEYANUS* Britton. *R. villosus humifusus* T. & G.

*RUBUS MILLSPAUGHI* Britton, Bull. Torr. Club, 18:366 (1891).

"Ascending, wand-like, entirely unarmed or with a very few weak prickles above, glabrous throughout or the younger shoots scurfy pubescent. Stems one and one-half to four meters long; leaves long petioled, pedately 5-foliolate or some of those on the twigs 3-foliolate; leaflets thin, oval, glabrous on both sides, long-acuminate at the apex, mostly rounded at the base, 12 to 15 cm. long, about 5 cm. wide, sharply, but not deeply serrate; stock of the terminal leaflet 7 to 10 cm. long; inflorescence loosely racemose; bracts linear-lanceolate, acuminate; fruit black, about 10 mm. long" (*Millspaugh*).

“Nearest to *R. villosus*, but evidently a distinct species. Curiously enough there is a leaf of this species glued down on the sheet of *R. Canadensis*, L. in herb Linn., and it appears to have been included in his description of that species—the specimens of which were furnished by Kalm.”

Near the summit of Point Mountain in Randolph County at an altitude of 3,500 ft., also along the Gandy in great profusion. Pendleton and Pocahontas, on Little Rich Mountains abundant. The mountaineers claim that it is upon this species that the bears grow fat for their period of hibernation, the fruit being late to ripen and very nutritious.

Mr. John K. Small remarks in his Studies of the Botany of the Southeastern United States, Bull. Torr. Club, 21:19 (1894) in reporting this species: “Grows on the walls of the cañon at Tallulah Falls, Ga. Collected in flower in April, 1893; alt. 1,600 ft. This most likely locates the southern limit of the geographical range of *R. Millspaughii*. Being a typical high mountain species, it is not to be expected much below 1,600 feet, and just south of the above-mentioned locality the Blue Ridge ‘runs out’ into the plains. Further south the land decreases gradually in altitude until the gulf is reached. According to my observations the above plant thrives most vigorously at altitudes ranging from 4,500 to 6,000 ft. It grows well at lower elevations, but never in such a prolific manner.” Mr. Small and Anna M. Vail also give in their “Report of the Botanical Exploration of Southwestern Virginia, 1892,” the following additional localities for this species: “Above: Fox Creek, on Pine Mountain in deep woods, alt. 3,000 ft. Summit and slopes of Mt. Rodgers, alt. 4,000 to 5,719 ft. Summit of White Top Mountain, alt. 5,678 ft. Slopes and summit of White Rock Mountain, alt. 3,000 to 4,400 ft.”

The species was also collected by Mr. T. H. Kearney, Jr., at the summit of Thunderhead (about 6,500 ft.), and on Chilhowee Mountain (at about 1,500 ft.), in Blount Co., Tennessee. Reports of the collection of the species in the mountains of North Carolina, New Hampshire, and northern Michigan have been made.

The species is reported as common in thickets at Wilton and Southport, Maine (*Fernald*), and at Deering, Maine (*Blankinship*).

#### RUBUS CANADENSIS Linn.

RUBUS CANADENSIS RORIBACCUS Bailey, Am. Gard., 11: 642 (1890).

Dry hillsides, Randolph Co., near Beverly. “Plant larger and stronger than the species; leaflets broad below, usually triangular-ovate, doubly serrate with small teeth, and more or less notched and jagged; peduncles longer, straighter and stouter, habituously more numerous and more conspicuously overtopping the leaves; flowers very large (sometimes two inches across); sepals uniformly larger, some of them much prolonged and leaf-like and conspicuously lobed (sometimes becoming an inch long and wide); fruit much larger.”

- RUBUS HISPIDUS Linn.  
 RUBUS TRIVALIS Michx.  
 DALIBARDA REPENS Linn.  
 GEUM CANADENSE Jacq.  
 GEUM VIRGINIANUM Linn. (*Nuttall*).  
 GEUM VERNUM (Raf.) T. & G.  
 WALDSTEINIA FRAGARIOIDES (Michx.) Tratt.  
 FRAGARIA VIRGINIANA Duchesne.  
 FRAGARIA VESCA Linn.  
 FRAGARIA INDICA Andr.  
 POTENTILLA MONSPELIENSIS Linn. *P. Norvegica* Linn.  
 POTENTILLA CANADENSIS Linn. *P. Canadensis simplex* T. & G.  
 AGRIMONIA STRIATA Michx. *A. Eupatoria* Linn.  
 AGRIMONIA PARVIFLORA Soland.  
 SANGUISORBA CANADENSIS Linn. *Poterium Canadense* A. Gray.  
 ROSA CAROLINA Linn. (*Nuttall*).  
 ROSA HUMILIS Marsh.  
 ROSA LUCIDA Ehrh. *Rosa humilis lucida* Best.  
 ROSA RUBIGINOSA Linn.  
 ROSA CANINA Linn.  
 PYRUS CORONARIA Linn.  
 PYRUS ANGUSTIFOLIA Ait.  
 ABRONIA ARBUTIFOLIA (Linn.) Ell. *Pyrus arbutifolia* Linn. f.  
 ABRONIA NIGRA (Willd.) Britton. *Pyrus melanocarpa* Hook.  
 SORBUS AMERICANA Marsh. *Pyrus Americana* DC.  
 CRATEGUS SPATHULATA Michx.  
 CRATEGUS CORDATA Ait.  
 CRATEGUS OXYACANTHA Linn.  
 CRATEGUS APIIFOLIA (Marsh.) Michx.  
 CRATEGUS COCCINEA Linn.  
 CRATEGUS MOLLIS (T. & G.) Scheele.  
 CRATEGUS TOMENTOSA Linn.  
 CRATEGUS PUNCTATA Jacq.  
 Large trees, of the form with bright yellow fruit, plentiful on Channel Ridge, Randolph Co. (*Millspaugh*).  
 CRATEGUS CRUS-GALLI Linn.  
 CRATEGUS FLAVA Ait. *C. flexispina* Sarg.  
 CRATEGUS FLAVA PUBESCENS A. Gray. *C. flexispina pubescens* Sarg.  
 CRATEGUS UNIFLORA Moench.  
**Crategus rotundifolia** (Ehrh.) Bork.  
 Greenbrier Co., on Kate's Mountain, alt. 3,300 ft. (*Small & Vail*).  
 AMELANCHIER CANADENSIS (Linn.) Medic.

## LEGUMINOSÆ.

BAPTISIA TINCTORIA (Linn.) R. Br.

BAPTISIA VILLOSA (Walt.) Nutt.

BAPTISIA AUSTRALIS (Linn.) R. Br.

LUPINUS PERENNIS Linn.

MEDICAGO SATIVA Linn.

MEDICAGO LUPULINA Linn.

MELILOTUS OFFICINALIS (Linn.) Lam.

MELILOTUS ALBA Lam.

TRIFOLIUM ARVENSE Linn.

TRIFOLIUM PRATENSE Linn.

TRIFOLIUM REPENS Linn.

**Trifolium Virginicum** Small, Mem. Torr. Club, 4:112 (1894)  
Tab. 75.

Perennial from a large and long root, diffusely branched from the summit of the root; branches 2 to 4 cm. long, strictly prostrate, pubescent; leaves 3-foliolate, petiole 4 to 8 cm. long; leaflets linear, linear-lanceolate or oblanceolate, 1 to 4 cm. long, acute or cuspidate, serrate-dentate, glabrous above, more or less silky beneath, conspicuously veined; sepals ovate, conspicuous; inflorescence in terminal, globose heads, about 2.5 cm. in diameter; flowers whitish, more or less crowded on slender pedicels, .2 to .4, standard emarginate-mucronate, striate; calyx clothed with long silky hairs, the teeth subulate, nearly half the length of the corolla. Pods and seeds not seen.

Growing on the rocky slopes of Kate's Mountain, Greenbrier Co., in company with *Clematis ovata*.

This species is most closely related to *T. stoloniferum* by its flower, but in all other respects it differs from that and all the other eastern American species.

TRIFOLIUM HYBRIDUM Linn.

TRIFOLIUM AGRARIUM Linn.

TRIFOLIUM PROCUMBENS Linn.

CRACCA VIRGINIANA Linn. *Tephrosia Virginiana* Pers.

Webster Co. on an "island" in Upper Glade (*Millsbaugh*).

ROBINIA PSEUDACACIA Linn.

ROBINIA HISPIDA Linn.

ASTRAGALUS CAROLINIANUS Linn.

**Astragalus distortus** T. & G.

Since the publication of the Preliminary Catalogue, Dr. Gamble has succeeded in securing fruits of this species, reported generically only in that work. This is the farthest eastern extension of the species, which ranges, according to the manual, from Illinois to Iowa, Missouri, Kansas and Texas.

STYLOSANTHES BIFLORA (Linn.) B. S. P.

STYLOSANTHES HAMATA (Linn.) Britton.

- MEIBOMIA NUDIFLORA (Linn.) O.K. *Desmodium nudiflorum* DC.  
 MEIBOMIA GRANDIFLORA (Walt.) O.K. *Desmodium grandiflorum* DC.  
 MEIBOMIA ROTUNDIFOLIA (Michx.) O.K. *Desmodium rotundifolium* DC.  
 MEIBOMIA OCHROLEUCA (M. A. Curtiss) O.K. *Desmodium ochroleucum*  
 M. A. Curtiss.  
 MEIBOMIA CANESCENS (Linn.) O.K. *Desmodium canescens* DC.  
 MEIBOMIA BRACTEOSA (Michx.) O.K. *Desmodium cuspidatum* Hook.  
 MEIBOMIA DILLENII (Darl.) O.K. *Desmodium Dillenii* Darl.  
 MEIBOMIA PANICULATA (Linn.) O.K. *Desmodium paniculatum* DC.  
 MEIBOMIA CANADENSIS (Linn.) O.K. *Desmodium Canadense* DC.  
 MEIBOMIA RIGIDA (Ell.) O.K. *Desmodium rigidum* DC.  
 MEIBOMIA OBTUSA (Muhl.) O.K. *Desmodium ciliare* DC.  
 MEIBOMIA MARILANDICA (L.) O.K. *Desmodium Marilandicum* Britt.  
 MEIBOMIA LINEATA (Michx.) O.K. *Desmodium lineatum* DC.  
 LESPEDEZA REPENS (L.) Bart.  
 LESPEDEZA FRUTESCENS (L.) Britton. *L. Stuevei intermedia* Watson.  
*Lespedeza reticulata* Pers.  
 LESPEDEZA VIOLACEA (L.) Pers.  
 LESPEDEZA VIRGINICA (L.) Britton.  
 LESPEDEZA STUVEI Nutt.  
 LESPEDEZA HIRTA (L.) Ell. *Lespedeza polystachya* Michx.  
 LESPEDEZA CAPITATA Michx.  
 LESPEDEZA STRIATA (Thumb.) Hook & Arn.  
 VICIA CAROLINIANA Walt.  
 Greenbrier Co., Kate's Mountain White Sulphur Springs, alt.  
 3,000 ft. (*Small & Vail*).  
 LATHYRUS VENOSUS Muhl.  
 Greenbrier Co., Kate's Mountain, alt. 3,300 ft. (*Small & Vail*).  
 FALCATA COMOSA (L.) O.K. *Amphicarpæa comosa* Ridd.  
 APIOS APIOS (L.) MacM. *Apios tuberosa* Moench.  
 PHASEOLUS POLYSTACHYUS (Linn.) B. S. P.  
 PHASEOLUS HELVOLUS Linn.  
 GLEDITSCHIA TRIACANTHOS Linn.  
 CASSIA MARILANDICA Linn.  
 CASSIA CHAMÆCHRISTA Linn.  
 CASSIA NICTITANS Linn.  
 CERCIS CANADENSIS Linn.  
 GYMNOCLADUS DIOICUS (Linn.) Koch. *Gymnocladus Canadensis* Lam.

GERANIACEÆ.

- GERANIUM MACULATUM Linn.  
 GERANIUM ROBERTIANUM Linn.  
 GERANIUM CAROLINIANUM Linn.

## OXALIDACEÆ.

- OXALIS ACETOSELLA Linn. (*Nuttall.*)  
 OXALIS VIOLACEA Linn.  
 OXALIS STRICTA Linn. *Oxalis corniculata stricta* Sav.  
 OXALIS RECURVA Ell.

## LINACEÆ.

- LINUM VIRGINIANUM Linn.  
 LINUM STRIATUM Walt.  
 LINUM USITATISSIMUM Linn.

## RUTACEÆ.

- ZANTHOXYLUM AMERICANUM Mill.  
 RUTA GRAVEOLENS Linn.  
 PTELEA TRIFOLIATA Linn.

## SIMARUBACEÆ.

- AILANTHUS GLANDULOSUS Desf.

## POLYGALACEÆ.

- POLYGALA VIRIDESCENS Linn. *Polygala sanguinea* Linn.  
 POLYGALA VIRIDESCENS ALBIFLORA Wheelock. *Polygala sanguinea albi-  
 flora* Millsp.  
 POLYGALA MARIANA Mill.  
 POLYGALA CURTISSII A. Gray.  
 POLYGALA CRUCIATA Linn.  
 POLYGALA VERTICILLATA Linn.  
 POLYGALA AMBIGUA Nutt.  
 POLYGALA NUTTALLII T. & G.  
 POLYGALA SENEGA Linn.  
 POLYGALA PAUCIFOLIA Willd.  
 Greenbrier Co., Kate's Mountain, alt. 3,300 ft. (*Small & Vail.*)

## EUPHORBIACEÆ.

- EUPHORBIA GLYPTOSPERMA PUBESCENS Engelm.  
 EUPHORBIA MACULATA Linn.  
 Mineral Co., opposite Cumberland, Md. (*J. K. Small.*)  
 EUPHORBIA PRESII Guss.  
 EUPHORBIA MARGINATA Pursh.  
 EUPHORBIA COROLLATA Linn.  
 EUPHORBIA DENTATA Michx.  
 EUPHORBIA DARLINGTONII Gray.  
 EUPHORBIA OBTUSATA Pursh.

Dr. W. F. G. ...

I have been ... along the ... and ...

PL. VII RHUS RADICANS Linn.

---

This species completely covers a half mile of fence along the Monongahela river below Morgantown. The growth here is remarkably rich in foliage and very characteristic in all its features.





*RHUS RADICANS* Linn.



EUPHORBIA CYPARISSIAS Linn.

EUPHORBIA COMMUTATA Engelm.

Greenbrier Co., near White Sulphur Springs (*J. H. Redfield*).

EUPHORBIA LATHYRIS Linn.

ACALYPHA VIRGINICA Linn.

ACALYPHA VIRGINICA INTERMEDIA Muell.

[COREMA CONRADI Torr., (Empetraceæ) mentioned in Botanical Gazette, Vol. II, p. 136, as occurring near Hawk's Nest, Fayette Co., is proven by Professor James to be another species, and should therefore not yet be credited to this state.]

### CALLITRICHACEÆ.

CALLITRICHÆ HETEROPHYLLA Pursh.

### LIMNANTHACEÆ.

FLGERKEA PROSERPINACOIDES Willd.

### ANACARDIACEÆ.

RHUS HIRTA (Linn.) Sudw. *Rhus typhina* Linn.

RHUS GLABRA Linn.

RHUS COPALLINA Linn.

RHUS VERNIX Linn. *Rhus venenata* Linn.

RHUS RADICANS Linn.

RHUS AROMATICA Ait. *Rhus Canadensis* Marsh.

### AQUIFOLIACEÆ.

ILEX OPACA Ait. —

ILEX MONTICOLA A. Gray.

ILEX MONTICOLA MOLLIS (A. Gray) Britton. *Ilex mollis* A. Gray.

ILEX VERTICILLATA (Linn.) A. Gray.

### CELASTRACEÆ.

EUONYMUS ATROPURPUREUS Jacq.

EUONYMUS AMERICANUS Linn.

**Euonymus obovatus** Nutt.

*E. Americanus* var. *obovatus* T. & Gray. Of this species Rev. A. Boutlou says: "In Marshall Co., at Board Tree, Cameron, Belton, etc., this is the most common form; while the trailing rooting form I have met almost everywhere I have been in the State."

CELASTRUS SCANDENS Linn.

### STAPHYLEACEÆ.

STAPHYLEA TRIFOLIA Linn.

## ACERACEÆ.

## ACER PENNSYLVANICUM Linn.

Although in most situations in the State this species is a mere shrub, yet near Welsh in Grant Co. it assumes the proportions of a beautiful tree 20 to 40 ft. high with a strict columnar trunk 5 to 8 inches in diameter (*Millspaugh*). Greenbrier Co., on Kate's Mountain, near White Sulphur Springs, alt. 3,300 ft. (*Small & Vail*).

## ACER SPICATUM Lam.

## ACER SACCHARUM Marsh.

In Randolph Co., near Pickens, the sugar maple grows to a diameter of 2 to 4 ft., and a height of 50 to 70 ft. (*Millspaugh*).

ACER NIGRUM Michx. f. *Acer saccharum* var. *nigrum* Britton.

## ACER SACCHARINUM Linn.

Large trees 2 to 4 ft. in diameter in Randolph Co., near Pickens. At one station in Webster Co., on Buffalo Bull Mountain, nearly every tree of this species is of the Bird's-eye type. (*Millspaugh*).

## ACER RUBRUM Linn.

ACER NEGUNDO Linn. *Negundo aceroides* Mœench. (*Nuttall*).

## HIPPOCASTANACEÆ.

## ÆSCULUS GLABRA Willd.

## ÆSCULUS OCTANDRA Marsh.

ÆSCULUS OCTANDRA HYBRIDA (DC.) Sarg. *Æsculus octandra purpurascens* A. Gray.

## ÆSCULUS PAVIA Linn.

## BALSAMINACEÆ.

## IMPATIENS AUREA Muhl.

## IMPATIENS BIFLORA Walt.

## RHAMNACEÆ.

## RHAMNUS CAROLINIANA Walt.

## CEANOTHUS AMERICANUS Linn.

## VITACEÆ.

## VITIS LABRUSCA Linn.

An immense specimen of this grape was found in Randolph Co., near Pickens; its trunk measured 22 inches in diameter and its two main branches 8 inches each. (*Millspaugh*).

## VITIS ÆSTIVALIS Michx.

## VITIS CORDIFOLIA Michx.

VITIS VULPINA Linn. *Vitis riparia* Michx.

## VITIS RUPESTRIS Scheele.

## VITIS ROTUNDFOLIA Michx.

PARTHENOCISSUS QUINQUEFOLIA (Linn.) Planch. *Vitis quinquefolia* Lam.  
 AMPELOPSIS CORDATA Michx. *Cissus Ampelopsis* Pers.

TILIACEÆ.

TILIA AMERICANA Linn.

Large growths of this species found at Pickens, Randolph Co., with trunk diameters of 26 to 38 inches, and height 60 to 100 feet. (*Millsbaugh*), (*Nuttall*).

TILIA HETEROPHYLLA Vent.

One specimen cut at Nuttallburg, Fayette Co., had a diameter of 26 inches. (*Millsbaugh*), (*Nuttall*).

MALVACEÆ.

ALTHÆA ROSEA Linn.

MALVA ROTUNDIFOLIA Linn.

MALVA MOSCHATA Linn.

SIDA SPINOSA Linn.

SIDA HERMAPHRODITA (Linn.) Rusby. *Sida Napæa* Cav.

ABUTILON ABUTILON (Linn.) Rusby. *Abutilon Avicenne* Gærtn.

HIBISCUS MOSCHEUTOS Linn.

HIBISCUS TRIONIUM Linn.

HYPERICACEÆ.

ASCYRUM HYPERICOIDES Linn. *Ascyrum Crux-Andree* Linn.

HYPERICUM PROLIFICUM Linn.

HYPERICUM DENSIFLORUM Pursh.

HYPERICUM VIRGATUM ACUTIFOLIUM Coulter.

HYPERICUM PERFORATUM Linn.

HYPERICUM MACULATUM Walt.

HYPERICUM MUTILUM Linn.

HYPERICUM CANADENSE Linn.

HYPERICUM GENTIANOIDES (Linn.) B. S. P.

HYPERICUM ADPRESSUM Barton.

HYPERICUM ELLIPTICUM Hook.

**Hypericum Virginicum** Linn. (*Nuttall*).

CISTACEÆ.

HELIANTHEMUM MAJUS (Linn.) B. S. P.

LECHEA MINOR Linn.

LECHEA LEGGETTII Britton & Holl.

VIOLACEÆ.

VIOLA PEDATA Linn.

Greenbrier Co., on Kate's Mountain, alt. 3,300 ft., near White Sulphur Springs (*Small & Vail*). Mercer Co., near Bluefield. MacDowell Co., at Fletcher's. (*Millsbaugh*).

VIOLA PEDATA BICOLOR Pursh.

Greenbrier Co., on Kate's Mountain, alt. 3,300 ft. (*Small & Vail*).

VIOLA PALMATA Linn.

VIOLA OBLIQUA Hill. *Viola cucullata* Ait.

VIOLA SAGITTATA Ait.

VIOLA BLANDA Willd.

VIOLA PRIMULÆFOLIA Linn.

VIOLA LANCEOLATA Linn.

VIOLA ROTUNDIFOLIA Michx.

VIOLA PUBESCENS Ait.

VIOLA SCABRIUSCULA (T. & G.) Schwein. *Viola pubescens scabriuscula* T. & G.

VIOLA HASTATA Michx.

VIOLA CANADENSIS Linn.

VIOLA STRIATA Ait.

**Viola Labradorica** Schrank.

*Viola canina* var. *Muhlenbergii*, Traut. Fayette Co., at foot of cliff at Nuttallburg (*Nuttall*).

VIOLA ROSTRATA Pursh.

VIOLA TENELLA Muhl.

SOLEA CONCOLOR (Forst.) Ging.

#### PASSIFLORACEÆ.

PASSIFLORA LUTEA Linn.

PASSIFLORA INCARNATA Linn.

#### CACTACEÆ.

OPUNTIA POLYACANTHA Haw. *Opuntia vulgaris* Linn.

#### THYMELÆACEÆ.

DIRCA PALUSTRIS Linn.

Kanawha Co., abundant on Blue Creek (*Rev. A. Boutlou*).

#### LYTHRACEÆ.

CUPHÆA PETIOLATA (Linn.) Kœhne.

#### MELASTOMACEÆ.

RHEXIA VIRGINICA Linn.

#### ONAGRACEÆ.

EPILOBIUM SPICATUM Muhl.

EPILOBIUM COLORATUM Muhl.

LUDWEGIA ALTERNIFOLIA Linn.

LUDWEGIA ALTERNIFOLIA LINEARIFOLIA Britton, Bull. Torr. Club, 17: 315 (1890).

"Two or three feet high, divergently branched, the branches ascending. Leaves linear, elongated, 2 to 4 in. long, 1.5 to 4 lines wide, acute; flowers solitary in the axils of the upper leaves or bracts, yellow; sepals ovate-lanceolate acute, narrower than those of *L. alternifolia*; branches and both sides of the leaves somewhat pubescent. Petals apparently remaining on the plant longer than those of *L. alternifolia*, which, as Dr. Millspaugh observes, commonly fall away when the plant is shocked."

"Appearing very distinct from typical *L. alternifolia*, but presumably but a variety of it." From the description it may be the *Rhexia linearifolia* Poir, in Lam. Encyl. vi. 2, said to come from Carolina."

Wood Co., near Lockhart's Run. (*Millspaugh*), (*Nuttall*).

LUDWEGIA PALUSTRIS (Linn.) Ell.

ONAGRA BIENNIS (Linn.) Scop. *Oenothera biennis* Linn. *O. biennis grandifolia* Lindl.

ŒNOTHERA\* PUMILA Linn.

ŒNOTHERA\* FRUTICOSA Linn.

ŒNOTHERA\* FRUTICOSA DIFFERTA Millsp. Fl. W. Va., 366 (1892).

Damp meadows, Wood Co., near Lockhart's Run, the most common form.

Stems 1 to 2 ft. high, nearly smooth, branching diffusely from every axil. Flowers profuse, large. Lower leaves ovate. Capsules narrowly winged, very short; apical inflorescence strongly cymose.

ŒNOTHERA\* LINEARIS Michx. *Oenothera fruticosa* var. *linearis* Watson.

GAURA BIENNIS Linn.

CIRCÆA LUTETIANA Linn.

CIRCÆA ALPINA Linn.

#### ARALIACEÆ.

ARALIA SPINOSA Linn.

ARALIA RACEMOSA Linn.

ARALIA NUDICAULIS Linn.

ARALIA HISPIDA Vent.

PANAX QUINQUEFOLIUM Linn. *Aralia quinquefolia* Dec. & Pl.

#### UMBELLIFERÆ.

HYDROCOTYLE AMERICANA Linn.

ERYNGIUM AQUATICUM Linn.

DAUCUS CAROTA Linn.

DAUCUS CAROTA *forma* ROSEA Millsp. Fl. W. Va., 369 (1892).

ANGELICA CURTISSII Buckley.

ANGELICA VILLOSA (Walt.) B. S. P.

\*See foot note page 168.

- ANGELICA ATROPURPUREA Linn.  
 OXYPOLIS RIGIDUS (Linn.) Britton. *Tiedemannia rigida* Coult. & Rose.  
 HERACLEUM LANATUM Michx.  
 PASTINACA SATIVA Linn.  
 THASPIUM TRIFOLIATUM (Nutt.) Britton. *Thaspium aureum* Nutt.  
*Thaspium aureum cordatum* Walt.  
 THASPIUM BARBINODE (Michx.) Nutt.  
 LIGUSTICUM CANADENSE (Linn.) Britton. *Ligusticum actaeifolium* (Michx).  
 DEERINGIA CANADENSIS (Linn.) O. K.  
 ZIZIA CORDATA (Walt.) DC.  
 ZIZIA AUREA (Linn.) Koch.  
 ZIZIA BEBBII (C. & R.) Britton.  
 Greenbrier Co., on Kate's Mountain, alt. 3,300 ft. (*Small & Vail*).  
 CICUTA MACULATA Linn.  
 CICUTA BULBIFERA Linn.  
 CHÆROPHYLLUM PROCUMBENS (Linn.) Crantz.  
 OSMORRHIZA CLAYTONI (Michx.) B. S. P.  
 OSMORRHIZA LONGISTYLIS (Torr.) DC.  
 ERIGENIA BULBOSA (Michx.) Nutt.  
 SANICULA MARYLANDICA Linn.  
 SANICULA CANADENSIS Linn.

## CORNACEÆ.

- CORNUS FLORIDA Linn.  
 Greenbrier Co., on Kate's Mountain, alt. 3,300 ft. (*Small & Vail*).  
 CORNUS CIRCINATA L'Her.  
 CORNUS AMONUM Mill. *Cornus sericea* Linn.  
 CORNUS CANDIDISSIMA Marsh.  
 CORNUS ALTERNIFOLIA Linn. f.  
 NYSSA AQUATICA Linn.  
*Nyssa salvatika* Marsh. Immense growths of this species at Pickens, Randolph Co., with trunks 3 to 4 ft. in diameter. (*Millsbaugh*).

## CLETHRACEÆ.

- CLETHRA ACUMINATA Michx.

## PYROLACEÆ.

- CHIMAPHILA UMBELLATA (Linn.) Nutt. *Pseva umbellata* O. K.  
 CHIMAPHILA MACULATA (Linn.) Pursh. *Pseva maculata* O. K.  
 MONESES UNIFLORA (Linn.) A. Gray. *Moneses grandiflora* Salisb.  
 PYROLA ELLIPTICA Nutt.  
 PYROLA ROTUNDIFOLIA Linn.



MONOTROPACEÆ.

MONOTROPA UNIFLORA Linn.

HYPOPYTIS HYPOPYTIS (Linn.) Small. *Hypopytis Monotropa* Crantz.

ERICACEÆ.

GAYLUSSACIA DUMOSA (Andr.) T. & G.

GAYLUSSACIA FRONDOSA (Linn.) T. & G.

GAYLUSSACIA RESINOSA (Ait.) T. & G.

OXYCOCCUS MACROCARPUS Pers.

OXYCOCCUS ERYTHROCARPUS Pers. *Vaccinium erythrocarpon* Michx.

VACCINIUM STAMINEUM Linn.

VACCINIUM PENNSYLVANICUM Lam.

VACCINIUM VACILLANS Soland.

VACCINIUM CORYMBOSUM Linn.

VACCINIUM PALLIDUM Ait. *Vaccinium corymbosum pallidum* A. Gray.

CHIOGENES HISPIDULA (Linn.) T. & G.

GAULTHERIA PROCUMBENS Linn.

EPIGÆA REPENS Linn.

XOLISMA LIGUSTRINA (Linn.) Britton. *Andromeda ligustrina* Muhl.

XOLISMA LIGUSTRINA PUBESCENS A. Gray. *Andromeda ligustrina pubescens* A. Gray.

PIERIS MARIANA (Linn.) Benth & Hook. *Andromeda Mariana* Linn.

OXYDENDRON ARBOREUM (Linn.) DC.

KALMIA LATIFOLIA Linn.

KALMIA AUGUSTIFOLIA Linn.

MENZIESIA PILOSA (Michx.) Pers. *Menziesia globularis* Salisb.

RHODODENDRON MAXIMUM Linn.

RHODODENDRON CATAWBIENSE Michx.

AZALEA ARBORESCENS Pursh. *Rhododendron arborescens* Torr.

AZALEA CANESCENS Michx. *Rhododendron canescens* Porter.

Greenbrier Co., Kate's Mountain, alt. 3,300 ft. (*Small & Vail*).

AZALEA VISCOSA Linn. *Rhododendron viscosum* Torr.

AZALEA VISCOSA GLAUCA Michx. *Rhododendron viscosum glaucum* A. Gray.

AZALEA VISCOSA NITIDA (Pursh.) Britton. *Rhododendron viscosum nitidum* A. Gray.

AZALEA NUDIFLORA Linn. *Rhododendron nudiflorum* Torr.

AZALEA LUTEA Linn. *Rhododendron calendulaceum* Torr.

DIAPENSIACEÆ.

GALAX APHYLLA Linn.

PRIMULACEÆ.

DODECATHEON MEADIA Linn.

TRIENTALIS AMERICANA Pursh.

STEIRONEMA CILIATUM (Linn.) Baudo.

Greenbrier Co., on the mountains about White Sulphur Springs, alt. 3,200 ft. (*Small & Vail*).

STEIRONEMA LANCEOLATUM (Walt.) A. Gray.

STEIRONEMA LANCEOLATUM ANGUSTIFOLIUM (Lam.) A. Gray.

LYSIMACHIA QUADRIFOLIA Linn.

LYSIMACHIA TERRESTRIS (Linn.) B. S. P.

LYSIMACHIA NUMMULARIA Linn.

NAUMBERGIA THYRSIFLORA (Linn.) Duby. *Lysimachia thyrsoflora* Linn.

ANAGALLIS ARVENSIS Linn. (*Nuttall*).

SAMOLUS FLORIBUNDUS H. B. K. *Samolus Valerandi floribundus* B. S. P.

#### EBENACEÆ.

DIOSPYROS VIRGINIANA Linn.

#### STYRACEÆ.

MOHRODENDRON CAROLINUM (Linn.) Britton. *Halesia tetraptera* Linn.

#### OLEACEÆ.

FRAXINUS AMERICANA Linn.

Fine specimens of very large growth in Randolph Co., especially on Channel ridge where the species grows very tall, and at Pickens where specimens were measured from 3 to 6 ft. in diameter. (*Millsbaugh*).

FRAXINUS PENNSYLVANICA Marsh. *F. pubescens* Lam.

FRAXINUS LANCEOLATA Borck. *F. viridis* Michx. f.

FRAXINUS NIGRA Marsh. *F. sambucifolia* Lam.

CHIONANTHUS VIRGINICA Linn.

LIGUSTRUM VULGARE Linn.

#### LOGANIACEÆ.

CYNOCTONUM MITREOLA (Linn.) Britton. *Mitreola petiolata* T. & G.

#### GENTIANACEÆ.

SABBATIA ANGULARIS (Linn.) Pursh.

GENTIANA QUINQUEFOLIA Linn.

GENTIANA ANDREWSII Griseb.

GENTIANA SAPONARIA Linn.

GENTIANA LINEARIS Frœl.

OBOLARIA VIRGINICA Linn.

#### APOCYNACEÆ.

APOCYNUM ANDROSÆMIFOLIUM Linn.

APOCYNUM CANNABINUM Linn.

APOCYNUM CANNABINUM PUBESCENS (R. Br.) A. DC.

ASCLEPIADACEÆ.

- ASCLEPIAS TUBEROSA Linn.  
 ASCLEPIAS RUBRA Linn.  
 ASCLEPIAS PURPURASCENS Linn.  
 ASCLEPIAS VARIEGATA Linn.  
 ASCLEPIAS INCARNATA Linn.  
 ASCLEPIAS PULCHRA Ehrh. *Asclepias incarnata pulchra* Pers.  
 ASCLEPIAS SYRIACA Linn. (*Nuttall.*)  
 ASCLEPIAS EXALTATA (Linn.) Muhl.  
 ASCLEPIAS QUADRIFOLIA Jacq.  
 ACERATES VIRIDIFLORA (Raf.) Eaton.  
 VINCETOXICUM GONOCARPUS LÆVIS (Michx.) Britton. *Gonolobus lævis*  
 Michx.

CONVOLVULACEÆ.

- IPOMŒA COCCINEA Linn.  
 IPOMŒA HEDERACEA Jacq.  
 IPOMŒA PURPUREA (Linn.) Roth.  
 Mineral Co., opposite Cumberland, Md. (*J. K. Small.*)  
 IPOMŒA PANDURATA (Linn.) Meyer.  
 IPOMŒA LACUNOSA Linn.  
 CONVULVULUS SPITHAMÆUS Linn.  
 CONVULVULUS SEPIUM Linn. (*Nuttall.*)  
 CONVULVULUS REPENS Linn. *C. sepium repens* A. Gray.

CUSCUTACEÆ.

- CUSCUTA EPITHYMUM Murr. *C. Trifolii* Weihe.  
 CUSCUTA GRONOVII Willd.  
 CUSCUTA GLOMERATA Choisy.

POLEMONIACEÆ.

- Phlox paniculata** Linn. (*Nuttall.*)  
 PHLOX PANICULATA ACUMINATA (Pursh.) Chapm.  
 PHLOX MACULATA Linn.  
 PHLOX AMENA Sims.  
 PHLOX REPTANS Michx.  
 PHLOX SUBULATA Linn.  
 Greenbrier Co., dry, stony ledges on Kate's Mountain, alt.  
 3,300 ft. (*Small & Vail.*)  
 POLEMONIUM REPTANS Linn.  
 POLEMONIUM VAN BRUNTIÆ Britton. *P. carulcum* Linn.

HYDROPHYLLACEÆ.

- HYDROPHYLLUM MACROPHYLLUM Nutt.  
 HYDROPHYLLUM VIRGINICUM Linn.

- HYDROPHYLLUM CANADENSE Linn.  
 HYDROPHYLLUM APPENDICULATUM Michx.  
 PHACELIA BIPINNATIFIDA Michx.  
 PHACELIA PURSHII Buckley.  
 PHACELIA DUBIA (Linn.) Small. *P. parviflora* Pursh.

## BORAGINACEÆ.

- CYNOGLOSSUM OFFICINALE Linn.  
 CYNOGLOSSUM VIRGINICUM Linn.  
 LAPPULA VIRGINIANA (Linn.) Greene. *Echinosperrnm Virginicum* Lehm.  
 MERTENSIA VIRGINICA (Linn.) DC.  
 ONOSMODIUM CAROLINIANUM (Lam.) A. DC.  
 MYOSOTIS PALUSTRIS (Linn.) Relh.  
 SYMPHYTUM OFFICINALE Linn.  
 LITHOSPERMUM ARVENSE Linn.  
 LITHOSPERMUM LATIFOLIUM Michx.  
 LITHOSPERMUM CANESCENS (Michx.) Lehm.  
 ECHIUM VULGARE Linn.

## VERBENACEÆ.

- VERBENA OFFICINALIS Linn.  
 VERBENA URTICÆFOLIA Linn.  
 VERBENA HASTATA Linn.  
 VERBENA ANGUSTIFOLIA Michx.  
 LIPPIA LANCEOLATA Michx.  
 PHRYMA LEPTOSTACHYA Linn.

## LABIATÆ.

- ISANTHUS BRACHIATUS (Linn.) B. S. P.  
 TEUCRIUM CANADENSE Linn.  
 COLLINSONIA CANADENSIS Linn.  
 PERILLA FRUTESCENS NANKINENSIS (Lour.) Britton. *P. ocyroides crispæ*  
 Benth.  
 MENTHA SPICATA Linn. *M. viridis* Linn.  
 MENTHA PIPERITA Linn.  
 MENTHA SATIVA Linn.  
 MENTHA CANADENSIS Linn.  
 LYCOPUS VIRGINICUS Linn.  
 LYCOPUS SINUATUS Ell.  
 CUNILA ORIGANOIDES (Linn.) Britton.  
 KÆLLIA FLEXUOSA (Walt.) Britton.  
 KÆLLIA VERTICILLATA (Michx.) O. K. *K. Torreyi* Benth.  
 KÆLLIA CLINIPODIOIDES (T. & G.) O. K.

- KÆLIA PYCANTHEMOIDES (Leavenw.) O. K. *K. Tullia* Benth.  
 KÆLIA INCANA (Linn.) O. K.  
 KÆLIA MONTANA (Michx.) O. K.  
 HEDEOMA PULEGIOIDES (Linn.) Pers.  
 CLINOPODIUM VULGARE Linn. *Calamintha Clinopodium* Benth.  
 MELISSA OFFICINALIS Linn.  
 SALVIA LYRATA Linn.  
 MONARDA DIDYMA Linn.  
 MONARDA FISTULOSA Linn. *M. fistulosa mollis* Benth.  
 MONARDA MEDIA Willd. *M. fistulosa rubra* A. Gray.  
 BLEPHILIA HIRSUTA (Pursh.) Torrey.  
 VLECKIA NEPETOIDES (Linn.) Raf. *Agastache nepetoides* O. K.  
 MEEHANIA CORDATA (Nutt.) Britton. *Cedronella cordata* Benth.  
 NEPETA CATARIA Linn.  
 GLECHOMA HEDERACEA Linn. *Nepeta hederacea* B. S. P.  
 SCUTELLARIA LATERIFLORA Linn.  
 SCUTELLARIA CORDIFOLIA Muhl. *S. versicolor minor* Chapm.  
 SCUTELLARIA SAXATILIS Riddell.  
 SCUTELLARIA SERRATA Andrews.  
 SCUTELLARIA INCANA Muhl. *S. canescens* Nutt.  
 SCUTELLARIA PILOSA Michx.  
 SCUTELLARIA PILOSA HIRSUTA (Short.) A. Gray.  
 SCUTELLARIA INTEGRIFOLIA Linn.  
 SCUTELLARIA PARVULA Michx.  
 SCUTELLARIA GALERICULATA Linn.  
 SCUTELLARIA GALERICULATA *forma* ALBIFLORA Millsp. Fl. W. Va. 428  
 (1892.)  
 SCUTELLARIA NERVOSA Pursh.  
 PRUNELLA VULGARIS Linn. *Brunella vulgaris* Linn.  
 PRUNELLA VULGARIS ALBIFLORA (Boggenh.) Britton.  
 PHYSOSTEGIA VIRGINIANA (Linn.) Benth.  
 MARRUBIUM VULGARE Linn.  
 STACHYS PALUSTRIS Linn.  
 STACHYS ASPERA Michx.  
 STACHYS ASPERA GLABRA A. Gray.  
 STACHYS CORDATA Ridd. (*Nuttall*).  
 GALEOPSIS TETRAHIT Linn.  
 LEONURUS CARDIACA Linn.  
 LAMIUM AMPLEXICAULE Linn.  
 TRICHOSTEMA DICHOTOMUM Linn.

SOLANACEÆ.

- SOLANUM DULCAMARA Linn.  
 SOLANUM NIGRUM Linn.

SOLANUM CAROLINENSE Linn.  
 PHYSALIS PHILADELPHICA Lam.  
 PHYSALIS ANGULATA Linn.  
 PHYSALIS PUBESCENS Linn.  
 PHYSALIS VIRGINIANA Mill.  
 PHYSALIS VISCOSA Linn.  
 PHYSALIS LANCEOLATA Michx.  
 PHYSALODES PHYSALODES (Linn.) Britton.  
 LYCIUM VULGARE (Ait. f.) Dun.  
 DATURA STRAMONIUM Linn.  
 DATURA TATULA Linn.  
 PETUNIA VIOLACEA Lindl.

## SCROPHULARIACEÆ.

VERBASCUM THAPSUS Linn.  
 VERBASCUM BLATTARIA Linn.  
 VERBASCUM LYCHNITIS Linn.  
 LINARIA LINARIA (Linn.) Karst. *L. vulgaris* Mill.  
 This species first appeared near Nuttallburg in Fayette Co.,  
 this year—1895 (*Nuttall*).  
 SCROPHULARIA MARILANDICA Linn. *S. nodosa Marilandica* A. Gray.  
 COLLINSIA VERNA Nutt.  
 CHELONE GLABRA Linn.  
 CHELONE OBLIQUA Linn.  
 PENTSTEMON HIRSUTUS (Linn.) Willd.  
 PENTSTEMON PENTSTEMON (Linn.) Britton. *P. lævigatus* Soland.  
 PENTSTEMON DIGITALIS (Sweet) Nutt. *P. lævigatus Digitalis* A. Gray.  
**Pentstemon canescens** Britton.  
 Slopes of Kate's Mountain, alt. 3,300 ft., near White Sulphur  
 Springs, Greenbrier Co. (*Small & Vail*).  
 MIMULUS RINGENS Linn.  
 MIMULUS ALATUS Soland.  
 GRATIOLA VIRGINIANA Linn.  
 GRATIOLA SPHEROCARPA Ell.  
 ILYSANTHES GRATIOLOIDES (Linn.) Benth.  
 LEPTANDRA VIRGINICA (Linn.) Nutt. *Veronica Virginia* Linn.  
 VERONICA ANAGALLIS AQUATICA Linn. *V. Anagallis* Linn.  
 VERONICA AMERICANA Schw.  
 VERONICA OFFICINALIS Linn.  
 VERONICA SERPYLLIFOLIA Linn.  
 VERONICA PEREGRINA Linn.  
 VERONICA ARVENSIS Linn.  
 BUECHNERA AMERICANA Linn.

- DASYSTOMA PEDICULARIA (Linn.) Benth. *Gerardia Pedicularia* Linn.  
 DASYSTOMA VIRGINICA (Linn.) Britton. *Gerardia Virginica* Linn.  
 DASYSTOMA FLAVA (Linn.) Wood. *Gerardia flava* Linn.  
 DASYSTOMA LÆVIGATA Raf. *Gerardia lævigata* Raf.  
 GERARDIA TENUIFOLIA Vahl.  
 GERARDIA AURICULATA Michx.  
 CASTILLEJA COCCINEA (Linn.) Spreng.  
 PEDICULARIS CANADENSIS Linn. (*Nuttall*).  
 MELAMPYRUM LINEARE Lam.

OROBANCHACEÆ.

- EPIPEGUS VIRGINIANA (Linn.) Bart.  
 CONOPHOLIS AMERICANA (Linn. f.) Wallr.  
 THALESIA UNIFLORA (Linn.) Britton. *Aphyllon uniflorum* A. Gray.

BIGNONIACEÆ.

**Bignonia crucigera** Linn.

*B. cupreolata* Linn. Kanawha Co., rich soil between Cannelton and Brownstown, where it grows luxuriantly along the banks of the Great Kanawha River. Its evergreen foliage makes it a conspicuous feature of the winter landscape (*Rev. A. Boutlou*).

- TECOMA RADICANS (Linn.) DC.  
 CATALPA CATALPA (Linn.) Karst. *C. Bignonioides* Walt.

**Catalpa speciosa** Warder.

Mason Co., banks of the Ohio River, near Point Pleasant. (*Millsbaugh*.)

ACANTHACEÆ.

- RUELLIA CILIOSA Pursh.  
 DIANTHERA AMERICANA Linn.

PLANTAGINACEÆ.

- PLANTAGO MAJOR Linn.  
 PLANTAGO RUGELII Decne.  
 PLANTAGO LANCEOLATA Linn.  
 PLANTAGO VIRGINICA Linn.

RUBIACEÆ.

- HOUSTONIA CÆRULEA Linn.  
 HOUSTONIA CÆRULEA *forma* ALBIFLORA Millsp. Fl. W. Va. 375 (1892).  
 HOUSTONIA SERPYLLIFOLIA Michx. (*Nuttall*).  
 HOUSTONIA PURPUREA Linn.  
 HOUSTONIA PURPUREA CALYCOSA A. Gray.  
 HOUSTONIA CILIOLATA Torr.  
 HOUSTONIA LONGIFOLIA Gærtn.

HOUSTONIA TENUIFOLIA Nutt.  
 CEPHALANTHUS OCCIDENTALIS Linn.  
 MITCHELLA REPENS Linn.  
 DIODIA TERES Walt.  
 GALIUM APARINE Linn.  
 GALIUM PILOSUM Ait.  
 GALIUM CIRCEZANS Michx.  
 GALIUM LANCEOLATUM Torr.  
 GALIUM LATIFOLIUM Michx.  
 GALIUM TRIFIDUM Linn.  
 GALIUM TINCTORIUM Linn. *G. trifidum latifolium* Torr.  
 GALIUM CONCINNUM T. & G.  
 GALIUM ASPRELLUM Michx.  
 GALIUM TRIFLORUM Michx.

## CAPRIFOLIACEÆ.

SAMBUCUS CANADENSIS Linn.  
 SAMBUCUS PUBENS Michx. *S. racemosa* Linn.  
 SAMBUCUS PUBENS ALBICOCCA Britt.  
 VIBURNUM ALNIFOLIUM Marsh. *V. lantanoides* Michx.  
 VIBURNUM ACERIFOLIUM Linn.  
 VIBURNUM DENTATUM Linn.  
 VIBURNUM NUDUM Linn.  
 VIBURNUM LENTAGO Linn.  
 VIBURNUM PRUNIFOLIUM Linn.  
 TRIOSTEUM PERFOLIATUM Linn.  
 SYMPHORICARPOS SYMPHORICARPOS (Linn.) MacM. *S. orbiculatus* Mœench.  
 LONICERA DIOICA Linn. *L. glauca* Hill.  
 LONICERA JAPONICA Thumb.  
 DIERVILLA DIERVILLA (Linn.) MacM. *Diervilla trifida* Mœench.

## VALERIANACEÆ.

VALERIANA PAUCIFLORA Michx.

## DIPSACEÆ.

DIPSACUS SYLVESTRIS Mill.

## CUCURBITACEÆ.

CUCURBITA OVIFERA Linn.  
 CITRULLUS VULGARIS Schrad.  
 CUCUMIS MELO Linn.  
 MICRAMPELIS LOBATA (Michx.) Greene. *M. echinata* Raf.  
 SICYOS ANGULATUS Linn.



CAMPANULACEÆ.

- LOBELIA CARDINALIS Linn.  
 LOBELIA SYPHILITICA Linn.  
 LOBELIA SYPHILITICA ALBIFLORA Britton.  
 LOBELIA PUBERULA Michx.  
 LOBELIA AMÆNA GLANDULIFERA A Gray.  
 LOBELIA LEPTOSTACHYS A. DC.  
 LOBELIA SPICATA Lam.  
 LOBELIA SPICATA PARVIFLORA A. Gray.  
 LOBELIA INFLATA Linn.  
 LOBELIA INFLATA SIMPLEX (Raf.) Millsp. Fl. W. Va. 398 (1892).  
 LEGOUZIA PERFOLIATA (Linn.) Britton. *Specularia perfoliata* A. DC.  
 CAMPANULA ROTUNDIFOLIA Linn.  
 CAMPANULA APARINOIDES Pursh.  
 CAMPANULA AMERICANA Linn.  
 CAMPANULA DIVARICATA Michx.

COMPOSITÆ.

- ELEPHANTOPUS CAROLINIANUS Willd.  
 ELEPHANTOPUS TOMENTOSUS Linn.  
 VERNONIA GIGANTEA (Walt.) Britton. *V. altissima* Nutt  
 VERNONIA NOVEBORACENSIS (Linn.) Willd.  
 VERNONIA GLAUCA (Linn.) Britton. *V. noveboracensis latifolia*. A. Gray.  
 EUPATORIUM PURPUREUM Linn.  
 EUPATORIUM MACULATUM AMÆNUM. (Pursh.) Britton. *E. purpureum amœnum* A. Gray.  
 EUPATORIUM HYSSOPIFOLIUM Linn.  
 EUPATORIUM PUBESCENS Muhl. *E. rotundifolium pubescens* B. S. P.  
 EUPATORIUM ALTISSIMUM Linn.  
 EUPATORIUM SESSILIFOLIUM Linn.  
 EUPATORIUM PERFOLIATUM Linn.  
 EUPATORIUM AGERATOIDES Linn. f.  
 EUPATORIUM AROMATICUM Linn.  
 EUPATORIUM CÆLESTINUM Linn.  
 LACINARIA SPICATA (Linn.) O. K.  
**Lacinaria scariosa squarrulosa** (Michx.) Small.  
 Greenbrier Co., White Sulphur Springs (*A. Brown*).  
 CHRYSOPSIS MARIANA (Linn.) Nutt.  
 SOLIDAGO FLEXICAULIS Linn. *S. latifolia* Linn.  
 SOLIDAGO CÆSIA Linn.  
 SOLIDAGO CURTISSII T. & G.  
 SOLIDAGO BICOLOR Linn.

- SOLIDAGO MONTICOLA T. & G.  
 SOLIDAGO PUBERULA Nutt.  
 SOLIDAGO SPECIOSA Nutt.  
 SOLIDAGO ODORA Ait.  
 SOLIDAGO RUGOSA Mill.  
 SOLIDAGO ULMIFOLIA Muhl.  
 SOLDAGO BOOTTH Hook.  
 SOLIDAGO ARGUTA Ait.  
 SOLIDAGO JUNCEA Ait.  
 SOLIDAGO JUNCEA SCABRELLA (T. & G.) A. Gray.  
 SOLIDAGO JUNCEA RAMOSA Porter & Britton.  
 SOLIDAGO SEROTINA Ait.  
 SOLIDAGO SEROTINA GIGANTEA (Ait.) A. Gray.  
 SOLIDAGO RUPESTRIS Raf.  
 SOLIDAGO CANADENSIS Linn.  
 SOLIDAGO NEMORALIS Ait.  
 EUTHAMIA GRAMINIFOLIA (Linn.) Nutt. *Solidago lanceolata* Linn. (*Nuttall*).  
 EUTHAMIA CAROLINIANA (Linn.) Greene. *Solidago Caroliniana* Linn.  
 BRACHYCHÆTA SPHACELATA (Raf.) Britton. *B. Cordata* T. & G.  
 SERICOCARPUS ASTEROIDES (Linn.) B. S. P.  
     Greenbrier Co., mountains about White Sulphur Springs, alt. 3,000 ft. (*A. Brown*).  
 ASTER DIVARICATUS Linn. *A. corymbosus* Ait.  
 ASTER MACROPHYLLUS Linn.  
 ASTER PATENS Ait.  
 ASTER PATENS PHLOGIFOLIUS (Muhl.) Nees.  
 ASTER LÆVIS Linn.  
 ASTER UNDULATUS Linn.  
 ASTER CORDIFOLIUS Linn.  
 ASTER LÓWRIEANUS Porter. *A. cordifolius lævigatus* Porter.  
 ASTER PURPURATUS Nees. *A. virgatus* Ell.  
 ASTER ERICOIDES Linn.  
 ASTER ERICOIDES DEPAUPERATUS Porter. *A. ericoides pusilus* A. Gray.  
 ASTER ERICOIDES PILOSUS (Willd.) Porter. *A. ericoides villosus* T. & G.  
 ASTER LATERIFLORUS (Linn.) Britton.  
 ASTER LATERIFLORUS HIRSUTICAULIS (Lindl.) Porter.  
 ASTER MULTIFLORUS Ait.  
 ASTER DUMOSUS Linn.  
 ASTER VIMINEUS Lam.  
 ASTER VIMINEUS FOLIOLOSUS (Ait.) A. Gray.  
 ASTER PANICULATUS Lam.  
 ASTER SALICIFOLIUS Ait.

- ASTER NOVI-BELGII Linn.  
 ASTER PRENANTHOIDES Muhl.  
 ASTER PUNICEUS Linn.  
 ASTER UMBELLATUS Mill.  
 ASTER INFIRMUS Michx.  
 ASTER ACUMINATUS Michx.  
 ASTER TENUIFOLIUS Linn.  
 ASTER LINARIIFOLIUS Linn.  
 ERIGERON CANADENSIS Linn.  
 ERIGERON ANNUUS (Linn.) Pers.  
 ERIGERON RAMOSUS (Walt.) B. S. P.  
 ERIGERON PULCHELLUS Michx.  
 ERIGERON PHILADELPHICUS Linn.  
 ANTENNARIA PLANTAGINIFOLIA (Linn.) Rich.  
 ANTENNARIA MARGARITACEA (Linn.) Hook. *Anaphalis margaritacea*  
     Benth & Hook.  
 GNAPHALIUM OBTUSIFOLIUM Linn.  
 GNAPHALIUM ULIGNOSUM Linn.  
 GNAPHALIUM PURPUREUM Linn.  
 INULA HELENIUM Linn.  
 POLYMNIA CANADENSIS Linn.  
 POLYMNIA CANADENSIS RADIATA A. Gray.  
 POLYMNIA UVEDALIA Linn.  
 SILPHIUM ASTERISCUS Linn.  
 SILPHIUM TRIFOLIATUM Linn.  
     Greenbrier Co., near White Sulphur Springs, alt. 3,000 ft.  
     (*A. Brown*).  
 CHRYSOGONUM VIRGINIANUM Linn.  
 PARTHENIUM INTEGRIFOLIUM Linn.  
 AMBROSIA TRIFIDA Linn.  
     Mineral Co., opposite Cumberland, Md. (*J. K. Small*).  
 AMBROSIA TRIFIDA INTEGRIFOLIA (Muhl.) T. & G.  
 AMBROSIA ARTEMISIÆFOLIA Linn.  
 XANTHIUM SPINOSUM Linn.  
     Mineral Co., opposite Cumberland, Md. (*J. K. Small*).  
 XANTHIUM STRUMARIUM Linn.  
 XANTHIUM CANADENSE Mill.  
 ECLIPTA ALBA (Linn.) Hassk.  
 HELIOPSIS SCABRA Dunal.  
 HELIOPSIS HELIANTHOIDES (Linn.) B. S. P. *H. laevis* Pers.  
 BRAUNERIA PURPUREA (Linn.) Britton. *Echinacea purpurea* Mœnch.  
 RUDBECKIA LACINIATA Linn.  
 RUDBECKIA LACINIATA HUMILIS A. Gray.

- RUDBECKIA FULGIDA Ait.  
 RUDBECKIA TRILOBA Linn.  
 RUDBECKIA HIRTA Linn.  
 RUDBECKIA SPECIOSA Wender.  
 HELIANTHUS LÆTIFLORUS Pers.  
 HELIANTHUS OCCIDENTALIS Riddell.  
 HELIANTHUS OCCIDENTALIS DOWELLIANUS T. & G.  
 HELIANTHUS TOMENTOSUS Michx.  
 HELIANTHUS GROSSE-SERRATUS Martens.  
 HELIANTHUS GIGANTEUS Linn.  
 HELIANTHUS LÆVIGATUS T. & G.  
 HELIANTHUS DORONICOIDES Lam.  
 HELIANTHUS MICROCEPHALUS T. & G. *H. parviflorus* Bernh.  
 HELIANTHUS DIVARICATUS Linn.  
 HELIANTHUS HIRSUTUS Raf.  
 HELIANTHUS STRUMOSUS Linn.  
 HELIANTHUS TRACHELIIFOLIUS Mill.  
 HELIANTHUS DECAPETALUS Linn.  
 VERBESINA OCCIDENTALIS (Linn.) Walt.  
 VERBESINA ALTERNIFOLIA (Linn.) Britton. *Ridania alternifolia* O. K.  
 COREOPSIS LANCEOLATA VILLOSA Michx.  
 COREOPSIS PUBESCENS Ell.  
 COREOPSIS AURICULATA Linn.  
 COREOPSIS MAJOR Walt. *C. senifolia* Michx.  
 COREOPSIS MAJOR CEMLERI (Ell.) Britton. *C. senifolia stellata* T. & G.  
 COREOPSIS TRIPTERIS Linn.  
 BIDENS FRONDOSA Linn.  
 BIDENS CONNATA Muhl. *B. connata comosa* A. Gray.  
 BIDENS LÆVIS (Linn.) B. S. P.  
 BIDENS BIPINNATA Linn.  
 BIDENS TRICHOSPERMA (Michx.) Britton. *Coreopsis trichosperma* Michx.  
 GALLINSOGA PARVIFLORA Cav.  
 HELENIUM AUTUMNALE Linn.  
 ANTHEMIS COTULA Linn.  
 ANTHEMIS ARVENSIS Linn.  
 ACHILLEA MILLEFOLIUM Linn.  
 CHRYSANTHEMUM LEUCANTHEMUM Linn.  
 MATRICARIA MATRICARIOIDES (Less.) Porter.  
 TANACETUM VULGARE Linn.  
 SENECIO VULGARIS Linn.  
 SENECIO AUREUS Linn.  
 SENECIO OBOVATUS Muhl. *S. aureus obovatus* T. & G.  
 SENECIO BALSAMITÆ Muhl. *S. aureus Balsamitæ* T. & G.

CACALIA SUAVEOLENS Linn.

CACALIA RENIFORMIS Muhl.

CACALIA ATRIPLICIFOLIA Linn.

ERECHTITES HIERACIFOLIA (Linn.) Raf.

ARCTIUM LAPPA Linn.

Mineral Co., opposite Cumberland, Md. (*J. K. Small*).

ARCTIUM MINUS Schk. *A. Lappa minus* A. Gray.

CARDUUS LANCEOLATUS Linn. *Cnicus lanceolatus* (Willd.)

CARDUUS ALTISSIMUS Linn. *Cnicus altissimus* Willd.

CARDUUS DISCOLOR (Muhl.) Nutt. *Cnicus discolor* A. Gray.

CARDUUS VIRGINIANUS Linn. *Cnicus Virginianus* Pursh.

CARDUUS MUTICUS (Michx.) Pursh. *Cnicus muticus* Pursh.

CARDUUS ODORATUS (Muhl.) Porter. *Cnicus odoratus* Muhl.

CARDUUS ARVENSIS (Linn.) Robs. *Cnicus arvensis* Hoffm.

ADOPOGON DANDELION (Linn.) O. K.

ADOPOGON VIRGINICUM (Linn) O. K. *A. amplexicaule* O. K.

CICHORIUM INTYBUS Linn.

HIERACIUM CANADENSE Michx.

HIERACIUM PANICULATUM Linn.

HIERACIUM VENOSUM Linn.

HIERACIUM SCABRUM Michx.

HIERACIUM GRONOVII Linn.

HIERACIUM LONGIPILUM Torr.

PRENANTHES ALTISSIMA Linn.

PRENANTHES ALBA Linn.

PRENANTHES SERPENTARIA Pursh. (*Nuttall*.)

TARAXACUM TARAXACUM (Linn) Karst. *T. officinale* Webb.

LACTUCA SCARIOLA Linn.

LACTUCA CANADENSIS Linn.

LACTUCA PULCHELLA (Pursh.) DC. *L. integrifolia* Nutt.

LACTUCA HIRSUTA Muhl.

LACTUCA SPICATA (Lam.) Hitch. *L. leucophæa* A. Gray.

LACTUCA VILLOSA Jacq.

LACTUCA FLORIDANA (Linn.) Gærtn.

SONCHUS OLERACEUS (Linn.)

SONCHUS ASPER (Linn.) All.

TRAGOPOGON PORRIFOLIUS Linn.

## Host Index of the Fungi.\*

ABIES BALSAMEA (L.) Miller.  
Peridermium Balsameum.

ABUTILON ABUTILON (L.) Rusby. (*A. Avicenne* Gærtm.)  
Macrosporium Abutilonis.

ACALYPHA VIRGINICA L.  
Cercospora Acalyphæ.

ACER *sp.*  
Daldinia concentrica.  
Helminthosporium persistens.  
Myxosporium seriatum.  
Trametes ambigua.

ACER NEGUNDO L. (*Negundo aceroides* Mœnch.)  
Ceganella Aceris.  
Diplodia atrata.  
Fusarium sarcochroum.  
Phoma negundinicola ramicola.  
Phyllosticta *sp.*  
Tubercularia vulgaris.

ACER PENNSYLVANICUM L.  
Botryodiplodia acerina.  
Cytispora exasperans.  
Phoma Pennsylvanica.  
Phoma samararum.  
Rhytisma punctatum.

ACER RUBRUM L.  
Botryosphæria quercuum.  
Bromella Ravenelii.  
Ceganella Aceris.  
Eutypella rugiella.  
Libertella acerina?  
Phyllosticta acericola.  
Sphæronæma acerinum.  
Steganosporium piriforme.  
Valsa etherialis.  
Valsa pauperata.

---

\* Species habiting fallen twigs, bark, branches, trunks, logs, stumps and leaves; soil, ground, sand and rocks; and excrementitious and decaying vegetable and animal substances, etc., etc., are not included in this index.

ACER SACCHARINUM L. (*A. dasycarpum* Ehrh.)

- Diaporthe Aceris.
- Fomes applanatus.
- Libertella acerina?
- Phyllosticta acericola.
- Scleroderris pallidula.
- Stagonospora collapsa.
- Valsa ceratophora.

ACER SACCHARUM Marsh. (*Acer saccharinum* Wang.)

- Hypoxylon fuscum.
- Pilacre Petersii.

ACTÆA ALBA (L.) Mill.

- Urocystis Anemones.

*Actinomeris squarrosa* see VERBESINA ALTERNIFOLIA.

AESCULUS OCTANDRA HYBRIDA (DC.) Sarg. (*Ae. octandra purpurascens* Gr.)

- Phyllosticta sphæropsidea.

AGARICUS *sp.*

- Nyctalis asterophora.
- Sporodinia Aspergillus.

AGRIMONIA STRIATA Michx. (*A. Eupatoria* L.)

- Sphærotheca Humuli.
- Uredo Agrimonix.

ALNUS RUGOSA (Ehrh.) Koch. (*A. serrulata* Willd.)

- Cryptospora femoralis.
- Diatrypella verruciformis.
- Eutypella stellulata.
- Glæosporus tremellosus.
- Glæosporium Alni. *sp. nov.*
- Glæosporium cylindrospermum.
- Hypoxylon fuscum.
- Phlebia radiata.
- Valsa femoralis.

AMALANCHIER CANADENSIS (L.) Medic.

- Dimerosporium Collinsii.

AMARANTHUS RETROFLEXUS L.

- Cystopus Bliti.

*Ampelopsis quinquefolia*. see PARTHENOCISSUS QUINQUEFOLIA.

AMYGDALUS PERSICA L. (*Prunus Persica* Bth. & Hook.)

- Cytispora leucostoma.
- Cytispora Persicæ.
- Exoascus deformans.
- Monilia fructigena.
- Sterigmatocystis nigra.

Streptothryx atra.  
Trichothecium roseum.  
Valsa leucostoma.

ANEMONE VIRGINIANA L.  
Puccinia Anemones-Virginiana.

*Andromeda ligustrina* see XOLISMA LIGUSTRINA.

ARALIA SPINOSA L.  
Botryosphæria Araliæ.  
Diaporthe Araliæ.  
Eutypella densissima.  
Haplosporella Araliæ.  
Hypoderma commune.  
Hypoxyton coccineum.  
Lecanidion atratum.  
Leptothyrium vulgare.  
Macrosporium commune.  
Nectria cinnabarina.  
Nematella nucleata.  
Phoma melaleuca.  
Phyllosticta Araliæ.  
Physarum sinuosum.  
Stagonospora petiolorum.  
Tubercularia vulgaris.  
Valsa ambiens.  
Vermicularia petiolorum.

ARCTIUM LAPPA L.  
Phyllosticta Lappæ.  
Vermicularia Arctii.

ARISÆMA TRIPHYLLUM (L.) Torr.  
Uromyces Caladii.

ASCLEPIAS SYRIACA L. (*A. cornuti* Dec.)  
Fusarium roseum.  
Macrosporium asclepiadeum.  
Phoma asclepiadea *sp. nov.*  
Phoma *sp.*  
Uromyces Howei.

ASIMINA TRILOBA (L.) Dunal.  
Anthostoma microëcium.  
Cytispora carphosperma.  
Dinemasporium hispidulum.  
Dothiorella Asiminæ *sp. nov.*  
Macrosporium olivaceum *sp. nov.*  
Mucor Mucedo.  
Phyllosticta Asiminæ.  
Rosellina microëcium.  
Sphæropsis Asiminæ *sp. nov.*  
Sphæropsis Asiminæ fructigena *var. nov.*



Tubercularia Asiminæ *sp. nov.*

Ustilago *sp.*

Valsa ambiens.

ASPARAGUS OFFICINALIS L.

Fusarium *sp.*

Gibberella Saubinettii.

Macrosporium commune.

Tilmadoche mutans.

Torula herbarum.

Vermicularia liliacearum.

*Aspidium spinulosum* see DRYOPTERIS SPINULOSA.

ASTER *sp.*

Thelephora sebacea.

ASTER CORDIFOLIUS L.

Coleosporium Sonchi.

Puccinia Asteris.

Septoria atro-purpurea.

ASTER INFIRMUS Mich.

Fusicladium *sp.*

AVENA SATIVA L.

Epicoccum neglectum.

Ustilago Avenæ lævis.

Ustilago segetum.

AZALEA VISCOSA L.

Sphærographum histricinum.

Pestalozzia Guepini.

BENZOIN BENZOIN (L.) Coulter. (*Lindera Benzoin* Meisn.)

Camorosporum Linderæ *sp. nov.*

Diaporthe sociata.

Diplodia Linderæ.

Phyllosticta lindericola.

Sphæropsis Linderæ.

Valsa Linderæ.

BETA VULGARIS L.

Cercospora beticola.

BETULA *sp.*

Corticium Petersii,

Hypoxyton atroviride?

Hypoxyton multiforme.

Polyporus pubescens.

BETULA LENTA L.

Gelatinosporum betulinum.

Hymenochæte corrugata.

Hymenochæte unicolor.

Polystictus hirsutus.

Poria tomento-cincta.  
 Septoria microsperma.  
 Stereum complicatum.  
 Stereum sericeum.  
 Stereum versicolor.

BETULA LUTEA Michx. f.

Discosia Artoceras.  
 Fomes fomentarius.  
 Lentinus strigosus.

BETULA NIGRA L.

Aposphæria pezizoides.  
 Cytispora betulina.  
 Glæosporium betularum.  
 Melanconium bicolor.  
 Steganosporium muricatum.

BOEHMERIA CYLINDRICA (L.) Willd.

Cercospora Boehmeriæ.

BOLETUS *sp.*

Sporodinia Aspergillus.

BRASSICA NIGRA (L.) Koch.

Cystopus candidus.  
 Septoria Brassicæ *sp. nov.*

BRASSICA OLERACEA L.

Botrytis vulgaris.

CARDUUS LANCEOLATUS L.

Puccinia Hieracii.  
 Puccinia suaveolens.

CALOPTENUS FEMUR RUBRUM (*grasshopper*).

Bacterium *sp.*  
 Empusa Grylli.

CAREX *sp.*

Puccinia Caricis.

CAREX FRASERI And.

Botrytis torta.  
 Epidochium melanochlorum?  
 Stachylidium caricinum.  
 Trichægum nodulosum.  
 Zygoesmus graminicola.

CARPINUS CAROLINIANA Walt.

Cytispora leucosperma.  
 Glæosporium Robergei.  
 Radulum orbiculare.  
 Thyrsidium hedericolum Carpini.  
 Xylaria flabelliformis.

*Carya sp.*, see HIKORIA *sp.*

CASSIA MARILANDIA L.

Vermicularia Dematium.

CASTANEA DENTATA (Marsh.) Sudw. (*C. sativa Americana* Sarg.)

Coryneum pustulatum.

Cryptospora cinctula.

Diatrype Stigma.

Discosia Artoceras.

Glonium parvulum.

Lenzites corrugata.

Leptothyrium Castanæ.

Melanconis modonia.

Microsphæria Alni.

Mucronoporus Gilbus.

Peniophora quercina.

Phyllactinia suffulta.

Septoria ochroleuca.

Sphærella maculiformis.

Steganosporium Castanæ.

Stereum versiforme.

Valsa coronata.

CASTANEA PUMILA Mill.

Phyllosticta Castanæ.

CATALPA CATALPA (L.) Karst. (*Catalpa Bignonioides* Walt.)

Epicoccum neglectum.

Microsphæria elevata.

Phyllosticta Catalpæ.

CATERPILLARS.

Empusa Grylli.

CAULOPHYLLUM THALICTROIDES (L.) Michx.

Cercospora Caulophylli.

Diaporthe Gladioli?

Vermicularia compacta.

CELASTRUS SCANDENS L.

Botryosphæria Hibiscii *var.*

Phyllosticta Celastri *sp. nov.*

Ramularia Celastri.

Sphæropsis celastrina.

Tubercularia Celastri.

CELTIS OCCIDENTALIS L.

Cytispora Celtidis.

Haplosporella Celtidis.

Macrosporium antennæforme.

Phyllosticta Celtidis.

Tubercularia hamata.

CEPHALANTHUS OCCIDENTALIS L.

Discosia Artoceras.

## CERCIS CANADENSIS L.

- Cytispora *sp.*
- Diplodia Cercidis.
- Glæosporium *sp.*
- Phyllosticta cercidicola *sp. nov.*
- Sarcinella heterospora.

## CHENOPODIUM ALBUM VIRIDE (L.) Moq.

- Cercospora dubia.

## CHENOPODIUM ANTHELMINTICUM L.

- Cercospora anthelmintica.
- Pyrenophora calvescens.

## CHIONANTHUS VIRGINICA L.

- Botryosphæria pyriospora.
- Cercospora Chionanthi *sp. nov.*
- Cytispora *sp.*
- Dermatea Chionanthi.
- Phyllosticta Chionanthi.
- Stagonospora *sp.*
- Valsa Chionanthi.

## CICUTA MACULATA L.

- Botrytis oleracea.
- Macrosporium commune.

## CIMICIFUGA RACEMOSA (L.) Nutt.

- Leptosphæria Ogilviensis.

## CLEMATIS VIRGINIANA L.

- Ascochyta clematidina.

## CLINTONIA UMBELLULATA (Michx.) Torr

- Vermicularia liliacearum.

*Gnicus lanceolatus* Willd see CARDUUS LANCEOLATUS L.CONVOLVULUS REPENS L. (*Calystegia sepium repens* Gray.)

- Puccinia Convolvuli.
- Puccinia Convolvuli *var.*

## CORNUS ALTERNIFOLIA L.f.

- Septoria cornicola.

## CORNUS FLORIDA L.

- Aschersonia *sp.*
- Dædalia confragosa.
- Discosia Artoceras.
- Phyllosticta cornicola.
- Phyllosticta globifera *sp. nov.*
- Septoria Corni-Maris.
- Sporodesmium toruloides.
- Tremella virens.
- Valsa ambiens.

CORYLUS AMERICANA Walt.

Gnomoniella Coryli.

CRATEGUS *sp.*

Aspergillus glaucus.

Cylindrosporium Cratægi.

Gymnosporangium clavariiforme (*Accid.*)

CRATEGUS OXYACANTHA L.

Podosphæria Oxyacanthæ.

CUNILA ORIGANOIDES (L.) Britton. (*C. Mariana* L.)

Puccinia Menthæ.

*Cuphœa petiolata* see PARSONSIA PETIOLATA.

DATURA STRAMONIUM L.

Macrosporium *sp.*

DENDROCTONUS FRONTALIS (*Insect.*)

Cylindrocolla Dendroctoni.

DENTARIA DIPHYLLA Michx.

Cystopus candidus.

*Desmodium paniculatum* see MEIBOMIA PANICULATA.

*Desmodium canescens* see MEIBOMIA CANESCENS.

DIANTHUS AMERICANA L.

Cercospora Diantheræ.

*Diarrhena Americana* see KORYCARPUS DIANDRUS.

DIOSCOREA VILLOSA L.

Phyllosticta Dioscoreæ.

Leptothyrium vulgare.

DIOSPYROS VIRGINIANA L.

Fusarium roseum *var.*

Macrosporium *sp.*

Podosphæria Oxyacanthæ.

Valsa Diospyri.

DISPORUM LANUGINOSUM (L.) Morong.

Discosia maculicola.

DRYOPTERIS SPINULOSA (Retz.) Kuntze. (*Aspidium spinulosum* Sw.)

Leocarpus fragilis.

Leptostromella filicina.

ELYMUS CANADENSIS GLAUCIFOLIUS (Willd.) Torr.

Phyllachora graminis.

EPIGÆA REPENS L.

Discosia Artoceras.

Erysiphe Vaccinii.

## EQUISETUM ARVENSE L.

Septosporium Equiseti.

## ERECTITES HIERACIFOLIA (L.) Raf.

Sphærotheca Castagnei.

## EUPATORIUM AGERATOIDES L.

Cercospora ageratoides.

Erysiphe communis.

Puccinia tenuis.

## EUPATORIUM CÆLESTINUM L.

Puccinia conoclinii.

## EUPATORIUM PURPUREUM L.

Oidium erysiphoides.

Erysiphe Cichoracearum.

FAGUS ATROPUNICEA (Marsh.) Sudw. (*F. ferruginea* Ait.)

Coryne urnalis.

Cryptosporella compta.

Horomyces fragiformis.

Hydnum coralloides.

Hydnum Erinaceus.

Lachnea scutellata.

Libertella faginea.

Microsphæria erineophila.

Polystictus hirsutus.

Scorias spongiosa.

FRAGARIA *cult.*

Phoma obscurans.

Ramularia Tulasnei.

Septoria aciculosa.

FRAXINUS *sp.*

Aposphæria pezizoides.

## FRAXINUS AMERICANA L.

Cucurbitaria Fraxini.

Diplodia infuscans.

Diplodia inquinans.

Fomes applanatus.

Hemiarcyria rubiformis.

Hypholoma sublateritium.

Sphærographum Fraxini.

Sphæroopsis phomatella.

FRAXINUS PENNSYLVANICA Mart. (*F. pubescens* Lam.)

Cytispora ceratophora.

Discosia Artoceras.

Glœosporium aridum.

## GALAX APHYLLA L.

Asterina Leemingii.

Dimerosporium Galactis.

Phyllosticta Galacis.

GALIMUM CIRCÆZANS Michx.

Septoria pilostega.

GAULTHERIA PROCUMBENS L.

Hypocrea tenerrima.

Sphærella Gaultheriæ.

GENTIANA ANDREWSII Griesb.

Phyllosticta gentianicola.

GEUM CANADENSE Jacq. (*G. album* Gm.)

Pestalozzia *sp.*

Phyllosticta *sp.*

*Halesia tetraptera* see MOHRODENDRON CAROLINUM.

HAMAMELIS VIRGINICA L.

Corticium leve.

Gonatobotryum maculicolum.

Lecanidion Hamamelidis.

Leptosphæria Doliolum.

Phyllosticta Hamamelidis.

Phyllosticta sphæropsidea.

HELIANTHUS DECAPETALUS L.

Cylindrocolla flagellaris *sp. nov.*

Puccinia Helianthi.

Septoria Helianthi.

HEUCHERA AMERICANA L.

Septoria Saxifragæ.

Vermicularia Dematium.

HICORIA *sp.*

Corticium lilacino-fuscum.

Scoriomyces Cragini.

Sphæropsis Caryæ.

Tremella intumescens.

HICORIA MICROCARPA (Nutt.) Britton. (*Carya microcarpa*.)

Phyllosticta caryiogena.

HICORIA MINIMA (Marsh.) Britton. (*Carya amara*.)

Valsa ceratophora.

HICORIA OVATA (Mill.) Britton. (*Carya alba*.)

Corticium subgiganteum.

Cytispora caryiogena.

Dædalia unicolor.

Diplodia caryiogena.

Melanoconium pallidum.

Myxosporium luteum.

Phyllosticta caryiogena.

Polyporus adustus.

Tilmadoche gyrocephala.

Trichoderma lignorum.

## HOLCUS LANATUS L.

Puccinia coronata.

## HOUSTONIA CÆRULEA L.

Æcidium Houstonianum.

## HOUSTONIA LONGIFOLIA Gærtm.

Æcidium Houstonianum.

HYDNUM *sp.*

Pencilium glaucum.

Stremonites fusca.

## HYDRANGÆA ARBORESCENS L.

Diaporthe Hydrangæ *sp. nov.*

Hypoderma commune.

Leptosphæria vagabunda.

Tilmadoche gyrocephala.

HYMENOCETUM *sp.*

Hypomyces rosellus.

## HYPERICUM MUTILUM L.

Uromyces Hyperici.

## HYPOXYLON COCCINEUM Bull.

Lasiosphæria ovina.

## HYPOXYLON RUBIGINOSUM.

Isaria Virginiensis.

## HYSTERIX HYSTERIX (L.) Millsp.

Phyllachora graminis.

## ILEX OPACA Ait.

Diaporthe cercophora.

Diplodia ilicicola.

Fusicoccum ilicinum.

Heliscus Lugdunensis.

Hypoxylon perforatum.

Leptothyrium foraminulatum.

Leptothyrium *sp.*

Phoma ilicicola.

Phyllosticta opaca *sp. nov.*

Physalospora Ilicis.

Pilacre Petersii.

## ILEX VERTICILLATA (L.) Gray.

Phyllosticta Hynaldi.

Rhytisma Prini.

Trichothecium roseum.

IMPATIENS BIFLORA Walt. (*I. fulva.*)

Septoria Nolintangere.

Æcidium Impatientis.



IPOMŒA PANDURATA (L.) Meyer.

- Botrytis vulgaris.
- Coleosporium Ipomœæ
- Pestalozzia funerea.
- Phlyctæna Ipomœæ.
- Phyllosticta Ipomœæ.
- Sphæropsis Ipomœæ.

IRIS CRISTATA Ait.

- Vermicularia liliacearum.

JUGLANS CINEREA L.

- Dasyscypha virginea.
- Diplodia Juglandis.
- Exidia glandulosa.
- Glonium simulans.
- Lasiosphæria ovina.
- Marsonia Juglandis.
- Melanconium oblongum.
- Mucronoporus ferruginosus.
- Orbilia coccinella.
- Propolis Faginea.
- Pseudohelotium fibrisedum.
- Sphæronæma infuscans *sp. nov.*
- Tapesia fusca.
- Tapesia sanguinea.
- Valsaria exasperans.

JUNIPERUS VIRGINIANA L.

- Gymnosporangium juniperinum (*Teleuto.*)
- Gymnosporangium macropus.

KALMIA LATIFOLIA L.

- Cercospora Kalmiæ.
- Dacryomyces minor.
- Hymenochæte unicolor.
- Hypocreopsis riccoidea.
- Hysterium Kalmiæ?
- Poria sinuosa.
- Rhabdospora Kalmianum.
- Septoria kalmiæcola.

KORYCARPUS DIANDRUS (Michx.) Kuntze. (*Diarrhena Americana.*)

- Phyllachora Caricis.

LACHNOCLADIUM SEMIVESTITUM.

- Aspergillus flavus.
- Aspergillus glaucus oblongisporus *var. nov.*

LACTUCA CANADENSIS L.

- Botryosporium pulchrum.
- Bremia Lactuæ.

LACTUCA HIRSUTA Muhl.

- Bremia Lactuæ.

## LESPEDEZA VIOLACEA (L.) Pers.

Uromyces Lespedezæ.

*Lindera Benzoin* see BENZOIN BENZOIN.

## LIQUIDAMBAR STYRACIFLUA L.

Aposphæria pezizoides.

Cytispora *sp.*

Grandinia crustosa.

Lophiotrema Nucula.

Monilia aureo-fulva.

Propolis faginea.

Trichia proximella.

## LIRIODENDRON TULIPIFERA L.

Dendroochium rubellum microsporum.

Dothiorella minor *sp. nov.*

Diplodia Liriodendri.

Eurotium herbariorum.

Erysiphe Liriodendri.

Leptothyrium Liriodendri.

Orbilina rubella.

Propolis faginea.

Phoma mixta.

Phyllosticta liriodendrica.

Phyllosticta macrospora.

Polystictus hirsutus.

Poria corticola.

Poria Tulipiferæ

Tubercularia vulgaris.

Valsa albopuncta *sp. nov.*

## LOBELIA CARDINALIS L.

Septoria Lobeliæ.

## LYCOPERSICUM ESCULENTUM L.

Fusarium Solani.

Macrosporium Tomato.

## MAGNOLIA ACUMINATA L.

Dendroochium rubellum microsporum.

Massaria Magnoliæ *sp. nov.*

Phyllosticta Cookei.

Sphæronæma Magnoliæ.

## MAGNOLIA FRASERI Walt.

Bombardia fasciculata.

Botrytis vulgare.

Calocera viscosa.

Cladosporium *sp.*

Collybia conigenoides.

Corticium leptaleum.

Corticium scutellare.

Diatrype platystoma.

Diatrypella verruciformis.  
 Discosia Artoceras.  
 Exidia glandulosa.  
 Helminthosporium macrocarpon.  
 Helminthosporium septemseptatum.  
 Hemiarcyria stipata.  
 Holwaya ophiobolus.  
 Hymenochæte corrugata.  
 Hypoxylon Howeanum.  
 Hypoxylon Nuttallii *sp. nov.*  
 Hypoxylon perforatum.  
 Irpex obliquus.  
 Kneiffia setigera.  
 Lachnella Virginica.  
 Leptothyrium petiolorum *var.*  
 Lycogala epidendrum.  
 Melanomma Pulvis-pyrius.  
 Nectria coccinea.  
 Nectria ditissima.  
 Nematella nucleata.  
 Orbilia Xanthostigma.  
 Peniophora cinerea.  
 Phlebia merismoides.  
 Phyllactinia suffulta.  
 Phoma pedunculi.  
 Physarum psittacinum.  
 Pilacre Petersii.  
 Polyporus *sp.*  
 Polyporus brunalis.  
 Poria ferruginosa.  
 Poria spissa.  
 Poria tomento-cincta.  
 Radulum Magnoliæ.  
 Stagonospora pedunculi.  
 Stilbum magnum.  
 Tapesia sanguinea.  
 Tilmadoche nutans.  
 Tilmadoche viridis.  
 Vermicularia subeffigurata.  
 Xylaria corniformis.

MEIBOMIA CANESCENS (L.) Kuntze. (*Desmodium canescens.*)  
 Uromyces Hedysari-paniculati.

MEIBOMIA PANICULATA (L.) Kuntze. (*Desmodium paniculatum.*)  
 Uromyces Hedysari-paniculati.

MENISPERMUM CANADENSE L.  
 Cercospora Menispermi.  
 Cladosporium herbarum.  
 Sphæroopsis Menispermi.

MOHRODENDRON CAROLINUM (L.) Britton. (*Halesia tetrap-  
tera.*)

Cytispora Halesiæ.  
Diaporthe Halesiæ.  
Diaporthe Tetrapteræ.  
Sirococcus Halesiæ.

MORUS RUBRA L.

Tubercularia *sp.*

MOSS.

Hypocrea tenerrima.  
Lycoperdon pedicellatum.  
Physarum citrinum.

*Nasturtium Armoracia* (L.) Fr. see RORIPA ARMORACIA.

NECTRIA VERRUCOSA.

Tubercularia *sp.*

*Negundo aceroides* see ACER NEGUNDO.

NYSSA AQUATICA L. (*N. sylvatica.*)

Phoma Nyssocarpa.  
Sphærella nyssæcola.  
Valsa Nyssæ.  
Valsa præstans.

*Ænothera biennis* see next.

ONAGRA BIENNIS (L.) Scop. (*Ænothera biennis.*)

Cercospora Ænotheræ.  
Leptothyrium Vulgare.  
Phlyctæna arcuata.  
Phlyctæna vagabunda.  
Phoma herbarum.  
Sphæronæma corneum.  
Sphæropsis Ænotheræ *sp. nov.*  
Vermicularia *sp.*

OPULASTER OPULIFOLIUS (L.) Kuntze. (*Physocarpus opuli-  
folius.*)

Botryosphæria Quercuum.  
Didymella Physocarpi.  
Pestalozzia Jefferisii.  
Phoma *sp.*  
Phoma leucostoma.  
Phoma Spirææ.  
Sphæronæma Physocarpi.  
Sphæropsis Physocarpi.  
Sporodesmium moriforme.  
Stagonospora Physocarpi.

ORANGE (*decaying.*)

Oospora fasciculata.

OSMORRHIZA CLAYTONII. (Michx.) B.S.P.

Puccinia Pimpinellæ.

OSTRYA VIRGINIANA (Mill.) Willd.

Thelephora pedicellata.

Corticium Oakesii.

OXALIS STRICTA L. (*O. corniculata stricta.*)

Microsphæria Russellii.

OXYDENDRON ARBOREUM (L.) DC.

Discosia maculicola.

Phyllosticta Oxydendri *sp. nov.*

Scleroderris pallidula.

Tympanis Oxydendri.

PARTHENOCISSUS QUINQUEFOLIA (L.) Planch. (*Ampelopsis quinquefolia.*)

Botryosphæria Quercuum.

Cercospora Ampelopsidis.

Coniothyrium Fuckelii.

Periconia pycnospora.

Phyllosticta Ampelopsidis.

Sphæropsis Ampelopsidis.

Uncinula Ampelopsidis.

PESTALOZZIA GUEPINI.

Dermatea lobata.

PHASEOLUS VULGARIS L.

Cercospora columnaris.

Colletotrichum Lindemuthianum.

Uromyces appendiculatus.

PHLOX AMCENA Sims.

Cercospora omphacodes.

Vermicularia phlogina.

PHRYMA LEPTOSTACHYA L.

Septoria Leptostachyæ.

*Physocarpus opulifolius* see OPULASTER OPULIFOLIUS.

PHYLLOSTICTA CARYIOGENA.

Discosia rugulosa.

PHYTOLACCA DECANDRA L.

Diaporthe aculeata.

Leptosphæria clavigera.

Macrosporium caudatum.

Periconia pycnospora.

Phlyctæna septorioides.

Phlyctæna vagabunda.

Phoma Phytolacçæ.

PICEA MARIANA (Mill.) BSP.

Fomes carneus.

Lenzites sepiaria.  
 Polystictus versicolor.  
 Scleroderma vulgare.

PINUS ECHINATA Mill. (*P. mitis.*)  
 Coleosporium Senecionis (*Æcid.*)  
 Hyphoderma Desmazieri.

PINUS RIGIDA Mill.  
 Amphispæria pinicola.  
 Dasycypha Ellisiana.  
 Lophodermium pinastri.  
 Pleosphæria corticola.  
 Stictis fimbriata.

PINUS STROBUS Linn.  
 Dacryomyces deliquescens.

PINUS VIRGINIANA Mill. (*P. inops.*)  
 Amphispæria pinicola.  
 Botryosphæria Quercuum.  
 Corticium scutellare.  
 Cytispora Curreyi.  
 Fomes volvatus.  
 Mollisia pinastri.  
 Polystictus abietinus.  
 Poria Xantha.  
 Rosellina abietina trichota.  
 Speira minor.  
 Stictis fimbriata.  
 Stereum Pini.  
 Tremella aurantia.

PLATANUS OCCIDENTALIS L.  
 Aposphæria pezizoides.  
 Cytispora Platani.  
 Dacryomyces Syringæ.  
 Dendrophoma Therryana.  
 Eutypella Platani.  
 Glæosporium nervisequum.  
 Helminthosporium macrocarpon.  
 Hendersonia Desmazieri.  
 Myxosporium platanicolum.  
 Phoma scabra.  
 Rosellinia millegrana.  
 Stilbum flavipes.

POA PRATENSIS L.  
 Erysiphe graminis.  
 Oidium monilioides.

PODOPHYLLUM PELTATUM L.  
 Puccinia Podophylli.

POLYGONATUM BIFLORUM (Walt.) Ell.

Cladosporium herbarum.

POLYGONUM AVICULARE L.

Cercospora avicularis.

POLYGONUM ERECTUM L.

Uromyces Polygoni.

POLYGONUM HYDROPIPER L.

Septoria Polygonorum.

POLYMNIA UVEDALIA L.

Helotium herbarum.

Leptosphaeria Doliolum.

Macrosporium *sp.*

Periconia pycnospora.

Pleurotus cyphelliformis.

Septoria Polymnæ.

POLYPORUS PERGAMENUS.

Calicium tigillare.

Cephalothecium roseum.

POLYPORUS VARIUS Fr.

Cladosporium epimyces.

Pencilium glaucum.

POLYSTICTUS VERSICOLOR.

Dactylium dendroides.

PORIA SPISSA.

Orbilina vinosa.

PORTULACCA OLERACEA L.

Cystopus Portulacæ.

POTENTILLA CANADENSIS L.

Phragmidium Potentillæ (*Uredo.*)

POTENTILLA MONSPELIENSIS L. (*P. Norvegica.*)

Mollisia Dehnii.

PRENANTHES SERPENTARIA Pursh.

Septoria Nabali.

PRUNUS *cult.* "Cherry."

Libertella faginea.

Monilia fructigena.

Plowrightia morbosa.

Podosphaeria Oxyacanthæ.

Podosphaeria tridactyla.

Polystictus cinnabarinus.

Volutella ciliata.

PRUNUS DOMESTICA L.

Cytispora leucostoma.

Exoascus Pruni.  
 Monilia fructigena.  
 Plowrightia morbosa.

*Prunus Persica* see AMYGDALUS PERSICA.

PRUNUS SEROTINA Ehrh.

Merulius ambiguus.  
 Polyporus nidulans.  
 Poria vaporaria.  
 Sphæropsis cerasina.

PYRUS COMMUNIS L.

Entomosporium maculatum.  
 Micrococcus amylovorus.

PYRUS CORONARIA L.

Gymnosporangium macropus (*Æcid.*)  
 Plowrightia morbosa

PYRUS MALUS L.

Fusicladium dendriticum,  
 Phyllosticta Pirina.  
 Polystictus hirsutus.  
 Schizophyllum commune.

QUERCUS *sp.*

Asterostoma cervicolor.  
 Botryosphæria Quercuum.  
 Chlorosplenium Schweinitzii.  
 Cryptospora trichospora.  
 Guepinia spathularia.  
 Hemiarcyria clavata.  
 Hypoxylon atroviride?  
 Hypoxylon stigmatum.  
 Nummularia punctulata.  
 Rosellinia corticium.  
 Stereum pustulosum.  
 Stereum versicolor.

QUERCUS ALBA L.

Bactridium flavum.  
 Coryne urnalis.  
 Fomes applanatus.  
 Hemiarcyria rubiformis.  
 Hemiarcyria clavata.  
 Lachnella erinaceus.  
 Lachnea scutellata.  
 Marsonia Martini.  
 Merulius tremellosus.  
 Merulius rubellus.  
 Phyllosticta phomiformis.  
 Polystictus versicolor.  
 Schizophyllum commune.  
 Stereum sulphuratum.



QUERCUS PALUSTRIS DuRoi.

Erysiphe quercina.  
 Fomes applanatus.  
 Lycoperdon pyriforme.  
 Trichopeziza capitata.

QUERCUS PRINUS L.

Botryodiplodia *sp.*  
 Cyathicula quisquillaris.  
 Marsonia Martini.  
 Phyllosticta phomiformis.  
 Phyllosticta Quercus-Prini.  
 Polyscytalum sericeum.  
 Pseudovalsa sigmoidea.  
 Stereum complicatum.

QUERCUS RUBRA L.

Fomes applanatus.  
 Leptothyrium dryinum.

QUERCUS VELUTINA Lam. (*Q. tinctoria*.)

Marsonia Martini.

RESEDA ODORATA L.

Cladosporium herbarum fasciculare.  
 Pyrenopeziza lacerta.

RHODODENDRON CATAWBIENSE Michx.

Pestalozzia Guepini.

RHODODENDRON MAXIMUM L.

Exobasidium Rhododendri.  
 Grandinia crustosa.  
 Heliotium castaneum.  
 Hypocrea Virginiensis.  
 Hypoxylon colliculosum.  
 Lophodermium Rhododendri.  
 Pestalozzia Guepini.  
 Sporocybe Azaleæ.  
 Stereum triste.  
 Tremella frondosa.

RHUS COPALLINA L.

Aponectria inaurata.  
 Calospora aculeans.  
 Polyporus pocula.  
 Sporocybe Rhois.  
 Stereum sericeum.

RHUS HIRTA (L.) Sudw. (*R. typhina*)

Calospora aculeans.  
 Calospora Rhoina.  
 Cladosporium herbarum.  
 Cytispora Rhoina.  
 Cytispora Rhois-hirtæ *sp. nov.*

Irpex lacteus.  
 Myxosporium Rhois.  
 Polyporus adustus.  
 Sphæropsis Sumachi.  
 Sporocybe Rhois.  
 Torula dimidiata.  
 Tubercularia vulgaris.

RHUS RADICANS L. (*R. toxicodendron*.)

Cylindrosporium Toxicodendri.  
 Macrosporium commune.  
 Pestalozzia Toxica.  
 Phyllosticta roicola.  
 Uromyces Terebinthi.  
 Vermicularia Toxica *sp. nov.*

RIBES *cult.*

Phyllosticta Ribis *sp. nov.*

ROBINIA PSEUDACACIA L.

Cladosporium epiphyllum.  
 Cladosporium nigrellum.  
 Cucurbitaria elongata.  
 Cytispora coccinea.  
 Cytispora orthospora.  
 Diaporthe oncostoma.  
 Dothiorella glandulosa.  
 Epicoccum Durianum.  
 Eutypella stellulata.  
 Fomes rimosus.  
 Hypoxylon rubiginosum.  
 Libertella *sp.*  
 Nectria ditissima.  
 Rosellinia subiculata.  
 Tubercularia vulgaris.  
 Vermicularia petiolorum.

RORIPA ARMORACIA (L.) Hitch. (*Nasturtium Armoracia*.)

Cercospora Armoraciæ.

ROSA *cult.*

Actinonema Rosæ.  
 Oidium leucoconium.

ROSA HUMILIS Marsh.

Phragmidium subcorticium.  
 Phyllosticta Rosæ.

RUBUS BAILEYANUS Britt. (*R. villosus humifusus*.)

Hypoderma virgultorum.  
 Scleroderris rhabbarbarina.

RUBUS CANADENSIS L.

Cercospora Rubi.

*Cercospora septorioides* *sp. nov.*  
*Phyllosticta* *sp.*  
*Septoria* Rubi.

RUBUS *culi*.

*Botryosphæria Quercuum* *var.*  
*Diplodia* Rubi.  
*Hainesia* Rubi.

RUBUS HISPIDUS L.

*Uredo* (*Ceoma*) *nitens*.

RUBUS ODORATUS L.

*Diaporthe rostellata*.  
*Oidium erysiphoides*.  
*Pericornia pycnospora*.  
*Pyrenopeziza lacerata*.  
*Vermicularia compacta*.

RUBUS STRIGOSUS Michx.

*Glæosporium rubicolum* *sp. nov.*  
*Glæosporium venetum*

RUBUS VILLOSOS Ait.

*Diaporthe obscura*.  
*Teichospora nitida*.  
*Uredo* (*Ceoma*) *nitens*.

RUMEX ACETOSELLA L.

*Coniosporium harknessioides*.

RUMEX OBTUSIFOLIUS L.

*Glæosporium Rumicis* *sp. nov.*  
*Macrosporium commune*.

SALIX NIGRA Marsh.

*Corticium salicinum*.  
*Cytispora Salicis*.  
*Diplodia Salicina*.  
*Melampsora farinosa*.  
*Trimmatostroma Americana*.

SALIX NIGRA FALCATA Torr.

*Aposphæria pezizoides*.  
*Hendersonia Lirella*.  
*Lophidium compressum*.

SAMBUCUS CANADENSIS L.

*Cercospora despazeoides*.  
*Corticium scutellare*.  
*Microsphæria Grossulariæ*.  
*Phyllosticta Sambuci*.

SAMBUCUS PUBENS Michx. (*S. racemosus*.)

*Tubercularia Sambuci*.  
*Tubercularia vulgaris*.

## SANGUINARIA CANADENSIS L.

Glœosporium Sanguinariæ.

Phyllosticta Sanguinariæ

## SAPONARIA OFFICINALIS L.

Macrosporium Saponariæ.

SASSAFRAS SASSAFRAS (L.) Karst. (*S. officinale.*)

Athoſtoma microplacum.

Colletotrichum *sp.*Cytispora Sassafras *sp. nov.*

Discosia Artoceras.

Glonium stellatum.

Hypoxyton Sassafras.

Phyllosticta Sassafras.

Sphæroopsis Sassafras.

Valsa subclypeata.

## SCLERODERMA VULGARE.

Boletus parasiticus.

## SCROPHULARIA MARILANDICA L.

Septoſia Scrophulariæ.

*Smilacina racemosa* see VAGNERA RACEMOSA.SMILAX *sp.*

Mycoporon Smilacis.

## SMILAX GLAUCA Walt.

Cercospora smilacina.

Puccinia Smilacis (*Uredo.*)

## SMILAX ROTUNDIFOLIA L.

Cercospora Smilacis.

Discosia maculicola.

Phoma smilacina.

Phyllosticta Smilacis subeffusa.

## SOLANUM TUBEROSUM L.

Phytophthora infestans.

## SOLIDAGO CANADENSIS L.

Coleosporium Solidaginis.

## SPERGULA ARVENSIS L.

Peronospora obovata.

## SPHÆROOPSIS ASIMINÆ.

Macrosporium olivaceum.

STEREUM *sp.*

Hypomyces aurantinus.

TARAXACUM TARAXACUM(L.) Karst. (*Taraxacum officinale* Web.)

Puccinia Hieracii.

Ramularia Taraxaci.

- TECOMA RADICANS (L.) DC.  
 Physalospora *sp.*  
 Septoria Tecomæ.
- THALICTRUM POLYGAMUM Muhl.  
 Leptostroma vulgare.
- TIARELLA CORDIFOLIA L.  
 Puccinia Heuchera.
- TILIA HETEROPHYLLA Vent.  
 Hercospora Tiliæ:
- TILIA AMERICANA L.  
 Phyllosticta Tiliæ.
- TRAUTVETTERIA CAROLINENSIS (Walt.) Vail. (*T. palamata.*)  
 Botrytis vulgaris.  
 Pericornia *sp.*  
 Sarcinella heterospora.  
 Septoria Trautvetteriæ.  
 Vermicularia *sp.*
- TRIFOLIUM PRATENSE L.  
 Uromyces Trifolii.
- TRIFOLIUM REPENS L.  
 Polythrincium Trifolii.
- TRIOSTEUM PERFOLIATUM L.  
 Cladosporium Triostei.
- TRITICUM VULGARE L.  
 Fusarium culmorum.  
 Hymenula cerealis.  
 Puccinia Rubigo-vera.  
 Tilletia Tritici.  
 Ustilago segetum.
- TSUGA CANADENSIS Carr.  
 Coryneum cupulatum.  
 Corticium albo-flavescens.  
 Corticium ochraceum.  
 Cytispora Curreyi.  
 Dacryomyces stellatus.  
 Dacryomyces corticioides.  
 Fomes lucidus.  
 Fomes pinicola.  
 Lenzites abietina.  
 Lenzites sepiaria *forma.*  
 Leocarpus fragilis.  
 Lindbaldia effusa.  
 Merulius Molluscus.  
 Microcera erumpens.  
 Odontia farinaceum.

Otidea onotica ochracea.  
 Panus dorsalis.  
 Peridermium Peckii.  
 Polystictus abietinus.  
 Sphærella conicola  
 Sporodesmium *sp.*  
 Stereum sangiunolentum.  
 Tapesia fusca.  
 Tremellodon gelatinosum.  
 Valsa Abietis.

ULMUS PUBESCENS Walt. (*U. fulva.*)

Corticium incarnatum.  
 Dothidella Ulmiæ.  
 Mucronoporus Everhartii.  
 Nectria vulpinia?  
 Phyllosticta Ulmi?

UREDO (*Cæoma*) NITENS.

Tuberculina persicina.

VAGNERA RACEMOSA (L.) Morong. (*Smilacina racemosa.*)

Septoria Smilaciniæ.  
 Vermicularia liliacearum.

VERBENA URTICÆFOLIA L.

Septoria Verbenæ.

VERNONIA NOVEBORACENSIS (L.) Willd.

Cercospora oculata.  
 Cercospora Vernoniæ.  
 Coleosporium Sonchi.  
 Coleosporium Vernoniæ.  
 Ophiobolus porphyrogonus.  
 Ophiobolus fulgidus.  
 Phlyctæna vagabunda.

VIOLA BLANDA Willd.

Puccinia Violæ (*Uredo.*)

VIOLA HASTATA Michx.

Puccinia Violæ (*Æcid.*)

VIOLA OBLIQUA Hill. (*V. cucullata.*)

Cercospora granuliformis.  
 Cercospora Violæ.

VIOLA PRIMULÆFOLIA L.

Septoria hyalina.

VIOLA ROTUNDIFOLIA Michx.

Puccinia Violæ (*Uredo.*)

VIOLA SAGITTATA Ait.

Septoria Violæ.

VITIS *sp.*

- Asterostoma corticolum.
- Botryosphæria Quercuum.
- Corticium Petersii.
- Diplodia viticola.
- Phialea virgultorum.
- Valsa Vitis.

VITIS ÆSTIVALIS Michx.

- Phyllosticta Labruscæ.

VITIS CORDIFOLIA Michx.

- Phyllosticta Labruscæ.
- Saccidium Vitis *sp. nov.*
- Uncinula Americana.

VITIS *cult.* "Concord."

- Dermatella viticola.
- Lophidium nitidum.
- Phoma uvicola.
- Phyllosticta Labruscæ.
- Uncinula Ampelopsidis.

VITIS LABRUSCA L.

- Plasmopara viticola.

VITIS RUPESTRIS Scheele.

- Phyllosticta Labruscæ.
- Trematosphæria vitigena *sp. nov.*

WEEDS (*in shade.*)

- Arcyria cinerea.

XANTHIUM CANADENSE Mill.

- Botrytis vulgaris.
- Erysiphe Cichoracearum.

XANTHORRHIZA APIIFOLIA L'Her.

- Phyllosticta Xanthorrhizæ *sp. nov.*
- Vermicularia Dematium.

XOLISMA LIGUSTRINA (L.) Britton. (*Andromeda ligustrina.*)

- Rhytisma decolorans.

YUCCA FILAMENTOSA L.

- Coniothyrum concentricum.
- Macrosporium caudatum.
- Vermicularia subeffigurata scapincola.

ZEA MAYS L.

- Colletotrichum lineola.
- Coniophora *sp.*
- Diplodia Maydis.
- Helminthosporium folliculatum.
- Leptosphæria orthogramma.

Macrosporium Maydis.

Ophiobolus acuminatus.

Puccinia Sorghi (*Teleut.*)

Ustilago Maydis.

ZIZIA CORDATA (Walt.) DC.

Oidium erysiphoides.



## Local Plant Names.\*

BEE-WEED	<i>Aster Lowrieanus</i>
BLUE DEVIL	<i>Echium vulgare</i>
BLUE-STEM	<i>Echium vulgare</i>
BLUE THISTLE	<i>Echium vulgare</i>
BLUE-WEED	<i>Echium vulgare</i>
BLISTER PINE	<i>Abies balsamea</i>
BROOM SEDGE	<i>Andropogon scoparius</i>
BUCK-HORN PLANTAIN	<i>Plantago lanceolata</i>
BULL'S-EYE	<i>Chrysanthemum Leucanthemum</i>
CATAWBA	<i>Catalpa Catalpa</i>
COLIC-ROOT	<i>Asarum Canadense</i>
COON-ROOT	<i>Sanguinaria Canadensis</i>
CROWD-WEED	<i>Brassica sinapistrum. Lepidium campestre</i>
CUCKLE-BURR	<i>Xanthium Canadense</i>
CUT-PAPER	<i>Papirus papyrifera</i>
DEER'S-TONGUE	<i>Panicum clandestinum</i>
DEVIL'S FLAX	<i>Linaria vulgaris</i>
DEVIL'S GRANDMOTHER	<i>Elephantopus tomentosus</i>
DEVIL'S GRASS	<i>Chondrilla juncea</i>
DEVIL'S IRON-WEED	<i>Lactuca pulchella</i>
DEVIL'S PLAGUE	<i>Daucus Carota</i>
DEVIL'S-WEED	<i>Lactuca Canadensis</i>
DOG BURR	<i>Cynoglossum officinale</i>
EVE'S THREAD	<i>Hemerocallis fulva</i>
FALL ASTER	<i>Aster Lowrieanus</i>
FAREWELL SUMMER	<i>Aster lateriflorus hirsuticaulis</i>
FEATHER GRASS	<i>Holcus lanatus</i>
FEDERAL-WEED	<i>Xanthium spinosum</i>
FIRE CHERRY	<i>Prunus Pennsylvanica</i>
GIPSY-WEED	<i>Veronica officinalis</i>
GLADE LILY	<i>Lilium Philadelphicum</i>

\*The significations of most of these names are given in the "Preliminary Flora," or in "Weeds of West Virginia."

GLENN PEPPER	<i>Lepidium campestre</i>
GLENN-WEED	<i>Lepidium campestre</i>
HIVE VINE	<i>Meibomia rotundifolia</i>
HOG BITE	<i>Chondrilla juncea</i>
HUTTON-WEED	<i>Dipsacus sylvestris</i>
IMPUDENT LAWYER	<i>Linaria vulgaris</i>
INDICATOR	<i>Botrychium Virginianum</i>
KILL COW	<i>Eleocharis tenuis</i>
KRAUT-WEED	<i>Brassica sinapistrum</i>
NAIL ROD	<i>Aster lateriflorus hirsuticaulis</i>
NAKED-WEED	<i>Chondrilla juncea</i>
NIGGER HEAD	<i>Rudbeckia hirta</i>
OLD FIELD SWEET	<i>Aster lateriflorus hirsuticaulis</i>
OLD VIRGINIA STICK-WEED	<i>Aster lateriflorus hirsuticaulis</i>
OLD WHITE-TOP	<i>Holcus lanatus</i>
PINK BLOOM	<i>Sabbatia angularis</i>
POPPLE	<i>Populus tremuloides</i>
POVERTY GRASS	<i>Eleocharis tenuis. Juncus tenuis</i>
QUEEN-WEED	<i>Pastinaca sativa</i>
QUILL-WORT	<i>Eupatorium purpureum</i>
RAMPS	<i>Allium tricoccum</i>
RIPPLE	<i>Plantago lanceolata</i>
RADICAL-WEED	<i>Solanum Carolinense</i>
RED-WEED	<i>Rumex acetosella</i>
RHEUMATISM-WEED	<i>Apocynum Canadense</i>
SEEDY BUCKBERRY	<i>Xolisma ligustrina</i>
SEEDY DEERBERRY	<i>Vaccinium corymbosum</i>
SHERIFF PINK	<i>Chrysanthemum Leucanthemum</i>
SHITTIM WOOD	<i>Mohrodendron Carolinum</i>
SHOO FLY	<i>Baptisia tinctoria</i>
SINK-FIELD	<i>Potentilla Canadensis</i>
SKELETON-WEED	<i>Chondrilla juncea</i>
SPRUCE PINE	<i>Picea Mariana</i>
STICK-WEED	<i>Aster Lowrieanus</i>
ST. JOHN	<i>Hypericum perforatum</i>
TAR-WEED	<i>Parsonsia petiolata</i>
TOBACCO-WEED	<i>Elephantopus tomentosus</i>
WALLINK	<i>Veronica Americana</i>
WATER THISTLE	<i>Dipsacus sylvestris</i>
WHITE SPRUCE	<i>Picea Mariana</i>
WHITE TOP	<i>Erigeron annuus</i>
WILD BEET	<i>Oenothera fruticosa</i>

WILD COTTON	<i>Apocynum cannabinum</i>
WILD SWEET POTATO	<i>Ipomœa pandurata</i>
WIRE-WEED	<i>Aster lateriflorus hirsuticaulis</i>
WOOL-MAT	<i>Cynoglossum Virginicum</i>
YELLOW DAISY	<i>Rudbeckia hirta</i>
YELLOW LIN	<i>Magnolia acuminata</i>
YELLOW TIP	<i>Solidago juncea</i>
YEW PINE	<i>Picea Mariana</i>

---

# Index.

	PAGE.		PAGE.
Abies .....	192	AMARYLLIDACEÆ.....	199
Abronia .....	211	Amblystegium.....	188
Abutilon.....	217	Ambrosia .....	231
Acalypha.....	215	Ampelopsis.....	217
ACANTHACEÆ .....	227	Amphicarpæa.....	213
Acanthostigma .....	142	Amphisphæria .....	141
Acer.....	216	ANACARDIACEÆ.....	215
Acerates .....	223	Anagallis.....	222
Achillea.....	232	<i>Anaphalis</i> .....	231
Achnanthes.....	85	<i>Andromeda</i> .....	221
Achroanthos.....	199	Andropogon.....	193
Aconitum.....	205	Anemone.....	205
Acorus.....	197	Aneura .....	184
Actæa.....	205	Angelica.....	219
Actinonema.....	116	Anomodon.....	188
Adiantum.....	190	ANONACEÆ .....	204
Adicea .....	202	Antennaria.....	231
Adlumia .....	206	Anthemis.....	232
Adopogon.....	233	Anthostoma .....	135
Æcidium.....	130	Anychia.....	204
Æsculus.....	216	<i>Aphyllon</i> .....	227
AGARICACEÆ .....	156	Apios.....	213
Agaricus.....	160	Aplectrum .....	199
<i>Agastache</i> .....	225	APOCYNACEÆ.....	222
Agrimonia.....	211	Apocynum.....	222
Agrostemma.....	204	Aponectria .....	144
Agrostis.....	194	Aposphæria.....	108
Agyrium.....	182	AQUIFOLIACEÆ.....	215
Ailanthus.....	214	Aquilegia.....	205
AIZOACEÆ.....	203	Aralia.....	219
Aletris.....	199	ARALIACEÆ.....	219
Alisma.....	192	Arabis.....	207
ALISMACEÆ .....	192	ARACEÆ .....	197
Allium.....	198	Archilejunea .....	183
Alnus.....	201	Arctium.....	233
Alsine.....	204	Arcyria .....	86
Althæa.....	217	Arisæma.....	197
Amalanchier .....	211	Aristida .....	193
Amanita .....	156	Aristolochia .....	202
Amanitopsis.....	157	ARISTOLOCHIACEÆ.....	202
AMARANTHACEÆ .....	203	Arrhenatherum .....	194
Amaranthus .....	203	Arthonia.....	181

	PAGE.
Aruncus.....	209
Asarum.....	202
Aschersonia.....	120
ASCLEPIADACEÆ.....	223
Asclepias.....	223
ASCOBOLACEÆ.....	152
Ascobolus.....	152
Ascochyta.....	116
Ascophanus.....	152
Ascyrum.....	217
Asimina.....	204
Asparagus.....	108
Aspergillus.....	88
<i>Aspidium</i> .....	190
Asplenium.....	190
Aster.....	230
Asterina.....	132
Asterostoma.....	171
Astilbe.....	208
Astragalus.....	212
Astragene.....	205
Astrophyllum.....	185
Azalea.....	221
Baccillus.....	83
Bactridium.....	100
Bæomyces.....	180
BALSAMINACEÆ.....	216
Baptisia.....	212
Barbarea.....	207
Barbula.....	186
Bartramia.....	185
BARTRAMIACEÆ.....	185
Bazzania.....	183
Bellincinia.....	183
Belonidium.....	152
Benzoin.....	206
BERBERIDACEÆ.....	206
Berberis.....	206
Betula.....	201
BETULACEÆ.....	201
Biatora.....	180
Bicuculla.....	206
Bidens.....	232
Bignonia.....	227
BIGNONIACEÆ.....	227
Blepharostoma.....	183
Blephila.....	225
Boehmeria.....	202
Boletus.....	161

	PAGE.
Bombardia.....	135
BORAGINACEÆ.....	224
Botrychium.....	190
Botryodiplodia.....	116
Botryosphæria.....	138
Botryosporium.....	88
Botrytis.....	88
Bovista.....	81
Bovistella.....	82
Brachychæta.....	230
Brachyelytrum.....	194
Brassica.....	207
Brauneria.....	231
Bremia.....	83
Bromus.....	195
Broomella.....	145
<i>Broussonetia</i> .....	202
<i>Brunella</i> .....	225
BRYACEÆ.....	186
Bryum.....	186
Buechnera.....	226
Buellia.....	181
Bulgaria.....	154
BULGARIACEÆ.....	153
Bursa.....	207
Buxbaumia.....	185
Cacalia.....	233
CACTACEÆ.....	218
Calamagrostis.....	194
CALICIACEÆ.....	182-155
Calicium.....	182-155
CALLITRICHACEÆ.....	215
Callitriche.....	215
Calocera.....	172
<i>Calopogon</i> .....	200
Calospora.....	142
Caltha.....	205
CALYCANTHACEÆ.....	206
Calycanthus.....	206
Camarosporium.....	117
Camassia.....	198
Camelina.....	207
Campanula.....	229
CAMPANULACEÆ.....	229
Camptosorus.....	190
Campylium.....	189
Cannabis.....	201
Cantharellus.....	158
Capnoides.....	206

	PAGE.		PAGE.
CAPPARIDACEÆ.....	207	<i>Cissus</i> .....	217
CAPRIFOLIACEÆ.....	228	CISTACEÆ.....	217
Cardamine.....	207	Citrullus.....	228
Carduus.....	233	Cladonia.....	179
Carex.....	196	CLADONIACEÆ.....	179
Carpinus.....	201	Cladosporium.....	91
CARYOPHYLLACEÆ.....	204	Clasterosporium.....	92
Cassia.....	213	Clavaria.....	172
Castalia.....	204	CLAVARIACEÆ.....	172
Castanea.....	201	Claytonia.....	203
Castilleja.....	227	Clematis.....	205
Catalpa.....	227	Cleome.....	207
Caulophyllum.....	206	Clethra.....	220
Ceanothus.....	216	CLETHRACEÆ.....	220
<i>Cedronella</i> .....	225	Climacium.....	189
CELASTRACEÆ.....	215	Clinopodium.....	225
Celastrus.....	215	Clintonia.....	198
Celtis.....	201	Clitocybe.....	157
Cenangella.....	153	Clitopilus.....	160
Cenchrus.....	193	<i>Cnicus</i> .....	233
Cephalanthus.....	228	Coleosporium.....	129
Cephalothecium.....	89	Collema.....	177
Cephalozia.....	183	COLLEMEÆ.....	177
Cerastium.....	204	Colletotrichum.....	124
Ceratium.....	98	Collinsia.....	226
Ceratodon.....	188	Collinsonia.....	224
Cercis.....	213	Collybia.....	157
Cercospora.....	93	Comatricha.....	86
Cetraria.....	175	Commelina.....	197
Chærophyllum.....	220	COMMELINACEÆ.....	197
Chamælorium.....	198	COMPOSITÆ.....	229
Chamæraphis.....	193	CONIFERACEÆ.....	192
Cheilanthes.....	190	Coniophora.....	172
Chelidonium.....	206	Coniosporium.....	90
Chelone.....	226	Coniothyrium.....	114
CHENOPODIACEÆ.....	203	Conocyphalum.....	184
Chenopodium.....	203	Conopholis.....	227
Chlorosplenium.....	150	CONVOLVULACEÆ.....	223
Chimaphila.....	220	Convolvulus.....	223
Chiogenes.....	221	Coprinus.....	161
Chionanthus.....	222	Corallorhiza.....	199
Chrosperma.....	199	Cordyceps.....	145
Chrysanthemum.....	232	Coreopsis.....	232
Chrysogonum.....	231	CORNACEÆ.....	220
Chrysopsis.....	229	Cornus.....	220
Cichorium.....	233	Corticium.....	170
Cicuta.....	220	<i>Corycarpus</i> .....	194
Cimicifuga.....	205	Corylus.....	201
Cinna.....	194	Coryne.....	154
Circeæ.....	219	Coryneum.....	125

	PAGE.
Cracca.....	212
CRASSULACEÆ.....	207
Crategus.....	211
Cribraria.....	86
Crucibulum.....	81
CRUCIFERACEÆ.....	206
Cryptospora.....	144
Cryptosporella.....	138
Cucumis.....	228
Cucurbita.....	228
CUCURBITACEÆ.....	228
Cucurbitaria.....	143
Cunila.....	224
Cuphea.....	218
Cuscuta.....	223
Cyathicula.....	150
Cylindrocolla.....	99
Cylindrosporium.....	124
Cylindrothecium.....	189
Cymbella.....	84
Cynoctonum.....	222
Cynoglossum.....	224
CYPERACEÆ.....	195
Cyperus.....	195
Cypripedium.....	200
Cystopteris.....	191
Cystopus.....	82
Cytispora.....	11
Dacryomyces.....	173
Dactylis.....	194
Dactylium.....	89
Dactyloctenium.....	194
Dædalia.....	165
Daldinia.....	137
Dalibarda.....	211
Danthonia.....	194
Dasyscypha.....	151
Dasystema.....	227
Datura.....	226
Daucus.....	219
Deeringia.....	220
Delphinium.....	205
DEMATIACEÆ.....	90
Dendroochium.....	98
Dendrophoma.....	109
Dentaria.....	207
Dermatea.....	153
DERMATEACEÆ.....	152
Dermatella.....	152

	PAGE.
<i>Desmodium</i> .....	213
Diachæa.....	85
Dianthera.....	227
Dianthus.....	204
DIAPENSIACEÆ.....	221
Diaporthe.....	139
DIATOMACEÆ.....	84
Diatrype.....	135
Diatrypella.....	135
Dicksonia.....	191
Dicranella.....	188
DICRANIACEÆ.....	186
Dicranodontium.....	186
Dicranum.....	187
Didymella.....	139
Didymium.....	85
Diervilla.....	228
Dimerosporium.....	132
Dinemasporium.....	122
Diodia.....	228
Dioscorea.....	199
DIOSCOREACEÆ.....	199
Diospyros.....	222
Diphyscium.....	185
Diplodia.....	115
Diplophyllum.....	183
Dipsacus.....	228
DIPSACEÆ.....	228
Dirca.....	218
Discosia.....	121
Disporum.....	198
Ditrichum.....	188
Dodecatheon.....	221
DOTHIDEACEÆ.....	146
Dothidella.....	146
Dothiorella.....	110
Draba.....	207
Drosera.....	207
DROSERACEÆ.....	207
Dryopteris.....	190
Dulichium.....	195
Durella.....	155
Eatonia.....	194
EBENACEÆ.....	222
<i>Echinacea</i> .....	231
<i>Echinosperrum</i> .....	224
Echium.....	224
Eclipta.....	231
Eleocharis.....	195

	PAGE.		PAGE.
Elephantopus.....	229	FISSIDENTACEÆ.....	185
Eleusine.....	194	Fistulina.....	162
<i>Elodea</i> .....	192	Flørkea.....	215
Elymus.....	195	Fomes.....	163
Empusa.....	83	Fragaria.....	211
Endocarpon.....	182	Fraxinus.....	222
Entodon.....	189	Frullania.....	183
ENTOMOPHTHORACEÆ.....	83	Fumaria.....	206
Entomosporium.....	122	Funaria.....	186
Epicoccum.....	101	FUNARIACEÆ.....	186
Epidochium.....	101	Fusarium.....	100
Epigæa.....	221	Fusicladium.....	91
Epilobium.....	218	Fusicoccum.....	111
Epiphegus.....	227		
EQUISITACEÆ.....	191	Galax.....	221
Equisetum.....	191	Galeopsis.....	225
Eragrostis.....	194	Galium.....	228
Erechtites.....	233	Gallinsoga.....	232
ERICACEÆ.....	221	Gaultheria.....	221
Erigenia.....	220	Gaura.....	219
Erigeron.....	231	Gaylussacia.....	221
Erigonum.....	202	Geaster.....	81
Erinella.....	152	Gelatinosporum.....	120
Eriophorum.....	196	Gentiana.....	222
Eryngium.....	215	GENTIANACEÆ.....	222
Erysimum.....	207	Geocalyx.....	183
Erysiptie.....	131	Geopyxis.....	148
Erythronium.....	198	Georgia.....	185
Euonymus.....	215	GEORGIACEÆ.....	185
Eupatorium.....	229	GERANIACEÆ.....	213
Euphorbia.....	214	Geranium.....	213
EUPHORBIACEÆ.....	214	Gerardia.....	227
Eurotium.....	131	Geum.....	211
Euthamia.....	230	Gibberella.....	145
Eutypa.....	134	<i>Gillenia</i> .....	209
Eutypella.....	134	Glaucium.....	206
EXCIPULACEÆ.....	122	Glechoma.....	225
Exidia.....	173	Gleditschia.....	213
EXOASCUS.....	155	Glœosporium.....	122
Exobasidium.....	172	Glœoporus.....	166
		Glonium.....	155
FAGACEÆ.....	201	Gnaphalium.....	231
Fagopyrum.....	203	Gnomoniella.....	138
Fagus.....	201	Gonatobotryum.....	90
Falcata.....	213	<i>Goodyera</i> .....	199
Favolus.....	165	GRAMINACEÆ.....	193
Festuca.....	195	Grandinia.....	168
FILICINÆ.....	190	GRAPHIDACEÆ.....	181
Fimbristylis.....	195	Graphis.....	181
Fissidens.....	185	Gratiola.....	226



	PAGE.
GRIMMIACEÆ.....	188
Guepinia.....	174
GYMNOASCACEÆ.....	155
Gymnocladus.....	213
Gymnosporangium.....	129
Gyromitra.....	147
Gyrostachys.....	199
Habenaria.....	200
Hainesia.....	122
<i>Halesia</i> .....	222
HAMAMELIDACEÆ.....	208
Hamamelis.....	208
Haplosporella.....	114
Harpanthus.....	184
Hedeoma.....	225
Hedwegia.....	189
Helenium.....	232
Helianthemum.....	217
Helianthus.....	232
Heliopsis.....	231
Heliscus.....	100
Helleborus.....	205
Helminthosporium.....	92
Helotiella.....	151
Helotium.....	149
Helvella.....	147
HELVELLACEÆ.....	147
Hemiarcyria.....	87
Hemorocallis.....	198
Hendersonia.....	116
Hepatica.....	205
Heracleum.....	220
Herbertia.....	183
Hercospora.....	139
Herpotrichia.....	142
Hesperis.....	207
Heuchera.....	208
Hibiscus.....	217
Hicoria.....	200
Hieracium.....	233
HIPPOCASTANACEÆ.....	216
Hirneola.....	173
Holcus.....	194
Holwaya.....	154
Homalocenchrus.....	193
Horomyces.....	174
Houstonia.....	227
Humaria.....	148
Humulus.....	201

	PAGE.
HYDNACEÆ.....	167
Hydnum.....	167
Hydrangea.....	208
Hydrastis.....	206
HYDROCHARITACEÆ.....	192
Hydrocotyle.....	219
HYDROPHYLLACEÆ.....	223
Hydrophyllum.....	223
Hygrophorus.....	158
Hylocomium.....	188
Hymenochæte.....	169
Hymenula.....	99
HYPERICACEÆ.....	217
Hypericum.....	217
Hyphelia.....	102
Hyphoderma.....	80
Hypholoma.....	160
HYPHOMYCETACEÆ.....	87
HYPNACEÆ.....	188
Hypnum.....	188
Hypocrea.....	144
HYPOCREACEÆ.....	144
Hypocreopsis.....	145
Hypoderma.....	156
Hypomyces.....	144
Hypopytis.....	221
Hypoxylon.....	136
HYSTERIACEÆ.....	155
Hysterium.....	155
Hysterix.....	195
Hysterographium.....	156
Ilex.....	215
Illosporium.....	99
Ilysanthes.....	226
Impatiens.....	216
Inula.....	231
Ipomœa.....	223
IRIDACEÆ.....	199
Iris.....	199
Irpex.....	167
Isanthus.....	224
Isaria.....	97
Isopyrum.....	205
Ithyphallus.....	81
Jeffersonia.....	206
Jubula.....	183
JUGLANDACEÆ.....	200
Juglans.....	200

	PAGE.		PAGE.
JUNCACEÆ .....	198	Leptosphæria .....	141
Juncoides .....	198	Leptostromella .....	122
Juncus .....	198	Leptothyrium .....	121
Jungermannia .....	184	LEPTROSTOMACEÆ .....	121
JUNGERMANNIACEÆ .....	183	Lespedeza .....	213
Juniperus .....	192	Leucobryum .....	186
		Leucodon .....	189
Kalmia .....	221	Libertella .....	125
Kantia .....	183	Ligusticum .....	220
Kneiffia .....	168	Ligustrum .....	222
Kœllia .....	224	LILIACEÆ .....	198
Korycarpus .....	194	Lilium .....	198
Kyllingia .....	195	LIMNANTHACEÆ .....	215
		Limodorum .....	200
LABIATEÆ .....	224	LINACEÆ .....	214
Lachnea .....	148	Linaria .....	226
Lachnella .....	151	Lindbaldia .....	86
Lachnocladium .....	173	Linum .....	214
Lacinaria .....	229	Lippia .....	224
Lactarius .....	158	Liquidambar .....	208
Lactuca .....	233	Liriodendron .....	204
Lamium .....	225	Lithospermum .....	224
Lamproderma .....	85	Lobelia .....	229
Laportea .....	202	LOGANIACEÆ .....	222
Lappula .....	224	Lolium .....	195
Lasiosphæria .....	142	Lonicera .....	228
Lathyrus .....	213	Lophidium .....	146
LAURACEÆ .....	206	LOPHIOSTOMACEÆ .....	146
Lecanora .....	178	Lophiotrema .....	146
LECANOREÆ .....	178	Lophodermium .....	156
Lecanidion .....	155	LORANTHACEÆ .....	202
Lechea .....	217	Ludwegia .....	218
Lecidea .....	181	Lupinus .....	212
LECIDEÆ .....	180	<i>Luzula</i> .....	198
Leersia .....	186	Lycium .....	226
Legouzia .....	229	Lycogala .....	86
LEGUMINOSÆ .....	212	LYCOPERDACEÆ .....	81
<i>Lejunea</i> .....	183	Lycoperdon .....	82
Lemna .....	197	LYCOPODIACEÆ .....	191
LEMNACEÆ .....	197	Lycopodium .....	191
Lentinus .....	159	Lycopus .....	224
Lenzites .....	160	Lysimachia .....	222
Leocarpus .....	85	LYTHRACEÆ .....	218
Leonurus .....	225		
Leotia .....	153	<i>Maclura</i> .....	202
Lepidium .....	207	Macrosporium .....	95
Leptandra .....	226	Magnolia .....	204
Leptobryum .....	186	MAGNOLIACEÆ .....	204
Leptogium .....	177	<i>Maianthemum</i> .....	198
Leptorchis .....	199	Malva .....	217

	PAGE.
MALVACEÆ	217
Marasmius	159
Marchantia	184
MARCHANTIACEÆ	184
Marrubium	225
Marsonia	125
Marsupelia	184
Massaria	141
Matricaria	232
Medicago	212
Mediola	199
Meehania	225
Meibomia	213
Melampsora	127
Melampyrum	227
MELANCONIACEÆ	122
Melanconis	139
Melanconium	125
Melanomma	142
Melanthium	199
MELASTOMACEÆ	218
Melica	194
Melilotus	212
Melissa	225
MENISPERMACEÆ	206
Menispermum	206
Mentha	224
Menziezia	221
Mertensia	224
Merulius	166
Metzgeria	184
Micrampelis	228
Microcera	101
Micrococcus	84
Microsphæria	131
MICROTHYRIACEÆ	146
Mimulus	226
Mitchella	228
Mitella	208
Mitremyces	81
Mitrula	147
MNIACEÆ	185
Mohrodendron	222
Mollia	186
Mollisia	150
Mollugo	203
Monarda	225
Moneses	220
Monilia	87
Monotropa	221

	PAGE.
MONOTROPACEÆ	221
MORACEÆ	201
Morchella	147
Morus	201
Mucor	82
Mucronoporus	164
Muhlenbergia	194
Muscari	198
Mycena	157
Mycoporon	146
Myosotis	224
Myriadoporus	165
Myriangium	179
MYXOMYCETEÆ	85
Myxosporium	123
Næmatella	173
NAIDACEÆ	192
<i>Nasturtium</i>	206
Naumbergia	222
Navicula	84
NECKERACEÆ	189
Neckeria	189
Nectria	144
NECTROIDACEÆ	120
<i>Negundo</i>	216
Nepeta	225
NIDULARIACEÆ	81
Nitzschia	85
Nummularia	138
Nyctalis	159
NYMPHACEÆ	204
Nymphæa	204
Nyssa	220
Obolaria	222
Odontia	168
Odontochisma	183
Oenothera	219
Oidium	87
OLEACEÆ	222
Omphalia	158
Onagra	219
ONAGRACEÆ	218
Onoclea	191
Onosmodium	224
Oospora	87
Opegrapha	181
Ophiobolus	143
OPHIOGLOSSACEÆ	190

	PAGE.		PAGE.
Ophioglossum.....	190	Peranium.....	199
Opulaster.....	208	Periconia.....	90
Opuntia.....	218	Peridermium.....	130
Orbilia.....	153	Perilla.....	224
ORCHIDACEÆ.....	199	PERISPORIACEÆ.....	130
Orchis.....	200	Peronospora.....	83
Orinthogallum.....	198	PERONOSPORACEÆ.....	82
OROBANCHACEÆ.....	227	Pertusaria.....	179
Orthotrichum.....	188	Pestalozzia.....	125
Osmorrhiza.....	220	Petunia.....	226
Osmunda.....	191	Peziza.....	148
Ostropa.....	156	PEZIZACEÆ.....	148
Ostrya.....	200	Phacelia.....	224
Otidea.....	148	PHACIDIACEÆ.....	154
OXALIDACEÆ.....	214	Phalaris.....	193
Oxalis.....	214	PHALLACEÆ.....	81
Oxycoccus.....	221	Phaseolus.....	213
Oxydendron.....	221	Phegopteris.....	190
Oxypolis.....	220	Phialea.....	149
Panax.....	219	Phlebia.....	168
Panæolus.....	161	Phleum.....	194
Panicularia.....	195	Phlox.....	223
Panicum.....	193	Phlyctæna.....	120
Pannaria.....	177	Phoma.....	106
PANNARIEÆ.....	177	Phoradendron.....	202
Panus.....	159	Phragmidium.....	129
Papaver.....	206	Phryma.....	224
PAPAVERACEÆ.....	206	Phyllachora.....	146
Papyrius.....	202	Phyllactinia.....	130
Parmelia.....	176	Phyllosticta.....	102
PARMELIEÆ.....	176	Physalis.....	226
Parnassia.....	208	Physalodes.....	226
Paronychia.....	204	Physalospora.....	138
Parthenium.....	231	Physarum.....	85
Parthenocissus.....	217	Physcia.....	176
Passiflora.....	218	Physcomitrium.....	186
PASSIFLORACEÆ.....	218	Physostegia.....	225
Paspalum.....	193	Phytolacca.....	203
Pastinaca.....	220	PHYTOLACCACEÆ.....	203
PATELLARIACEÆ.....	155	Phytophthora.....	83
Pedicularis.....	227	Picea.....	192
Pellæa.....	190	Pieris.....	221
Pellia.....	184	Pilacre.....	97
Peltigera.....	177	<i>Pilea</i> .....	202
PELTIGERIEÆ.....	177	Pinus.....	192
Pencillium.....	88	Placodium.....	178
Peniophora.....	171	Plagiochila.....	184
Penthorum.....	207	Plagiothecium.....	189
Pentstemon.....	226	PLANTAGINACEÆ.....	227
		Plantago.....	227

	PAGE.
Plasmopara .....	83
PLATANACEÆ .....	208
Platanus .....	208
Pleonectria .....	145
Pleurotus .....	158
Plowrightia .....	146
Poa .....	194
Podophyllum .....	206
Podospæria .....	130
Pogonia .....	200
Polygala .....	214
POLYGALACEÆ .....	214
POLYGONACEÆ .....	202
Polygonatum .....	198
Polygonum .....	202
Polymnia .....	231
POLEMONIACEÆ .....	223
Polemonium .....	223
Polypodium .....	190
POLYPORACEÆ .....	161
Polyporus .....	162
Polyscytulum .....	87
Polystictus .....	163
Polythrincium .....	91
POLYTRICHIACEÆ .....	185
Polytrichum .....	185
Populus .....	200
<i>Porella</i> .....	183
Poria .....	164
Porothelium .....	166
Porteranthus .....	209
Portulaca .....	203
PORTULACACEÆ .....	203
Potamogeton .....	192
Potentilla .....	211
<i>Poterium</i> .....	211
Prenanthes .....	233
PRIMULACEÆ .....	221
Propolis .....	154
Prunella .....	225
Prunus .....	208
Psathyrella .....	161
Pseudohelotium .....	149
Pseudovalsa .....	142
Ptelea .....	214
Pteris .....	190
Ptilium .....	189
Puccinia .....	128
Pyrenopeziza .....	150
Pyrenophora .....	143

	PAGE.
Pyrenula .....	182
Pylaisia .....	189.
Pyrola .....	220
PYROLACEÆ .....	220
Pyularia .....	202
Pyrus .....	211
Quercus .....	201
Radula .....	183
Radulum .....	167
Ramalina .....	175
Ramularia .....	90
RANUNCULACEÆ .....	205
Ranunculus .....	205
Raphanus .....	207
Reticularia .....	86
Rhabdospora .....	120
Rhamnus .....	216
RHAMNACEÆ .....	216
Rhexia .....	218
Rhizina .....	148
Rhododendron .....	221
Rhus .....	215
Rhynchospora .....	196
Rhytisma .....	154
Ribes .....	208
<i>Ridania</i> .....	232
Rinodia .....	170
Robinia .....	212
Roripa .....	206
Rosa .....	211
ROSACEÆ .....	208
Rosellinia .....	135
RUBIACEÆ .....	227
Rubus .....	209
Rudbeckia .....	231
Ruellia .....	227
Rumex .....	203
Russula .....	158
Ruta .....	214
RUTACEÆ .....	214
Sabbatia .....	222
Saccharomyces .....	84
SACCHAROMYCETACEÆ .....	84
Sacidium .....	121
Sagittaria .....	192
SALICACEÆ .....	200
Salix .....	200

	PAGE.		PAGE.
Salvia.....	225	Sonchus.....	233
Sambucus.....	228	Sorbus.....	211
Samolus.....	222	SPARGANIACEÆ.....	192
Sanguinaria.....	206	Sparganium.....	192
Sanguisorba.....	211	Spartina.....	193
Sanicula.....	220	Spathularia.....	147
SANTALACEÆ.....	202	Spathyema.....	197
Saponaria.....	204	<i>Specularia</i> .....	229
Sarcinella.....	97	Speira.....	95
Sarcoscypha.....	148	Spergula.....	204
Sassafras.....	206	Sphærella.....	138
SAURURACEÆ.....	200	SPHÆRIACEÆ.....	132
Saururus.....	200	Sphærocephalus.....	185
Saxifraga.....	208	Sphærographium.....	120
SAXIFRAGACEÆ.....	208	SPHÆROIDACEÆ.....	102
Scapania.....	183	Sphæronæma.....	109
SCHIZOMYCETACEÆ.....	83	SPHÆROPSIDEÆ.....	102
Schizophyllum.....	160	Sphæropsis.....	113
Scirpus.....	196	Sphærotheca.....	130
Scleroderma.....	82	SPHAGNACEÆ.....	189
Scleroderris.....	153	Sphagnum.....	189
Sclerotina.....	149	Spirillum.....	84
Scorias.....	132	Sporocybe.....	98
Scoriomyces.....	100	Sporodinia.....	82
Scrophularia.....	226	Sporodesmium.....	95
SCROPHULARIACEÆ.....	226	Spiræa.....	208
Scutellaria.....	225	Stachyidium.....	91
Sedum.....	207	Stachys.....	225
Selaginella.....	191	Stagonospora.....	116
SELAGINELLACEÆ.....	191	STAPHYLACEÆ.....	215
Senecio.....	232	Staphylea.....	215
Septoria.....	117	Stauroneis.....	84
Septosporium.....	96	Steganosporium.....	126
Sericocarpus.....	230	Steironema.....	222
Sicyos.....	228	<i>Stellaria</i> .....	204
Sida.....	217	Stenophragma.....	207
Sieglingia.....	194	Stereodon.....	189
Silene.....	204	STEREODONTACEÆ.....	188
Silphium.....	231	Stereum.....	168
SIMARUBACEÆ.....	214	Sterigmatocystis.....	88
Sirococcus.....	109	Sticta.....	177
Sisymbrium.....	207	STICTACEÆ.....	154
Sisyrinchium.....	199	Stictis.....	154
SMILACEÆ.....	199	STILBACEÆ.....	97
Smilax.....	199	Stremonitis.....	86
SOLANACEÆ.....	225	Streptococcus.....	84
Solanum.....	225	Streptopus.....	198
Solea.....	218	Streptothyx.....	90
Solenia.....	166	Strobilomyces.....	162
Solidago.....	229	Stropharia.....	160

	PAGE.
Stylosanthes.....	212
Stylophorum.....	206
STYRACEÆ.....	222
Symphoricarpos.....	228
Symphytum.....	224
Syndesmon.....	205
Synedra.....	85
Tanacetum.....	232
Tapesia.....	150
Taraxacum.....	233
Taxus.....	192
Tecoma.....	227
Teichospora.....	143
<i>Tephrosia</i> .....	212
Teucrium.....	224
Thalesia.....	227
Thalictrum.....	205
Thaspium.....	220
Thelephora.....	168
THELEPHORACEÆ.....	168
Thelia.....	188
Theloschistes.....	175
Thelotrema.....	179
Therofon.....	208
Thuidium.....	188
Thuja.....	192
THYMELEACEÆ.....	218
Thyrsidium.....	125
Tiarella.....	208
<i>Tiedmannia</i> .....	220
Tilia.....	217
TILIACEÆ.....	217
Tilletia.....	127
Tilmadoche.....	85
Tortula.....	186
TORTULACEÆ.....	186
Torula.....	90
Toxylon.....	202
Tragopogon.....	233
Trametes.....	165
Trautvetteria.....	205
Trematosphæria.....	142
Tremella.....	173
TREMELLINACEÆ.....	173
Tremellodon.....	167
Trichægum.....	96
Trichia.....	86
Trichocolea.....	183
Trichoderma.....	88

	PAGE.
Tricholoma.....	157
Trichopeziza.....	151
Trichosphæria.....	138
Trichostema.....	225
Trichothecium.....	89
Trientalis.....	221
Trifolium.....	212
Trillium.....	199
Trimmatostroma.....	102
Triosteum.....	228
Tripsacum.....	193
Trogia.....	159
Trypethelium.....	182
Tsuga.....	192
Tubercularia.....	98
TUBERCULARIACEÆ.....	98
Tuberculina.....	99
Tubulina.....	86
Tylostoma.....	81
Tympanis.....	153
Typha.....	192
TYPHACEÆ.....	192
Udora.....	192
ULMACEÆ.....	201
Ulmaria.....	209
Ulmus.....	201
Ulocolla.....	173
UMBELLIFERÆ.....	219
Umbilicaria.....	176
UMBILICARIÆ.....	176
Uncinula.....	131
Unifolium.....	198
Uniola.....	194
Uredo.....	130
Urnula.....	152
Urocystis.....	127
Urtica.....	202
URTICACEÆ.....	202
Urticastrum.....	202
USNACEÆ.....	175
Usnea.....	175
USTILAGINACEÆ.....	126
Ustilago.....	126
Ustulina.....	136
Uvularia.....	198
Vaccinium.....	221
Vagnera.....	198
Valeriana.....	228

	PAGE.		PAGE.
VALERIANACEÆ .....	228	VITACEÆ .....	216
Valsa .....	132	Vitis .....	216
Valsaria .....	141	Vleckia .....	225
Veratrum .....	199	Volutella .....	100
Verbascum .....	226	Volvaria .....	160
Verbena .....	224		
VERBENACEÆ .....	224	Waldstenia .....	211
Verbesina .....	232	Weissia .....	188
Vermicularia .....	109	Woodsia .....	191
Vernonia .....	229		
Veronica .....	226	Xanthium .....	231
VERRUCARIACEÆ .....	182	Xanthorrhiza .....	206
Verticillium .....	89	Xolisma .....	221
Vibrissea .....	147	Xylaria .....	136
Viburnum .....	228		
Vicia .....	213	Zanthoxylum .....	214
Vincetoxicum .....	223	Zizia .....	220
Viola .....	217	Zygodesmus .....	90
VIOLACEÆ .....	217		





